

## **Brightest Stars, expanded version**

**by Toomas Karmo ([Toomas.Karmo@gmail.com](mailto:Toomas.Karmo@gmail.com))**

### **PREFACE: Scope and purpose of this essay**

This online essay is an extended version of the essay in the printed-edition Handbook, containing all the material of its printed-edition accompaniment, but adding material of its own. The accompanying online table is likewise an extended version of the printed-edition table, (a) with extra stars (after providing for multiplicity, as we explain below, the brightest MK-classified 325, allowing for variability, where the printed edition has about 30 fewer, allowing for variability: our cutoff is mag.  $\sim 3.55$ ), and (b) with additional remarks for most of the duplicated stars. We use a dagger superscript ( $\dagger$ ) to mark data cells for which the online table supplies some additional information, some context, or a caveat.

The online essay and table try to address the needs of three kinds of serious amateur: amateurs who are also astrophysics students (whether or not enrolled formally at some campus); amateurs who, like many in the RASC, assist in public outreach, through some form of lecturing; and amateurs who are planning their own private citizen-science observing runs, in the spirit of such “pro-am” organizations as AAVSO. Additionally, we would hope that the online project will help serve a constituency of sky-lovers, whether professional or amateur, who work with the heavens in an unambitious and contemplative spirit, seeking to understand at the eyepiece, or even with the naked eye, the realities behind the little that their limited circumstances may allow them to see. (This is the same contemplative exercise as is proposed for the Cyg X-1 black hole, with its gas-dumping supergiant companion HD226868, in the Handbook “Expired Stars” essay: with a small telescope, or even with binoculars, we first find HD226868, and then take a moment to ponder in awe the accompanying unobserved realities of gas-fed hot accretion disk, event horizon, and spacetime singularity.)

Our online project, started as a supplement to the 2017 Handbook, must be considered still in its rather early stages. We cannot claim to have fully satisfied the needs of our various constituencies. Above all, we cannot claim to have covered all the appropriate points from stellar-astronomy news in our “Remarks” column, important though news is to amateurs of all three types. We would hope in coming years to remedy our deficiencies in several ways, most notably by relying more in our writing on recent primary-literature journal articles, with appropriate explicit citations.

In our citations, we favour the now-preferred astrophysics “bibcode” formalism. The formalism is documented in [simbad.u-strasbg.fr/guide/refcode/refcode-paper.html](http://simbad.u-strasbg.fr/guide/refcode/refcode-paper.html), and again in section 1.2.3 (headed “Bibliographic Identifiers”) in [adsabs.harvard.edu/abs\\_doc/help\\_pages/data.html](http://adsabs.harvard.edu/abs_doc/help_pages/data.html).

A bibcode can be transformed into the display of a more human-readable bibliography entry, often with clickable hyperlink to an underlying online full-text, all-illustrations PDF publication, in various ways. We illustrate some possibilities by taking an extreme case, namely our bibcode reference to the classic 1910 Joel Stibbins *Astrophysical Journal* paper that reports the electric-photometry discovery of a secondary minimum in the Algol light curve. Old though the paper is, it is nevertheless available online. The bibcode (as we state again in our “Remarks” for the Algol entry in our table) is [1910ApJ....32..185S](https://ui.adsabs.harvard.edu/abs/1910ApJ....32..185S). A browser display with hyperlink to the desired full-text, all-illustrations PDF is available from the Centre de Données Stellaires (CDS) server (probably in Strasbourg) as [simbad.u-strasbg.fr/simbad/sim-ref?bibcode=1910ApJ....32..185S](http://simbad.u-strasbg.fr/simbad/sim-ref?bibcode=1910ApJ....32..185S). If something has gone wrong—and experience suggests that things can go wrong, even when a bibcode appears to casual inspection to be correctly typed, at any rate in some such autonomous-agent computing environment as Microsoft Office—then one can recover through CDS as [simbad.u-strasbg.fr/simbad/sim-fid](http://simbad.u-strasbg.fr/simbad/sim-fid) if the star of interest and year of publication are known. In this particular case, recovery involves giving [simbad.u-strasbg.fr/simbad/sim-fid](http://simbad.u-strasbg.fr/simbad/sim-fid) some convenient identifier, for instance the IAU-promulgated name “Algol” or the Bayer identifier “beta Per.” In the Algol-specific input form generated, one next asks, in the “References” section of the form, for all references from 1910 to 1910. The duly displayed bibcode, [1910ApJ....32..185S](https://ui.adsabs.harvard.edu/abs/1910ApJ....32..185S), for the sole 1910-through-1910 reference, is shown as a clickable hyperlink. Upon further clicking, the hyperlink eventually yields the PDF. A similar browser display is available from a (probably North American) ADS-NASA server as [ui.adsabs.harvard.edu/abs/1910ApJ....32..185S/abstract](http://ui.adsabs.harvard.edu/abs/1910ApJ....32..185S/abstract). As a fourth possibility, the PDF is

retrievable through a self-evident set of steps that starts by copying and pasting the bibcode into the “Bibliographic Code Query” box at the paper-workflow, as distinct from the more obviously accessible paperless-workflow, online form [ui.adsabs.harvard.edu/paper-form](http://ui.adsabs.harvard.edu/paper-form). This fourth method has the advantage that multiple bibcodes can be entered within a single query. As a fifth possibility, which in our view cannot be guaranteed to work (but there seem to be intermittent problems with the fourth possibility as well; and in general, servers should not be presumed fully reliable, in any discipline) is simply to put [1910ApJ...32..185S](https://ui.adsabs.harvard.edu/abs/1910ApJ...32..185S) naively into a general Google search, and to explore the ensuing chain of hyperlinks: in the case of at least a heavily cited paper, one is likely soon enough to reach an abstract at ADS-NASA or some similar authority, with accompanying PDF.

The bibliographic support of [simbad.u-strasbg.fr/simbad](http://simbad.u-strasbg.fr/simbad) and [ui.adsabs.harvard.edu](http://ui.adsabs.harvard.edu), as the principal tools for our primary-literature searching, is herewith gratefully acknowledged, as are *Wikipedia* (in exact-science topics, generally careful and up to date); *Sky & Telescope*, the web materials of Prof. James Kaler, and at a more technical level, two key sources of data, the Washington Double Star Catalog (WDS) and AAVSO. Helpful at AAVSO are not only the general graphing facility and the general AAVSO record of observations, but also a more recent offering, the VSX online database.

Thanks are additionally due both to the RASC family in North America and to the Tartu Observatory dark-sky campus in Estonia for encouragement and support. At the dark-sky campus, particular mention should be made of conversations regarding photometry, notably with Dr. T. Eenmäe and Dr. I. Kolka.

It has been necessary in the photometry section (“Section 6”) to make an unusually large number of judgement calls or professions of uncertainty. It seems for this reason particularly advisable for the present author, Toomas Karmo, to stress that the various inevitable errors in this work (notably in Section 6, but also in the other sections, and the table) are his sole responsibility, and to stress his desire to receive feedback. Feedback is best communicated in e-mail, to [toomas.karmo@gmail.com](mailto:toomas.karmo@gmail.com), using some such hard-to-miss subject line as “RASC brightest-stars online: some possible errors noted; some recommendations.”

### **SECTION 1: Selection bases for our 317 nominal “bright stars,” strictly 325 MK-classified bright stars**

Of our selected 317 nominal stars, five call for extra comment pertinent to this mag. ~3.55 naked-eye selection criterion. (1)  $\kappa$  (kappa) CMa (at RA ~6h50) brightened in the 1960s or 1970s, just managing to meet the cutoff, and has remained bright. This change was unfortunately not noted in the RASC Handbook until 2019. Should  $\kappa$  CMa now once again fade, we propose to keep listing it for at least a few years, since it is a variable of the  $\gamma$  Cas type (and may therefore be liable to yet further episodes of brightening during the 21st century; in general, the  $\gamma$  Cas variables, whether temporarily bright or temporarily faint, are desirable targets for ongoing, regular, citizen-science spectroscopy, and even naked-eye monitoring, being closely associated with the amateur-relevant “Be phenomenon,” which we discuss at the end of Section 5 in this essay). (2) We discontinued listing L<sub>2</sub> Pup (at RA ~07h14) in the 2017, 2018, and 2019 Handbooks. Since then, however, we have reverted to our pre-2017 policy, since L<sub>2</sub> Pup is a semi-regular pulsator, occasionally bright. (3) T CrB A (at RA ~16h00) has shown nova behaviour, brightening from its current very faint state (mag. ~10) to mag. 2.0 in 1866 and to mag. 3.0 in 1946. The Handbook has for years or decades listed this T CrB A in its brightest-stars table. We propose to continue listing it, since its history suggests the possibility of a 21st-century outburst. (4) The Mira-type variable  $\chi$  Cyg A (at RA~19h50) was considered by visual observers to surmount our brightness threshold in 2006, 2008, and 2013, even while not being assessed as quite this bright by CCD observers in those three years. From January 2023 Handbook work onward, we regard  $\chi$  Cyg A as a star (just barely) meriting inclusion. (5) The supergiant  $\mu$  Cep A, “Herschel’s Garnet Star” (at RA~21h44) occasionally surmounts our brightness threshold, being (e.g.) as bright as mag. 3.4 in a V-passband CCD observation from 2022 June 28. From January 2023 Handbook work onward, we therefore regard  $\mu$  Cep A as a star (more than just barely) meriting inclusion.

Two omissions from our selection of 317 nominal stars also call for comment. (1) While mindful of the fact that  $\eta$  Car brightened greatly, attaining even mag. 0 for a few years from 1837 onward, we omit it from our table since there is no firm prognosis of a 21st-century repetition of that outburst. In February 2022,  $\eta$  Car was being reported as at mag. 4.1 in the V band, and in January 2021 was being variously reported visually as at

mag. 4.3 and 4.5, in all cases being decidedly fainter than our naked-eye cutoff. The star was not reported by AAVSO visual observers as any brighter than mag. 3.9 in the second half of 2020. (2) P Cyg temporarily surmounted the Sample S brightness threshold around the year 1600, and perhaps again on one or two occasions later in the 1600s. The entire available AAVSO record, however, reaching back to 1891, shows P Cyg at all times no brighter than mag. 4.0. The AAVSO archive shows that in December 2022, P Cyg was being observed with photometric equipment in the V passband as at mag. 4.7.

We may now explain in what sense the set of 317 is nominal. In a strict accounting, the selection is a set of 317 objects that are in naked-eye terms “bright stars,” i.e. are bright, unresolved point sources of starlight. Three kinds of situation need to be distinguished here, as we move from naked-eye impressions to underlying physical realities:

(1) In some cases (to cite an example at random,  $\beta$  Tau [Elnath]) what is to the naked eye a point source actually is, so far as is known, a solitary star.

(2) Very common is a situation in which a bright star is a component in a multi-star system, with the other member(s) making either a very small or a negligible contribution to the naked-eye retinal signal. An instance of the former type of pairing is  $\gamma$  And A, from which at a distance of just over 9" lie two fainter stars,  $\gamma$  And B and  $\gamma$  And C, themselves separated by a mere 0.2", and so faint that the BC pairing shines at around mag. 5. This has the consequence that BC makes just a modest contribution to the overall  $\gamma$  And ABC naked-eye neuron response. An instance of the latter type of pairing is  $\alpha$  CMa A (Sirius), with  $\alpha$  CMa B a white dwarf shining at mag. 8.5, in other words shining so feebly as to play essentially no role in the signal generated by the naked-eye retina. This binary constitutes a not trivial, and yet also at the present favourable time a not hopeless, project for the small telescope. (At [www.rasc.ca/sirius-observing-challenge](http://www.rasc.ca/sirius-observing-challenge), RASC notes that with apastron due in 2025, “an extremely difficult feat has become merely a very demanding one.”) Since our table is officially a table of bright stars, we take care, at any rate in our various table revisions from early 2021 onward, to write in our first table column “ $\gamma$  And A” (not “ $\gamma$  And AB” or “ $\gamma$  And”), and “ $\alpha$  CMa A” (not “ $\alpha$  CMa AB” or “ $\alpha$  CMa”). Helpfully, the naming rules promulgated since around 2016 at IAU, and reflected in our concluding “Remarks” column, stipulate, in parallel with our first-column decision, that a name such as “Sirius” applies to a star such as  $\alpha$  CMa A, rather than to the binary system  $\alpha$  CMa AB.

(3) In eleven other cases, the naked-eye point, shining at mag.  $\sim 3.55$  or brighter, is the combined light of two binary-system components, each individually bright enough to count as a “bright star”—perhaps with each component exceeding our mag.  $\sim 3.55$  cutoff, but also perhaps with one or both components just a little fainter than our mag.  $\sim 3.55$  cutoff, and yielding a “star” brighter than mag.  $\sim 3.55$  upon combining the light.

These eleven, so-to-speak awkward, cases (awkwardly forcing us to write the binary designations “AB” or “Aa,Ab” in the first column) are the following:

- $\beta$  Phe AB (with each of A, B individually around mag. 4, yielding an aggregated naked-eye impression of mag. 3.2)
- $\gamma$  Per Aa,Ab (with each of Aa, Ab a little brighter than mag. 4, yielding an aggregated naked-eye impression of mag. 2.91)
- $\alpha$  Aur Aa (Capella), Ab (with each of Aa, Ab very close to mag. 0)
- $\beta$  Aur Aa (Menkalinan), Ab (with magnitudes nearly equal, yielding an aggregated naked-eye impression a little brighter than mag. 2)
- $\zeta$  CMa Aa,Ab (magnitudes nearly equal, at 3.6 and 3.8; this very tight binary is not as yet well observed, with as of at any rate 2022 March 2 just 4 WDS-documented measurements, from 2019 and 2020)
- $\gamma$  Vir A (Porrina), B (magnitudes nearly equal, and with each individually very close to our mag.  $\sim 3.55$  cutoff, yielding an aggregated naked-eye impression a little brighter than mag. 3)
- $\beta$  Cen Aa (Hadar), Ab (magnitudes nearly equal, with each individual star much brighter than our mag.  $\sim 3.55$  cutoff)
- $\eta$  Oph A (Sabik), B (with B at mag. 3.5)
- $\lambda$  Sco Aa (Shaula), Ab (with even Ab well above our cutoff, at mag.  $\sim 2.8$ )
- $\zeta$  Sgr A (Ascella), B (with B at mag. 3.5)

- $\pi$  Sgr A (Albaldah), B (a poorly documented pairing, with the faint outlier C also poorly documented: WDS implies that B is of nearly the same magnitude as A, with each of these two stars very close to our mag.  $\sim 3.55$  cutoff)

It is tempting to consider the  $\eta$  Peg system to be a twelfth case, requiring entry as “ $\eta$  Peg Aa (Matar), Ab.” But since Ab is decidedly fainter than our mag.  $\sim 3.55$  cutoff, and Aa only slightly fainter than that, we are obliged instead to enter this case simply as “ $\eta$  Peg Aa (Matar),” drawing attention in the table “Remarks” column to the fact that our stated magnitude of 2.93 is for the combined light. Somewhat like the  $\eta$  Peg system is the o (omicron) Leo system, where o Leo Aa (Subra) is very close to mag. 3.5, and where o Leo Ab is, while fainter than mag. 3.5, nevertheless bright enough to make a non-trivial contribution to the overall visual impression. Before 2021, our table unfortunately had the erroneous information that o Leo Aa,Ab is a binary system in which the components are of equal magnitude. Also somewhat like the  $\eta$  Peg system is  $\theta$  Tau Aa, shining a bit below our magnitude cutoff at 3.74, and rising to just above the cutoff in the combined light of  $\theta$  Tau Aa,Ab (where Ab is well below our cutoff, at mag. 4.86).

We thus have a table of nominally 317 stars, comprising a more refined, i.e. less nominal, analysis  $317 + 11 = 328$  bright stars. With 3 exceptions, each of the 328 has a known (at worst, an uncertainty-flagged) MK temperature type and MK luminosity class (with the Sun, of course, better observed than any of the others). (The case of  $\zeta$  CMa Aa,Ab is admittedly rather indeterminate. Here we have a longstanding MK type, from decades before the 2019 discovery of binarity. Should it be assumed now that the type applies only to  $\zeta$  CMa Aa, or should it be understood as approximately correct also for  $\zeta$  CMa Ab? Since the magnitude difference is small, around 0.2 mag., and since the stars are likely to be of the same age, as products of co-genesis from an ISM molecular cloud, we take the latter option.) The final result is accordingly a set of  $328 - 3 = 325$  bright stars of known MK classification.

## **SECTION 2: General characteristics of our 325 MK-classified bright stars**

Our 325-element sample is found to lie in a region, around 3000 ly in radius, essentially confined to the sandwich-filler, or “thin disk,” part of the overall galactic disk, within the Orion Arm. Of the few Sample-S interlopers born outside the sandwich filling, and now temporarily passing through it on orbits oblique to the thin disk, the best known is  $\alpha$  Boo (Arcturus). It is convenient here to use the term “Population P” for the ensemble of non-brown-dwarf, non-white-dwarf stars in the much larger, 3000-ly radius, subdisk-of-the-thin-disk from which our (tiny) Sample S is drawn. This P-region is itself only a (tiny) fraction of the overall galactic thin-disk region of stars,  $\sim 50,000$  ly in radius. The various pages at [atlasoftheuniverse.com](http://atlasoftheuniverse.com) are a useful resource for visualization of the Orion Arm, furnishing both a zooming-out to the wider galactic context and a zooming-in to detailed features within the Arm.

Sample S, being formally defined by an apparent-magnitude cutoff as opposed to a distance cutoff, is itself far from statistically representative of Population P. (a) In P, the O stars are vanishingly rare. A tabulation by Glenn Ledrew, in *JRASC* **95** (2001), pp. 32ff (bibcode [2001JRASC..95...32L](https://ui.adsabs.org/2001JRASC..95...32L)) suggests an O-star frequency within P of just 0.00003%. By contrast, O stars comprise a hefty  $\sim 2\%$  of S. A similar overrepresentation occurs for the B, A, F, G, and K stars, with Ledrew’s tabulation suggesting that these MK temperature types might have a respective frequency within P of 0.1%, 0.6%, 3.2%, 8.0%, and 12.9%. By contrast, the first three of these five rare types comprise  $\sim 30\%$ ,  $\sim 20\%$ , and  $\sim 10\%$ , respectively, of S, and the last  $\sim 20\%$  of S. (b) In P, something on the order of 76% or 78%—different authorities are perhaps mildly discrepant—must be M stars. (Ledrew’s tabulation, in particular, suggests an M-star frequency of 78.2%.) Only a few of these (the Ledrew tabulation suggests 0.04%) have evolved to beyond the main-sequence stage of stable-core hydrogen fusion. By contrast, the M stars comprise just  $\sim 5\%$  or  $\sim 10\%$  of S. All of them have evolved beyond the Main Sequence, having started their lives as types hotter than M or K.

The statistically anomalous character of S is further illustrated by the fact that in S, in each of the Big Six MK temperature types hotter than M, the numerical majority comprises the stars that have ended stable-core hydrogen fusion (and so have, as a generally reliable rule—we return below to a necessary caveat regarding reliability—evolved out of MK luminosity class V into one of the brighter MK luminosity classes IV, III, II, or I). In Ledrew’s tabulation, the percentages of evolved stars in F, G, and K, as a percentage of the overall respective F, G, and K populations, are just 2.0%, 2.5%, and 3.8%. Consistently with this, the 1991 Gliese-

Jahreiss catalogue of the nearest 1000 stars (containing, admittedly, not only the local OBAFGKM VI, V, IV, III, II, and I stars, but also at least many of the local white dwarfs) assigns less than 1% of its population to MK luminosity classes IV, III, II, or I.

Sample S—so rich in varieties of star statistically infrequent within Population P—harbours physical extremes. Although the extremes are for the most part not written into our table, they can be studied easily from such sources as Prof. James Kaler’s [stars.astro.illinois.edu/sow/sowlist.html](http://stars.astro.illinois.edu/sow/sowlist.html).

At least 58 of our 325-star set each radiate, across the full spectrum from X-ray through UV and optical to IR and radio, at least as much power as is radiated by 10,000 Suns. Possibly the most dramatic is  $\zeta$  Ori, with a bolometric luminosity of 375,000 Suns—making  $\zeta$  Ori notable not within S alone, but even within the overall galaxy. Several others are not far behind, among them  $\zeta$  Pup (360,000 Suns, suggests Kaler, as of July 2008 revising his earlier, circa-1999, suggestion of ~750,000 Suns). We believe that just two stars in Sample S, nearby  $\tau$  Cet and nearby  $\alpha$  Cen B, radiate more feebly than our Sun, each at about half of the Sun’s bolometric luminosity.

The principal determinant of stellar luminosity, for any given phase in stellar evolution, is mass, with even small variations in mass translating into large variations in energy output. The exceptional luminosities of  $\zeta$  Ori and  $\zeta$  Pup, in particular, are a consequence of their exceptionally high respective masses, 20  $M_{\odot}$  and 40  $M_{\odot}$ . (Kaler now suggests 40  $M_{\odot}$  for  $\zeta$  Pup, while having previously suggested 60  $M_{\odot}$ . He additionally notes from the literature the lower suggested value of 22.5  $M_{\odot}$ .)

Theory does predict, although our small Sample S does not succeed in illustrating, the possibility of masses up to the Eddington stellar-mass limit, somewhere above 100  $M_{\odot}$ , and even of some “super-Eddington” stars. (Eddington’s limit is by definition attained when luminosity rises so high as to make the outward radiation push, tending to tear a star apart, exceed the inward gravitational pull.)

Rotation periods in Sample S vary from far in excess of our Sun’s to far short of our Sun’s (which we may here take as a nominal 27 d; refined treatments of solar rotation provide for rotation-period variations both with solar latitude and with solar depth). Spectroscopy yields for  $\gamma$  Cep a period of 781 d, i.e. of 2.14 y. Kaler suggests that the respective rotation periods of  $\alpha$  Hya and  $\epsilon$  Crv could be as long as 2.4 y and 3.9 y. Perhaps our slowest rotator, however, is  $\alpha$  Ori, now (cf [2009A&A...504..115K](#)) assigned the period of 8.4 y. At the other extreme, Kaler suggests for  $\zeta$  Aql A,  $\alpha$  Aql, and  $\zeta$  Lep, respectively, 16 h, at most 10 h, and around 6 h.

Radii (as distance from centre to outermost opaque layer, perpendicular to the axis of stellar rotation) are typically greater than the solar radius. Two notable instances of stellar expansion—in other words, of notably tenuous stellar atmosphere—are  $\alpha$  Sco (with a radius of 3.4 au, not far short of the Sun-Jupiter distance) and  $\alpha$  Ori (with a radius of 4.1 au or 4.6 au from interferometry, or alternatively 3.1 au or 3.4 au from luminosity-temperature deductions). A still more notable, but also very hard-to-determine, case is “Herschel’s Garnet Star,”  $\mu$  Cep A, with radius variously estimated as 4.5 au, 5.6 au, 6.6 au, or 7.7 au. Results in these extreme cases depend strongly on the wavelength selected for evaluating opacity. Observations within Population P do indicate, although our Sample S does not succeed in illustrating, the possibility of still more-extreme stellar radii, to values approaching ~10 au.

The broad range of temperatures (a topic whose MK conceptual subtleties we examine in subsection 4.1, below) is reflected in the fact that all of the Big Seven temperature-type bins in the traditional MK temperature sequence are well occupied, however statistically skewed (as we have argued above) is the distribution in the MK Big Five luminosity-class bins. At the MK temperature extremes are the hot  $\zeta$  Pup (O5; 42,000 K) and the cool Miras, most famously  $\omicron$  (omicron) Cet (M5–M10; a typical temperature for this variable is variously suggested as ~2000 K or ~3000 K).

Interesting spectral anomalies in Sample S include the “Be phenomenon” and “shell spectrum” stars, as discussed at length in our final subsection.

### **SECTION 3: Initial user guide to the columns in our 317-entry table**

In our first column, we use the flags “+nP” (n = 1, 2, ... ) for companions of sub-stellar mass, such as have been found outside our Solar System, in an accelerating sequence of discoveries, from the 1990s onward, that has now reached even the tiny Sample S. Such companions are typically planets but could in principle also be brown dwarfs. We do not attempt here to define formally the difference between a planet and a brown-dwarf companion.

In this same column, we apply the WDS naming scheme for multiplicity, both in the case of true binarity

and in the case of mere optical doubles (in all but eleven awkward cases, as noted in section 1 above, putting into the first column just the name of the brightest WDS-catalogued component; but we additionally try to supply particulars, at any rate in the online table, for binary and mere-optical companions brighter than mag. 10, in the “Remarks” column, while for the most part regarding binary and mere-optical companions fainter than this limit), using underlining, as discussed in Section 4 below, to flag instances in which a binary system possesses a published orbital solution.

*Apparent Visual Magnitude ( $m_v = V$ ):* Apparent magnitudes, with “v” appended for large-amplitude variables, are from *HIPPARCOS*. In the case of variable, we take as authoritative the ranges (where possible, in V), and also the periods, published in the online AAVSO(VSX) database. Our reasoning here is that AAVSO has critically appraised and filtered data originally presented in more upstream sources, such as the primary (journal-article) literature. Our “V” is the usual “V” of UBV photometry, as introduced by H.L. Johnson and W.W. Morgan in [1953ApJ...117..313J](#). The (yellow) V filter corresponds roughly to the response of the eye. We retain, without having attempted our own independent error analysis, the assertion of our Handbook predecessor R.F. Garrison (working essentially before *HIPPARCOS*) that the “probable error” of each of our cited V values is at most 0.03 mag. (in other words, that of the actually and potentially available V measurements from the world’s duly competent photometry facilities, at least half will lie within 0.03 mag. of our own cited V values). Some small inaccuracies in magnitudes may be present in cases of combined light: readers needing confirmation may check our values against WDS, or where possible against the magnitude-specifying atlas pages of AAVSO. (By the nature of its mission, AAVSO is constrained to supply in its cartography not only details of variables, but also magnitudes of stars that are constant, and that can be used by amateur photometrists as comparison stars and check stars.)

*Spectral Classification (MK Type):* The “MK temperature type” (O, B, A, F, G, K, M) is given first, followed by a finer subtype (0–9) and an “MK luminosity class” (Roman numerals I–V, with “a” or “b” added occasionally to indicate slightly brighter or fainter stars within the class). As we discuss in detail in subsection 4.1 below, O stars are the hottest, M stars coolest; Ia stars are termed the most luminous “supergiants”; III stars are termed “giants”; and V stars are termed “dwarfs.” V stars form the largest class in the cosmos, comprising the observational Main Sequence (MS) (as a region in two-dimensional MK-luminosity-class-versus-MK-temperature-type classification space). Other MK symbols include “e” for hydrogen emission; “f” for broad, non-hydrogen emission in hot stars; “m” for strong metallic absorption; “n” or “nn” for unusually broad absorption; “p” for peculiarities; “s” for a mixture of broad and sharp lines; and “:” for a minor uncertainty. (The flags “n” and “nn” are a signature of rotation. It seems that historically “n” and “nn” signified “nebulous,” as references to the photographic-plate appearance of a rotationally broadened absorption line.) Where a single star (e.g.  $\alpha$  CMa A) is given two types, with the second flagged “m,” the first is the type that best characterizes the hydrogen lines, the second the type that best characterizes the metal lines.

MK classifications are in some cases controverted. We have inherited our own types for the most part from the judgements of our predecessor R.F. Garrison, who, as a principal historical authority in MK classification, drew both on what he judged to be the best of the literature and on some of his own unpublished classifications. As of 2021 Jan. 13, we have made a modest beginning at flagging the cases of controverted MK phenomenology (in our online, but not in our printed-edition, “Remarks” column), in two ways: (a) Where the literature suggests a real difficulty in MK classification, we draw attention to the difficulty, discussing it in a few words; (b) Where we have not found reason in the literature to suspect an MK-classification uncertainty, but nevertheless find our assigned MK type diverging (even in a small way) from the type assigned as of epoch 2021.5 in the official United States Naval Observatory and HM Nautical Almanac Office publication *Astronomical Almanac*, Section H (bright stars), we document the divergence, without further discussion.

*Parallax ( $\pi$ ), Proper Motion ( $\mu$ ), and Position Angle (PA):* Parallaxes, in milliarcseconds (mas), proper-motion vector norms ( $''/y$ ), and vector position-angles (degrees, from N through E) are derived from the *HIPPARCOS* 2007 data reduction, with a few exceptions. It may be hoped that in future years more precise parallaxes will be forthcoming from the *Gaia* mission, which has now found an engineering solution significantly easing its initial restriction to the fainter stars. (Detector overload had been feared.) Like *HIPPARCOS*, *Gaia* has to cope with the special challenges posed in measuring to high precision (i) the parallax of a (orbitally wobbling) star possessing a gravitationally bound, and not necessarily well documented, companion, and (ii) the parallax of a star with perturbed photosphere, and consequently with displaced

photocentre (as when a tight binary system contains a bright mass-transfer stream).

*Absolute Visual Magnitude* ( $M_V$ ) and *Distance in Light-Years* ( $D$ ): Absolute magnitudes and distances are determined from parallaxes, except where a colon follows the absolute magnitude; in these cases, both quantities are determined from a calibration of the spectral classification. The absolute magnitude is left uncorrected for interstellar absorption. The appropriate correction is typically  $\sim +0.06$  mag. per 100 ly outside the (admittedly very-far-from-spherical) Local Bubble, i.e. beyond  $\sim 100$  ly. A special difficulty, not fully grasped by us, arises in the case of the controverted  $\epsilon$  Aur system distance (for which we now use *Gaia* DR2, additionally supplying references to the recent literature).

We take account of uncertainties in parallaxes by stating the derived distances, in ly, to no more than the appropriate number of significant figures (rounding where necessary). In cases where rounding would itself be misleading, we use a tilde as an indicator of imprecision.

*Radial Velocity* ( $V_{\text{rad}}$ ): Radial velocities are from BSC5. “SB” indicates a spectroscopic binary, an unresolved system whose duplicity is revealed by periodic Doppler oscillations in its spectrum and for which an orbit is possibly known. If the lines of both stars are detectable, “SB2” is used; “+” and “-” indicate, respectively, motion away from and toward the observer. “V” indicates a variable velocity in a star not observable as a spectroscopic binary. (In most “V” cases, the orbit is unknown.)

**Remarks:** Remarks include data on variability and spectra, particulars of any companions, and (for the most part, only in our online table) prominent bits of observational-astronomy news. We are often a little casual with rounding, stating physical quantities for a given star (as, to take a random example, the angle between the  $\alpha$  Cen AB orbital plane and the plane of the sky) to a lower precision than is now available from the primary literature. In a departure from our practice prior to 2017, we now give star names in all and only those cases in which star names are formally promulgated in the International Astronomical Union (IAU) star-naming project, as launched in 2016 at [www.iau.org/public/themes/naming\\_stars](http://www.iau.org/public/themes/naming_stars). Readers requiring further information on names could start with the individual star descriptions in [stars.astro.illinois.edu/sow/sowlist.html](http://stars.astro.illinois.edu/sow/sowlist.html). Richard Hinckley Allen’s 1899 book *Star Names: Their Lore and Meaning* has been much cited over the decades. More recent scholarship, with due professional attention to Arabic philology, is, however, presented in Paul Kunitzsch and Tim Smart, *Short Guide to Modern Star Names and their Derivations* (Wiesbaden, 1986), and (by the same pair of authors) *Dictionary of Modern Star Names: A Short Guide to 254 Star Names and their Derivations* (Cambridge, MA, circa 2006). In the **Remarks** column, a **boldface** star name indicates a navigation star.

## **SECTION 4: Supplementary user guide, concerning our treatment of double-star astrometry**

### **4.1: General background remarks on double-star astrometry**

#### **4.1.1: Introductory remarks; prevalence of binarity in System S:**

The observer at the eyepiece seeks physical understanding. Here is a speck of starlight, and here at medium power, around 10” (around a quarter-Jupiter) away is another, with perhaps an intriguing tint difference between pure white and yellowish white, over and above a notable magnitude difference: how far away is each of these two stars, and what stages have they attained in their respective lives, and how do their masses and photospheric temperatures differ?

When the specks are paired, as in this imagined example, a further question arises, however, no less important than the ones already mentioned. Are these two physically unrelated stars, neighbours on the two-dimensional celestial sphere through coincidence (with one star perhaps twice or five times as far from Earth as the other, but perhaps with the stars even at rather similar distances from Earth)? Or is it, on the contrary, the case that each star experiences the gravitational attraction of the other so strongly as to keep the pair in a mutual orbit, constraining them to move through galactic space as a binary system?

Most stars in our galaxy, and in particular most stars in that portion of our galaxy that is our nearby Population P, not only are red dwarfs in the stable core-hydrogen-fusion phase of their lives but are solitary. (Our own galaxy is a barred spiral. In elliptical galaxies, red dwarfs are still more common, and solitary stars therefore presumably likewise still more common.) However, binarity becomes more prevalent as stellar mass increases. In our Sample S, high-mass stars predominate, and binarity is correspondingly more evident within Sample S than in the overall Population P.

#### **4.1.2: Hierarchically organized systems; contrast with clusters:**

A situation by no means rare in binarity-rich Sample S is the many-star system hierarchically organized, with binarity at each hierarchy level. One of the good instances is the six-star system whose combined light becomes, in a compact amateur atlas, the starlight point  $\alpha$  Gem, and whose most prominent star is Castor (with the ancient name “Castor” since 2016, under IAU rules, designating just a single star, but before that ruling often used loosely, as a name marking merely the overall naked-eye point of light).

We may have, for instance, stars  $w$  and  $x$  in a mutual orbit (with  $w$  describing an ellipse about the  $wx$  centre of mass  $C_{wx}$ , and  $x$  for its part also describing an ellipse about  $C_{wx}$ , but with some remote star  $y$  experiencing the entire  $wx$  system as essentially a point mass). In an appropriately chosen rest frame,  $y$  describes some rather wide, rather slow orbit in tandem with the  $wx$  pair. In this situation,  $C_{wx}$  moves in an ellipse about the  $wxy$  centre of mass  $C_{wxy}$ , and  $y$  for its part also moves in an ellipse about  $C_{wxy}$ . There can now be a still more remote star  $z$ , experiencing the  $wxy$  trio as essentially a point mass. In such a case,  $C_{wxy}$  will, in an appropriately chosen rest frame, describe an ellipse around  $C_{wxyz}$ , and  $z$  will for its part also describe an ellipse about  $C_{wxyz}$ .

Here “appropriateness of choice” consists in taking some inertial frame of mass in which the centre of mass for the given pairing is held at rest. For present purposes, it is enough first to confine our discussion to classical Newtonian mechanics (neglecting the general relativity analysis of gravitation as a geometrical feature of spacetime), and then within this confined framework to further bypass a conceivable deep conceptual problem, saying—simply and loosely—that an inertial frame anchored on some point in space is one in which that point is held at rest (for instance, by being made the origin of Cartesian coordinates for that frame), and in which additionally the frame is “stipulated not to rotate.” (A deep question, extraneous to our own limited purposes in this Handbook article, does admittedly arise even in the confined Newtonian-mechanics framework: with respect to what physical standard is it asserted that a given frame “does not rotate”? Without venturing into details here, we remark that in our Handbook view, the deep question—historically prominent in the Leibniz-Clarke correspondence—can be given a conceptually satisfactory answer, by defining “does not rotate” in terms of “absence of centrifugal pseudo-forces violating Newton’s force-as-product-of-mass-and-acceleration principle.”)

Still further geometries are possible in the case of four stars. In particular, as an alternative to the situation just described,  $wxyz$  could instead be a double double, with  $wx$  a tight pair, and  $yz$  a tight pair, and with these two tight pairs sufficiently distant from each other to make each experience the other as essentially a point mass (with, in this case,  $C_{wx}$  and  $C_{yz}$  describing, in an appropriate choice of rest frame, elliptical orbits about  $C_{wxyz}$ ).

Does an intricate system of, say, eight stars, organized into a four-level hierarchy as “{{{1,2},{3,4}},{5,6}},{7,8}” or into a six-level hierarchy as “{1,2,3,4},{5,6},7,8” differ in kind, or only in degree, from a stellar cluster (be it an open cluster, such as the Hyades, or a globular, such as M13)? We argue that the difference is one of kind.

In developing our argument, we first run through some formal preliminaries.

Consider any pair of point masses  $P$ ,  $Q$  in three-dimensional space, subject to any arbitrary assemblage of forces, and never occupying the same location in space at the same time, and with their respective masses constant through time. For each member of the pair, in any frame of reference, its velocity at any instant (no matter how simple or intricate may be the assemblage of acting forces) is perpendicular to its angular momentum at that instant. We call this the “Basic Momentum-Geometry Fact.”

For any binary point-mass system  $P$ ,  $Q$  (perhaps experiencing some very intricate assemblage of forces, some of them exerted somehow by bodies other than  $P$  and  $Q$ ), call a force experienced by  $P$  or by  $Q$  “central” if and only if it is either parallel or antiparallel to the vector from  $P$  to  $Q$ . Newtonian gravitation in an isolated two-body system is a central force, as also are electrostatic forces of attraction and repulsion for isolated two-body systems in particle physics. (More concretely,  $P$  experiences in the astronomical or particle-physics isolated-binary-system case a force exerted by  $Q$ , acting along the  $P$ – $Q$  line, whereas  $Q$  experiences an equal and opposite force exerted by  $P$ .) If, as is the case for an isolated two-point-mass gravitational or electrostatic system,  $P$  experiences at every moment nothing but the one central force exerted on it by  $Q$ , and  $Q$  experiences nothing but the equal-and-opposite central force exerted on it by  $P$ , then (via a one- or two-line proof in differential vector calculus) the angular momentum vector of  $P$  is constant and the angular momentum vector of  $Q$  is constant. But then, by the “Basic Momentum-Geometry Fact,” each of  $P$  and  $Q$  describes, in any inertial frame in which  $C_{PQ}$  is at rest, a curve confined to a plane.

In astronomy, gravitating bodies are often close to possessing a spherically symmetric mass distribution. It



is provable (with a many-line argument; Newton held up publication of *Principia* by a couple of decades until he at last had the proof) that such bodies behave as point masses.

With these formal preliminaries complete, we now give our argument.

In a hierarchical system, each component (whether a single star, a pair of stars, a double double, or a lone outlier in tandem with a tight double, ...) experiences as non-negligible only a central force exerted by its nearest companion. It therefore describes, in any inertial centre-of-mass frame for that binary within the hierarchy of nested binaries, a curve that is confined to a plane. In an arbitrary stellar cluster, on the other hand, the described curves are not in general plane curves, but can be so-to-speak warped (“twisted,” “skewed”).

We add here that warped-curves cases might arise even in systems simpler than open clusters and globular clusters, including groupings that are not stable: one example of such a small, unstable, grouping is the handful of stars at the heart of the Orion Nebula, whose four brightest members constitute the Trapezium (and whose very brightest member shines at mag. 5.1, less than two magnitude steps below some of our own “Brightest Stars”).

#### **4.1.3: Two-body systems and the conic sections, and orbiting as a kind of oblique falling:**

None of this argument makes any assumptions about the mathematical form of the central force. It is, however, probable (again as a non-trivial theorem, requiring a multi-line proof) that in the special situation in which the central force obeys an inverse-square law (as is the case for gravitation, and indeed also for the electrostatics dominant in particle physics), the curve is not only confined to a plane, but assumes the specially simple form of a conic section. The section is a hyperbola if the central force happens to be repulsive and is a hyperbola or parabola or ellipse if the central force happens to be attractive. The hyperbola case is illustrated in the mechanics of our Solar System by those comets that are moving too quickly to be captured into closed-curve orbits around the Sun. Ellipses are of course common in celestial mechanics: for an isolated binary in an inertial centre-of-mass frame, there is a plane P such that each of the two components describes an ellipse in P with the centre of mass at one focus, with the two components always on opposite sides of the centre of mass, and with the more massive component moving in the smaller of the tandem ellipses, and the tandem ellipses being of the same shape even if quite different in size (being, that is, figures in P that are similar-even-if-not-congruent). That conic section that is the parabola is, on the other hand, from the point of view of real-life celestial mechanics, a mere mathematical idealization, constituting the so-to-speak infinitely thin boundary between the cases of the hyperbola and the cases of the ellipse. More formally, a parabolic trajectory is realized in an inertial centre-of-mass rest frame of an isolated two-point-mass system when and only when the relative speed of the two masses exactly equals the least upper bound of those various relative speeds, relative to the centre of mass, which are low enough to yield an ellipse.

The soccer ball, as a projectile on the sports ground, constitutes a two-body system with Earth. It is often stated, as an approximation, that the impelled ball describes a parabola. Here, however, the truth is that the trajectory is a segment of an ellipse, minutely divergent from a parabola, and that the trajectory would become a perfect parabola if the gravitational field on the sports ground were, contrary to fact, to be of constant direction. (Take the Earth to be a sphere of perfectly spherical mass distribution: then the gravitational field across the soccer ground, exerted by Earth on the ball, changes everywhere in direction, pointing everywhere in the soccer ground to that single point that is the Earth’s centre.)

In the limiting case in which a soccer ball is impelled almost directly upward, and so falls almost directly downward, the trajectory becomes a segment of some almost-degenerate ellipse, with its minor axis of almost negligible length. Since Earth is so overwhelmingly more massive than the soccer ball, the common centre of mass of the Earth-ball system nearly coincides with the centre of the Earth, and the Earth’s ellipse around the centre of mass becomes correspondingly of sub-sub-atomic dimensions. (Admittedly, this is the situation in a Newtonian setting. It will be interesting to see what becomes of such entailed sub-subatomic ellipses if, at some future era, general relativity becomes successfully unified with quantum mechanics.) Upon reflecting on this sportsground example as an extreme case, it can rather soon be seen (we omit details) that a binary system in stellar astronomy is a case of falling—in which, however, the two bodies fall toward each other in such a way as to stray a little off the line at any instant connecting them, and so are destined never to meet up. If the falling is at all times only a little off the instantaneous connecting line, the ellipse is severely elongated (has an “eccentricity” just slightly less than 1; in our brightest-stars table “Remarks,” we write “*e*” in our occasional reports of known orbit geometries); if, on the other hand, the falling is at each instant as far off the

instantaneous connecting line as geometrically possible, so that at each instant each body is moving perpendicularly to what is at that instant the inter-body connecting line, then the ellipse is a circle (with  $e=0$ ).

#### **4.1.4: Binary systems and the determination of individual masses:**

For foundational astrophysical reasons, much effort has historically been, and is still now being, expended on documenting binary stars. This work was pioneered with the filar micrometry of William Herschel (1738–1822) and (more systematically) F.G.W. von Struve (1793–1864). The work took on fresh vigour with the late-Victorian advent of radial-velocity spectroscopy, as spectrograms began to be measured under the microscope for Doppler shifts. Since 1980 or so, it has taken on still greater vigour with the rise of interferometry—speckle interferometry and aperture-masking interferometry in the case of observatories possessing lone telescopes of large aperture, but more notably with optical tables and beam combiners, at those few facilities in the astronomical community possessing multi-telescope interferometric arrays (notably, at CHARA, NPOI, and VLTI).

Always, from the pioneering filar micrometry onward, the astrophysical motivation has been the same. Once a full orbital solution for a binary system of known distance is determined, the individual masses of the two components are known—both (*a*) as multiples of Solar System quantities and (*b*) in the absolute *Système international d’unités* (SI) laboratory unit, which is the kilogram. Here a “full orbital solution” is a set of half a dozen geometrical parameters, or “orbital elements,” in essence angles, describing the ellipticity of the orbit and its orientation in three-dimensional space (including its angle of inclination with respect to the plane of the sky). With these, plus a determination of the distance to the binary system, the orbital trajectory of the binary (in any inertial centre-of-mass rest frame) is fully described, in particular with the length of its semimajor axis determined in the laboratory SI unit of metres.

It is, admittedly, an intricate task to proceed to the orbital elements from the little that is available at the telescope. In the case of traditional filar microscopy, orbital elements are in principle obtainable from some years or decades of raw astrometry, with each night supplying just the angular separation of the components and their position angle (as an angle in the half-open interval  $[0^\circ, 360^\circ]$ , taken from sky north through east, south, and finally west; in our table “Remarks,” we write “PA”). Also in principle obtainable are orbital solutions for binaries of known distance in which there is no filar-micrometer astrometry, and also no other (for instance, interferometer-procured) astrometry, but in which the plane of the orbit happens to be exactly perpendicular to the plane of the sky, and in which spectroscopic Doppler-shift measurements have over many successive observing sessions supplied the changing radial velocities of each of the two components. Although the situation in which the plane of a binary orbit is exactly perpendicular to the plane of the sky is seldom, if ever, realized in observational work, the situation is approximated to usable precision by cases in which the two stars are found to eclipse each other (as with, e.g.  $\beta$  Per Aa1 (Algol) and  $\beta$  Per Aa2). Finally, an orbital solution may be obtained from a combination of (perhaps imperfect) astrometry and (perhaps imperfect) spectroscopy.

Let it, in any case, now be taken that a binary system of known distance has had its orbital elements determined, by some means or other.

The solution to problem (*a*) (individual stellar masses to be determined as multiples of Solar System quantities) rests on the fact that there exists a “universal gravitational constant”  $G$ , such that for any two-body system of constant point masses  $m_1$  and  $m_2$ , in the gravitationally bound, elliptical-orbit case the sum  $m_1+m_2$  obeys, under Newton’s generalization of Kepler’s Third Law, the equality “ $m_1+m_2 = (4\pi^2 a_{1,2}^3)/(G P_{1,2}^2)$ ” (with  $a_{1,2}$  the length of the semimajor axis of the orbit (in any convenient inertial rest frame of the  $m_1, m_2$  centre of mass), and  $P_{1,2}$  the  $m_1, m_2$  orbital period). The law as stated here is independent of units: masses could be measured in kilograms, or in any other convenient units; the distance, which is  $a_{1,2}$ , could be measured in metres, in light-seconds, or in any other convenient units; and time could be measured in any convenient units. Let, now,  $M_{EM}$  and  $M_\odot$  be the respective masses of the Earth-Moon binary and the Sun. It then follows as a special case, and under the (in practice sufficiently good) idealization of the Earth-Moon binary and the Sun as an isolated system of two point masses that  $M_{EM}+M_\odot = (4\pi^2 a_{EM,\odot}^3)/(G P_{EM,\odot}^2)$  (where  $P_{EM,\odot}$  is the orbital period, in any convenient inertial rest frame of the centre of mass of those two entities, which are the tight Earth-Moon binary and the Sun, of that wide binary, which is the Earth-Moon centre-of-mass and the Sun). Equating the ratio of the left-hand sides of this pair of equations with the ratio of the right-hand sides of this pair of equations, and additionally equating  $a_{EM,\odot}$  to the physical quantity 1 au as defined since 2012 in the SI unit of metres at IAU (this equating, while not exact, is an excellent approximation), we have  $(m_1+m_2)/(M_{EM}+M_\odot) =$

$(a_{1,2}/1 \text{ au})^3/(P_{1,2}/1 \text{ y})^2$ . Conveniently, however,  $M_{EM}$  is to one significant figure a mere 3-millionths the mass of the Sun. This justifies replacing, for most ordinary astrophysical purposes,  $M_{EM}+M_{\odot}$  with  $M_{\odot}$ , yielding as a final, good approximation, the following solution to problem (a):  $m_1+m_2 = M_{\odot} (a_{1,2}/1 \text{ au})^3/(P_{1,2}/1 \text{ y})^2$ .

It remains to determine not just  $m_1+m_2$  in terms of the quantities 1 au, 1 y, but the individual stellar masses  $m_1, m_2$  in terms of this pair of quantities. This, however, is a comparatively modest further step. Once the orbit of the binary system, in some convenient centre-of-mass inertial rest frame, is given, the mass ratio  $m_1/m_2$  can be found by comparing the respective dimensions of the two similar-though-in-general-not-congruent ellipses (the smaller ellipse for the larger of the two masses) traced around the common centre of mass in any convenient inertial centre-of-mass rest frame. With  $m_1+m_2$  known and  $m_1/m_2$  known, the individual values of  $m_1$  and  $m_2$  follow.

Problem “(a)” has thus been solved without recourse to a laboratory determination of the troublesome constant  $G$ . It is for problem “(b)” (determination of  $m_1, m_2$  individual values in kilograms) that  $G$  itself is needed. Work on the laboratory problem has been proceeding for a little over two centuries. Google or YouTube searches under such terms as “Cavendish experiment apparatus” reveal the possibilities for repeating, under a constrained high-school budget or a frugal lone hobbyist’s budget, the result published by Henry Cavendish in 1798, and falling within around 1% of the now-accepted value. As the current state of the art, where expense is surely not spared, [www.pnas.org/content/113/36/9949](http://www.pnas.org/content/113/36/9949) cites a *Phys. Rev. Lett.* year-2000 determination of  $G$  to an uncertainty of 0.0014%. Even this modern level of precision compares unfavourably with the precision attainable for, e.g. the electron charge-to-mass ratio, the speed of light, the electrical permittivity of free space, and the magnetic permeability of free space. Nevertheless,  $G$ , while continuing to be something of a laboratory embarrassment, is sufficiently well known to facilitate work in those areas of astronomy (notably in planetary science) where actual kilogram masses are useful. Already in Cavendish’s day, for instance, it was determined (we here rephrase Cavendish’s result in modern terminology, while conserving its substance) that the mass-per-unit-volume of Earth is higher than the mass-per-unit-volume of ordinary rock (planet Earth 5515 kg/m<sup>3</sup>, but basalt and granite merely ~3000 kg/m<sup>3</sup>), and that therefore the rocks familiar to geology are not representative of Earth’s deeper interior. SI-unit density determinations, resting on the determination of masses in kilograms, are now needed not in geophysics alone but in exoplanet work, for instance for supporting hypotheses regarding a given exoplanet’s composition (gas, in the manner of Jupiter? or something more dense, in the manner of Earth?).

#### **4.1.5: Some further reading:**

Tutorial resources on the Web include a conspicuously thorough source of pages from an author of the *Cambridge Double Star Atlas* (2009, second edition 2015), Bruce MacEvoy (the colleague author for this book is the celestial cartography authority, Wil Tirion), at [www.handprint.com/ASTRO/index.html](http://www.handprint.com/ASTRO/index.html).

#### **4.2: Our notational conventions in table “Star Name” column for double-star astrometry:**

Our treatment of double stars follows the WDS naming rules, but with additionally our own (purely Handbook-local) underline-flagging convention.

Suppose, as a hypothetical case, that a certain bright naked-eye point source has been familiar from Johann Bayer’s 1603 atlas onward as “omega FooBaris,” or  $\omega$  FBr. Suppose  $\omega$  FBr to have been discovered by some 1830s filar micrometrists to be a tight double, with components separated on the celestial sphere by an angular distance of 0.7”. It does not matter whether the pair is a binary or a mere line-of-sight coincidence: in either case, at the 1960’s launch of WDS, the pairing is catalogued as  $\omega$  FBr A and  $\omega$  FBr B.

Now suppose, as a refinement of this basic scenario, that around 1910,  $\omega$  FBr A was found by some spectroscopist to be a spectroscopic binary (in our penultimate-column notation, to be an “SB”), and that nothing further was known about  $\omega$  FBr A until 1974. What are the 1973 WDS implications of the 1910 discovery? Under WDS rules,  $\omega$  FBr A had at that early stage in the development of WDS to be  $\omega$  FBr A (not  $\omega$  FBr Aa,  $\omega$  FBr Ab), since as of 1973 its components had not been measured in the celestial-sphere terms of PA and angular separation.

Stellar interferometry was launched in a modest way in the 1920s. It is perhaps reasonable to say that a “Second Generation” of optical interferometers was ushered in by the team of Robert Hanbury Brown, operating the Narrabri Stellar Intensity Interferometer from 1963 to 1974. Suppose, then, that in 1974 some interferometer, such as Narrabri, succeeded in resolving  $\omega$  FBr A into two components, say at a measured

separation of 0.1". At this stage, the WDS multiplicity catalogue was at last able (and under its self-imposed rules was required) to refer not to " $\omega$  FBr A" but to  $\omega$  FBr Aa and  $\omega$  FBr Ab.

Finally, suppose that in the current, arguably "third," generation of optical interferometry, some such instrument as CHARA or NPOI or VLTI, perhaps working in the year 2020 or 2030, measures  $\omega$  FBr Ab itself as a (very tight, very rapid) binary, with the separation even at apastron found to be just a few tens of milliarcseconds. At this stage, WDS is able (and under its self-imposed rules is required) to refer not to  $\omega$  FBr Aa and  $\omega$  FBr Ab but, rather, to  $\omega$  FBr Aa,  $\omega$  FBr Ab1, and  $\omega$  FBr Ab2.

In the leftmost column of our table, we indicate with underlining that a published orbital solution is asserted to exist in WDS. In our notation, " $\omega$  FBr Aa" signifies the existence of a published orbital solution for  $\omega$  FBr Aa and  $\{\omega$  FBr Ab1,  $\omega$  FBr Ab2 $\}$  (where the star  $\omega$  FBr Aa experiences the outlying pair of stars  $\omega$  FBr Ab1,  $\omega$  FBr Ab2 as essentially a point mass), whereas " $\omega$  FBr Aa" signifies in our notation the existence of a published orbital solution for the entire  $\{\omega$  FBr Aa,  $\omega$  FBr Ab1,  $\omega$  FBr Ab2 $\}$  three-star system, considered as a point mass, in its wide and slow orbit with the remote  $\omega$  FBr B. In various cases in which this notation is, whether definitely or at least arguably, unclear in its intent, we explain in the "Remarks" what is and is not available in the published  $\omega$  FBr orbital-solutions literature.

Although the presence of underlining in our leftmost column is a safe indication that a given double is a binary, the absence of underlining is not a safe indication that a given double is a mere line-of-sight coincidence. In some cases lacking underlining, it is a known fact that the given double is a binary (typically with some very wide, slow, orbit, that will defy mathematical modelling until some further centuries or millennia of astrometry become available); in other such cases, it is a known fact that the given double is a mere line-of-sight coincidence (for instance, because either the parallaxes or the proper motions of the two stars are severely discrepant); and in very many other such cases, the answer to the question "Binary, or not?" is currently unknown. Although we do not here try to flag the first and the second of these three possibilities in our leftmost column, WDS does try to track the current state of knowledge with its own (duly elaborate) flagging system.

## **SECTION 5: Supplementary user guide, concerning the more detailed interpretation of our MK-classification column**

### **5.1: Conceptual underpinnings of the MK classification system**

In strict conceptual accuracy, the MK temperature types are a purely phenomenological record of which elements are present (*a*) in which stages of ionization, and (*b*) at what densities (in other words, under what local strength of the local downward-directed gravitational field) in the photosphere of the given star.

Decades before the 1943 Morgan-Keenan-Kellman publication of the full two-dimensional MK scheme, it had already been found possible to set up the phenomenological spectral types under our heading "*(a)*" in a single orderly OBAFGKM sequence, in which individual types gave way smoothly to their neighbouring types. (This process was itself not quite straightforward. First came a simple Harvard "A, B, C, D, ..." scheme. This was followed by the realization that "A," for example, linked smoothly in its phenomenology with "B" and "F," with some of the old alphabet having to be altogether dropped or repurposed. In working out this ordering, it was found necessary by the Harvard pioneers to subdivide the OBAFGKM categories, for instance in the sense of "G rather similar to F" and "G rather similar to K" and "G about equidistant between F and K." Hearnshaw's *Analysis of Starlight*, now in its second edition as [2014anst.book.....H](#), is the definitive history both of the MK scheme and of its predecessors.)

It was then not a matter of definition, but of astrophysical discovery (cf, e.g. the already-cited [2014anst.book.....H](#), or again [1994AJ.....107..742G](#), or again the detailed MK reference-work exposition [2009ssc..book.....G](#)), that the OBAFGKM sequence corresponded to a temperature-ordered sequence of stellar groupings, running from the hottest photospheres to the coolest, with each of the various subdivisions within each of the O, B, A, F, G, K, and M types corresponding to a particular temperature range.

With the 1943 introduction of the two-dimensional MK scheme, the luminosity classes I, II, III, IV, V likewise had strictly a phenomenological, not an astrophysical, definition (proceeding now from our heading "*(b)*," as opposed to the "*(a)*" that yielded O, B, A, F, G, K, and M). It was then once again conceptually speaking not a matter of definition, but of astrophysical discovery, that the I-through-V sequence corresponded to a luminosity-ordered sequence of stellar groupings, running from the most luminous to the least luminous.

Admittedly, this conceptual picture, for the history of work under our heading “(b),” is idealized. It was evident on the theoretical front even some decades before 1943 that the “(b)”-heading phenomenological features highlighted in 1943 by the developers of the MK taxonomic system and signalling differences in photospheric gas densities (i.e. to differences in the strength of the local downward-pointing gravitational field) in fact correspond to differences in stellar luminosities. The developers of the MK taxonomy thus had a theoretical motivation for their definitions of classes I, II, III, IV, and V, resolutely phenomenological though their definitions were required to be, under observational-astronomy methodology—the MK system now serves as a paradigm of successful taxonomy, even for fields outside astronomy. A classification system is defined in terms of mere phenomenological fieldwork, and yet in the expectation (successfully realized in the case of MK) that the phenomenological classification bins will in due course be discovered by the theoreticians to correspond to relevant, important, physical differences in the materials observed. (Parallels might be suggested with, e.g. 18th- or 19th-century medicine: whereas (i) the old clinical-phenomenology definition of “tertian fever” and “quartan fever,” in terms of the observed duration of body-temperature anomalies, have been found in physiology theory not to correspond to useful fundamental realities at the level of microbiology, (ii) the gross empirical observation, as with the pre-Victorian stethoscope, of heartbeat anomalies has been found to correspond to useful fundamental realities at the level of cardiac neuroanatomy.)

When the MK system was introduced, it was already evident that if the classes I through V signalled a progressive decrease in stellar luminosities, then they had to signal a corresponding progressive decrease in stellar radii. The temperature of a given photosphere determines the amount of energy that photosphere radiates per unit time per unit of photosphere area. Consequently, if two stars in the same temperature type are found to differ in luminosity class, the one in the brighter luminosity class must have a larger total photosphere area, and so must be of greater radius.

It was therefore natural to adopt theory-informed, but nevertheless in official terms purely mnemonic, labels for the phenomenologically conceived luminosity classes, with I called for convenience the “supergiants,” II the “bright giants,” III the “giants,” and IV the “subgiants.” V had to be given some mnemonic label opposed to “giant,” with “dwarf” consequently pressed into service, and “subdwarf” used for the underluminous class VI (important in studies of congenital metallicity, but irrelevant to our own Sample S). (It is admittedly troublesome that the terms “white dwarf”—and nowadays also “brown dwarf”—prove necessary in other contexts, with the “white dwarfs” and the now-celebrated “brown dwarfs” radiating at luminosities far below even classes V and VI.)

## **5.2: MK classification and stellar evolution: preliminary remarks**

In 1943, when the MK system was introduced, stellar-evolution theory was not yet on a sound footing. Only the broad outline, that a star may be expected to increase in photospheric radius after completing the fusion of hydrogen in its innermost portion, was at that point known. With the theoretical nuclear-physics advances of the 1950s and 1960s, and with the advent of increasingly detailed computer modelling from the 1960s onward, it became possible to map the elaborate excursions (we outline these in subsections 4.7 and 4.8 below) that evolving stars perform in the two-dimensional luminosity-class-versus-temperature-type phenomenologically defined MK plane (the “observational HR diagram”). In particular, it is now known that every star in the phenomenological class V in our 325-star set from our 317-entry table is still performing stable fusion of hydrogen in its innermost portion. (We repeat that this class V is best termed, with correct deference to the MK classification conceptual underpinning, not simply the “Main Sequence” (MS), but the “observational MS”—as at p. 342 of the authoritative [2006ima.book....C](#).) Further, membership in the phenomenological class IV is a good (though even in our small 325-star set not an infallible) indicator that stable hydrogen fusion in the innermost portion is over, with the subject star now having performed at least some part of its (in general, elaborate) later-life excursions over the MK phenomenological plane.

The distribution of the set of 325 stars across MK luminosity classes I through V accordingly turns out to be a reasonable indication of the evolutionary spread of the set.

It follows that the naked-eye bright-star night sky is a different place from the daytime sky, with its lone proximate class-V star. Something on the order of a mere fifth of our 325 MK-classified bright nighttime stars (for the most part stars in luminosity class V) resemble the Sun (the sole daytime object in our set of 325 MK-classified bright stars) in stably burning hydrogen at their centre. Even most of these are far hotter than the Sun and are consequently destined to spend less time than the Sun in this process of stable burning. All the rest have

in one way or another moved beyond that stage, as shown by their luminosity classes—with the nocturnal 324 falling overwhelmingly into classes III and IV, but with classes I and II also rather well populated.

### **5.3: MK classification and stellar evolution: starbirth and MS**

A star has at birth (i.e. has upon condensing sufficiently from its local ISM cloud to begin hydrogen fusion) four key characteristics. If the star happens not to be in the disturbing environment of some proximate star (most notably, in the disturbing environment of a binary companion so close as to transfer matter) then these four characteristics jointly entail its various other characteristics, for each point in its entire subsequent career. Prominent among those other characteristics are the duration of overall life, and at each point in the overall life additionally those time-varying key characteristics, which are radius, luminosity, and its photosphere effective temperature. Here, then, are the “Governing Four”: (a) birth-epoch mass (the more massive stars are also the hotter, the more luminous, and the shorter-lived); (b) birth-epoch elemental composition (the most important aspect of composition is simply the birth-epoch “metallicity”—i.e. the extent to which, thanks to the specific properties of the local gestating ISM cloud, the subject star contains at the time of birth any elements, in whatever detailed proportions, heavier than hydrogen and helium); (c) absence or (possible) presence of inherited fossil magnetism, from (possible) magnetism in the gestating ISM cloud; and (d) birth-epoch speed of rotation.

Of the four listed properties, the first is the most important, accounting, along with the accidental circumstances of distance-from-Earth and time-elapsed-since-gestation, for essentially all the stellar variety that the un-aided eye can discern.

Regarding the accidental circumstance of time-elapsed-since-gestation, a parenthetical caveat, relevant even to interpreting the casual naked-eye experience, is needed: stars condensed from the same ISM cloud are of the same age. This is the case not only with binaries but also, more dramatically, with associations (such as the dramatic naked-eye association in the northern sky whose most familiar members comprise  $\beta$  UMa (Merak),  $\gamma$  UMa A (Phecda),  $\delta$  UMa A (Megrez),  $\epsilon$  UMa A (Alioth), and  $\zeta$  UMa Aa (Mizar), in other words comprise all but the first and last of the seven Big Dipper stars).

In contrast with mass and present age, congenital elemental composition does not vary greatly across our set of 325 MK-classified bright stars. The pronounced chemical differences across the set of 325 (evident from the notations for chemical peculiarities in many of the 325 bright-star MK types in our 317-entry table) are due, rather, to processes of stellar aging, notably (i) gravitational settling and radiational lofting of selected elemental species in cases in which the outer layers are quiet (in particular, not rotationally disturbed), and (ii) processes known as “Dredge-Up” (discussed again in subsection 5.8, below), when convection in an evolving star raises such elements as carbon or nitrogen into the photosphere from the buried thermonuclear furnaces.

We will not attempt to discuss congenital magnetism. But we do remark that, like chemical peculiarities, magnetism can develop and change as a star ages (with, for instance, convection in outer layers, under rotation, producing a dynamo, and with the dynamo in turn generating the kind of looping-field locally dipolar magnetic structures present in the Sun, and hinted at in the small telescope by the Sun’s appearance through a hydrogen Balmer- $\alpha$  filter).

The fourth property in our list, congenital rotation, is a consequence of the vagaries of possible motions in the gestating ISM. The local part of the condensing gas was likely to have some kind of coordinated spin, and this spin tended to increase, under conservation of angular momentum, as the gas became more and more condensed—even though some angular momentum also was possibly shed via gas outflows, as the condensation proceeded toward starbirth.

We will not discuss congenital rotation further. We do, however, remark that the rotation speed of a solitary, undisturbed star is once again a property that can evolve as the given star ages, under the combined influence of its evolving mass distribution (although the mass of all but the hottest stars remains rather constant until late in life, after cessation of core hydrogen fusion the mass gets distributed over larger radii, forcing (under conservation of angular momentum) an increase in rotation period and its (possibly, as already noted, evolving) magnetism).

The process of change has two aspects. On the one hand, as an aging star evolves out of luminosity class V into IV, III, and in the case of congenitally massive stars even into II or I, increases in its radius cause (because angular momentum is conserved) a slowing of rotation.

On the other hand, and quite apart from this general slowing-through-bloating, a spin-braking mechanism

exists within class V for those stars that succeed in generating the right kind of local, looping, dipole magnetic-field structures. The mass shed by such a star in winds, although modest, is nevertheless constrained by magnetic fields not to orbit the star freely, but to rotate at about the same angular velocity as the star itself. Under conservation of angular momentum, this so-called “magnetic braking” then slows the rotation. In the overall galactic population of V stars, those cooler than MK temperature type F5 are capable of achieving magnetic braking, and those hotter than F5 are not. The F5 type thus constitutes a so-called “rotation break” within class V.

In our set of 325 MK-classified bright stars, all but 6 of the class V stars lie on the hot side of the break. The brightest V-class stars in Earth’s sky have to be either the most luminous, and therefore the hottest, or those nearest to Earth. The scarcity of V-class bright stars on the slow side of the rotation break therefore indicates that it is the first of these two brightness-promoting characteristics that predominates, in our overall set of 325.

Although we here largely neglect stars in the disturbing environment of other proximate stars, we do have to remark parenthetically that in the case of a close binary, rotation (like also chemical composition) can be affected by processes of mass transfer. This is very notably the case with one of the more heavily studied stars in the 325-member set,  $\alpha$  Leo A (Regulus). Here the rapid rotation is the result of a now-completed spinning-up process, involving a copious mass transfer, from the now diminutive, and therefore now observationally elusive, pre-white dwarf. In the “Remarks” for  $\alpha$  Leo A in the table, we point out that this elusive companion, having for decades escaped observation, is at last reported in [2020ApJ...902...25G](#) as detected spectroscopically.

The F5 “rotation break” within MK luminosity class V is ultimately due to, and is nearly coincident with, a transition (as one proceeds along the observational MS from the hottest stars to the coolest, i.e. as one advances in the sense OBAFGKM) from stars in which the hydrogen fusion is predominantly the work of the carbon-nitrogen-oxygen (CNO) cycle to stars in which the hydrogen fusion is predominantly the work of the proton-proton chain. The point at which the two processes deliver, per unit of fusion-depth mass, roughly equal energy-per-unit time is at or near a total stellar mass of  $1.2 M_{\odot}$ .

To what extent are the four key properties reflected in the MK type of a young star (in observational terms, a star found to lie in MK luminosity class V)?

(a) Mass is well correlated with MK temperature type, in the sense that the OBAFGKM progression within class V proves to be a progression from the most massive stars to the least massive. This fact is itself far from obvious. It was, however, established in the early decades of the 20th century by spectrally classifying the elements of binary systems, of known distance, in which the orbit is not so tight as to allow the disturbing feature of mass transfer, and yet in which the orbit is tight enough, and consequently fast enough, to permit determination of orbital geometry and orbital period. For such binaries, individual masses can be determined from Newtonian mechanics.

(b) Birth-epoch elemental composition is not really reflected in the observationally assigned MK class. We have already remarked that the elemental-composition flags present in many of the 325 bright MK types are due, if not to “Dredge-Up” in the case of an aging star, then to segregation of elements through gravitational settling and radiative lofting (processes that can occur even for a young star, provided its atmosphere is quiet, as in cases where rapid rotation is absent).

(c) The MK scheme does not attempt to flag magnetism, even though magnetism is observed spectroscopically, through the Zeeman splitting of emission and absorption lines when a magnetic field is strong.

(d) Rotation can be inferred in favourable cases, but not in all cases, from the presence of the MK-type flags “n” and “nn.” In a favourable case, a rapidly rotating star is seen more or less equator-on, causing its emission and absorption lines to be Doppler-broadened (since half of the photosphere is rapidly receding from the spectrograph, and the other half rapidly approaching it). In, however, the unfavourable case in which the star is seen more or less pole-on, there is no rotational broadening. A particularly well-known example of a rapid pole-on rotator (with “n” and “nn” therefore absent from the observed MK type) is  $\alpha$  Lyr A (Vega).

We might add by way of background that it is only in recent decades that the detection of pole-on rotators has become feasible at all. If the star is close and bright enough, interferometry, while powerless to detect the shape deformation of a pole-on rapid rotator, may nevertheless succeed in picking up the equatorial darkening that accompanies rotational flattening (in the pole-on case, as an anomalous darkening, over and above the normal “limb darkening,” toward the edges of the interferometrically discerned stellar disk, at whose centre is the Earth-facing stellar pole).

#### **5.4: MK classification and stellar evolution: rotation largely neglected here**

It is now helpful to outline the various possibilities for stellar evolution, as experienced by that majority of stars in the 325 MK-classified set that are already in MK luminosity classes IV, III, II, or I, as opposed to the “observational MS,” which is class V. But an initial caveat is needed: we here largely neglect the disturbing influence of stellar rotation, important though that influence is.

Regarding rotation, we do remark at this point that rotation can produce flows of matter along lines of stellar longitude (“meridional flows”), and that where such flows extend some significant distance into the stellar interior, they help replenish the supply of hydrogen, as a thermonuclear fuel, in the stellar depths. The effect of rotation is in general to somewhat shift the evolutionary track of a star on the phenomenological MK plane (by promoting mixing of stellar layers that would otherwise be more sharply separated) without radically changing the shape of the track.

Difficulties in constructing an evolutionary model for the interior of a rapid rotator are among the themes of Section 1 in [2011ApJ...732...68C](#). This same paper discusses difficulties involved in deducing the mass and age of a rapid rotator, and the problem of deviations from the von Zeipel 1925 gravity-darkening law for oblate-spheroid stars. The law would give the correct result for gravity darkening if the flattened star had a purely radiative envelope. With rotation, however, gravity darkening can lower the photosphere effective temperature at the equator, causing convection to set in there even when the envelope is radiative at the poles. In our 325-star set, this pathology is present in at least  $\alpha$  Aql A (Altair) and  $\alpha$  Cep A (Alderamin).

Even where the convective regime is uniform, it is not possible to assign a single photospheric effective temperature to a rapid rotator: its observed MK temperature type is now a mongrel, the result of light entering the spectrograph from the differing temperature regimes of (hot) poles and (cool) equator.

#### **5.5: MK classification and stellar evolution: structure, energy flows**

As a further preface to details of evolution, it is now necessary to introduce discussion-guiding concepts of stellar structure and stellar energy flows.

A star still stably fusing hydrogen in its innermost portion (whether predominantly via the CNO cycle or predominantly via the proton-proton chain) is said to have a hydrogen-fusing “core.” The layers outside the energy-producing “core” of such a star are said to comprise its “envelope.” Under this definition of “envelope,” the envelope is not a place of energy generation, but merely a place of energy transport. This transport involves a cascade, in which a single core-produced photon is absorbed by some envelope atom, causing the envelope atom to re-radiate multiple photons, each individually less energetic, and with the same aggregate energy as the now-vanished input photon. Each of these less energetic photons is in turn absorbed by some envelope atom in a still higher layer, which in its turn re-radiates a plurality of correspondingly less energetic photons. Eventually, as that outer-skin part of the envelope that is the photosphere is reached, photons begin travelling freely, without processes of absorption and re-radiation.

Those young stars with cores hot enough to have the CNO cycle as their principal mode of hydrogen fusion have convective cores. In the case of the very hottest O stars (perhaps hotter than any of the 35 or 40 or so O stars in our set of 325 MK-classified bright stars), not only the core but even the envelope is convective. The more usual case, however, for a CNO-dominated star, and perhaps the only case appearing for the CNO-dominated subset of our 325-star set, involves a convective core overlain by a radiative envelope.

Where the temperatures at the core are low enough for the proton-proton chain to predominate, the core of a young star is radiative. High envelope opacities in this low-temperature case make radiation an inefficient mode of energy transport, causing envelopes to be convective. As one advances along the temperature sequence in the sense OBAFGKM, stars at first present just a thin convective layer (settling in at a photosphere effective temperature of  $\sim 8300$  K), with convection then running deeper and deeper (and in particular, in the case of our own Sun, as a G2V star, pervading the entire envelope).

Here (once again) a caveat is necessary regarding rotation. A rapid rotator can straddle the  $\sim 8300$  K boundary, with convection absent at its (hot) poles, and at least a thin convective layer present at its (cooler) equator.

As an irony of nature, an extreme case exists at the cool end of the OBAFGKM progression, just as for its already-discussed hot end. In the coolest young M stars, convection extends all the way down to the core. As for the extreme O stars, so also, however, the extreme-M case is irrelevant for us: our set of 325 MK-classified



bright stars contains no young M stars at all.

### **5.6: MK phenomenology of early evolution within the theoretically defined MS**

Having so far mentioned just the “observational MS,” we may now proceed to the theoretical definition of the MS, or more strictly of departure-from-MS (and soon we shall also be relating this bit of theory to the already-presented observational MS concept). The theoretical MS will turn out (subsection 5.8, below) to be defined in such a way that departure perhaps can occur already within class V, but can also be delayed until an aging star has brightened enough to take it into class IV.

It is a sufficient, although not a necessary, condition for a star lying within the theoretical MS that it be still fusing hydrogen within its core.

Even within this early, seemingly placid, stage of a star’s life, large changes can occur. While our own Sun has another four or five gigayears of life before its core-hydrogen fusion is over, the placid process of early MS evolution will, after just a single gigayear, already drive its luminosity high enough to destroy Earth’s biosphere.

At the heart of this early MS process is a gradual change in core composition, as helium ash accumulates. With the core becoming progressively helium richer even while core hydrogen nuclei continue to fuse, the number of particles constituting the aggregate of gas that is the core progressively falls. Given this rise in the mean mass of the core-gas particles (the free electrons, and a diminishing number of hydrogen nuclei, and a rising number of helium nuclei: but the increased helium comes at the expense of the hydrogen, with two hydrogens yielding one helium) the core, while maintaining the pressure needed to support the overlying envelope, is under the Ideal Gas Law forced to contract. Under a physical principle known as the Virial Theorem, half of the gravitational potential energy liberated by the contraction is translated into thermal energy, i.e. into a rise in the temperature of the core. With this rise in temperature, core hydrogen fusion (a process already decidedly dependent on temperature in the case of the proton-proton chain, and very strongly dependent on temperature in the case of the CNO cycle) becomes more vigorous. As a result, the star overall becomes more luminous, and also experiences a modest increase in radius.

It is now convenient to distinguish in our set of 325 MK-classified bright stars between (*A*) the very massive ones (possessing at birth a mass greater than around  $8 M_{\odot}$  or  $10 M_{\odot}$ ) and (*B*) all the others. The very massive stars are destined to die as supernovae (leaving behind perhaps a black hole, perhaps a neutron “star”). The others are destined to die as white dwarfs.

### **5.7: MK phenomenology of evolving high-mass stars (eventual supernovae)**

In observational terms, the very massive MS stars are of MK temperature class O, or else of the hot B subdivisions B0, B1, or B2. In our set of 325 MK-classified bright stars, at least the following ten (in order of increasing RA) can be said with confidence to meet this condition:  $\eta$  Ori Aa (B0.5 V),  $\theta$  Car (B0.5 V),  $\alpha$  Cru B (B1 V),  $\beta$  Mus Aa (B2 V),  $\pi$  Sco A (Fang; B1 V),  $\beta$  Sco Aa (Acreb; B0.5 V),  $\tau$  Sco (Paikauhale; B0 V),  $\zeta$  Oph (O9.5 V), and  $\alpha$  Ara A (B2 V). Additionally, 31 are observed to be on the borderline for meeting this condition (being in IV, or being classified “IV–V,” or being of MK temperature class B2.5).

In the process leading up to the supernova climax, these massive stars will eventually rise in observational terms into the MK “supergiant” luminosity class I. In the set of 325, 35 are clearly now at that late stage in their development.

We will not discuss at any length the details of massive-star evolution once core hydrogen is exhausted, instead contenting ourselves with just five brief points:

(i) The very concept of MS is a little misleading for the most extreme of the massive stars, since in the most extreme cases scarcely has starbirth (the commencing of core hydrogen fusion) been achieved before gross observable evolutionary changes have set in. We will not here attempt to chart this territory (and in particular will not attempt to define for this group of stars the tricky theoretical concept of “departure from MS”). We remark only that a safe early life theoretical concept for the most massive stars is the concept of a mere instant, as opposed to an interval—namely arrival on the “Zero Age [Theoretical] MS,” as the instant at which core hydrogen fusion starts.

(ii) In their so-short lives, these very massive stars fuse progressively heavier elements, in a central aggregation and in shells overlying the aggregation. The fusion after helium is finished is fuelled first by

carbon, then by oxygen, and neon, and magnesium, and finally by sulphur, and silicon, yielding the eventual dumping of iron ash, from sulphur-silicon burning in a shell, onto a growing inert central aggregate of iron.

(iii) A “core-collapse” supernova eventuates after the iron aggregate exceeds the “Chandrasekhar limit” of  $\sim 1.4 M_{\odot}$

(iv) The complexities of core and shell burning, with burning at various levels switching itself on and off in the process leading up to the supernova, translates in observational terms into movements across the MK luminosity-class-vs-temperature-type surface, with luminosity not changing much, but with temperature type changing dramatically (and with changes possible both in the redward, or OBAFGKM, sense and in the blueward, or MKGFABO, sense). Each of the MK types OBAFGKM is represented in our group of 35 supergiants, with at the hot (blue) extreme  $\zeta$  Pup (Naos; O5 Ia) and  $\zeta$  Ori Aa (Alnitak; O9.5 Ib), and at the cool (red) extreme at least by  $\alpha$  Ori Aa (Betelgeuse; M2 Iab),  $\alpha$  Sco A (Antares; M1.5 Iab), and  $\mu$  Cep (“Herschel’s Garnet Star,” M2<sup>-</sup> Ia).

(v) In its redward or blueward progressions, an evolving supergiant can pass, possibly more than once, through the “Instability Strip” (IS) in the luminosity class-vs-temperature type MK plane, thereby temporarily becoming a Cepheid variable. This possibility is presently actualized in our set of 35 class-I stars by (in order of increasing RA)  $\alpha$  Umi Aa (Polaris),  $\beta$  Dor, I (ell) Car,  $\eta$  Aql A, and  $\delta$  Cep A.

### **5.8: MK phenomenology of evolving lower-mass stars (eventual white dwarfs)**

(B) We may now proceed to explain the sense in which, extreme cases of lower-mass cases of rotation aside (where rotation yields gas flows so violent as to leave no gas unmixed), all stars in the 325-star set with masses below  $\sim 8 M_{\odot}$  or  $\sim 10 M_{\odot}$ , and not disturbed by mass transfer from some companion star, proceed from a readily definable theoretical-MS interval of life to the theoretical Sub-Giant Branch (SGB), then to the theoretical Red Giant Branch (RGB), then to either the theoretical Horizontal Branch (HB) or the theoretical Red Clump, then to the theoretical Asymptotic Giant Branch (AGB), and finally (as almost-corpses or corpses) to a post-theoretical-AGB phase, which, in the fullness of time, yields a white dwarf.

It might seem natural to set up a definition of “theoretical MS” for our eventual-white-dwarf stars on which such a star is deemed to leave the theoretical MS upon finishing core hydrogen fusion. The definition actually employed is, however, different (Carroll-and-Ostlie [2006ima.book....C](#), pp. 452, 453). That definition has (surely?) been motivated, over the past few decades of theory construction, by a desire to make the theoretical-astrophysics demarcations correspond as closely as possible to the actual spectrograph-observable changes of direction (i.e. to the actual observed bends) as a star traces its path, over a span of megayears or tens or hundreds or thousands of megayears, on the phenomenological I-through-V vs O-through-M surface. Under the standardly employed definition, a star is said to remain on the theoretical MS not only through the process of luminosity increase attributed in Subsection 4.6 to the Ideal Gas Law, but somewhat later, even a little after the depletion of core hydrogen has brought core fusion to a halt.

The matchup of theory and phenomenology is, despite efforts at fine-tuning the theoretical definitions, imperfect. Awkwardly enough, not only can a star be on the theoretical MS even after finishing core-hydrogen fusion: conversely, a star can even have left the observational MS, in other words can have left the MK luminosity class V, while residing so far within the theoretical MS as to be still burning its core hydrogen. In terms of our table, this awkward converse possibility is illustrated by at least the following (in order of increasing RA):  $\chi$  Car (B3 IV (p?)),  $\lambda$  UMa (Tania Borealis; A1 IV),  $\beta$  Cru A (Mimosa; B0.5 III),  $\nu$  Cen (B2 IV),  $\zeta$  Cen (B2.5 IV),  $\iota$  Lup (B2.5 IVn),  $\alpha$  Tel (B3 IV), and the celebrated variable  $\beta$  Cep Aa (Alfirk; B1 III). Additionally,  $\alpha$  Lyr A (Vega) is still far within the theoretical MS, and yet might erroneously be thought to have evolved to the edge of the observational MS, since its MK class is A0 Va. Here the cause of the “Va,” as distinct from “V,” is rotation (with Vega presenting itself to the spectrograph pole-on while rotationally flattened, in other words presenting a misleadingly increased radius).

At the moment when the depletion of core hydrogen has brought core fusion to a halt, the luminosity of the star derives from fusion in a core-surrounding hydrogen shell, now raised to a fusion-capable temperature by the increased temperature of the inactive helium-ash core. For some modest time after core-hydrogen fusion has ceased, nothing dramatic happens from an observational MK standpoint. Departure from the MS is defined as occurring when the central deposit of non-fusing helium ash becomes so massive as to trigger a rapid internal reorganization of the star, with one or the other of two possible types of rapid contraction, to be distinguished below as “(B.a)” and “(B.b).” This is the point at which something MK-noteworthy, i.e. something that registers

strongly in the spectrograph, finally happens.

(*B.a*) For stars in the 325-star set of mass below  $\sim 1.25 M_{\odot}$ , the growing central deposit of still-inert helium ash becomes so massive as to trigger a further, this time rapid, contraction of the core. Some of the gravitational potential energy present before the abrupt contraction, and now liberated by infall, is under the Virial Theorem translated into an increase in the thermal energy of the shell (in which fusion of hydrogen is therefore in turn speeded up). Paradoxically, although the core has decreased in radius, the rise in shell temperature causes the shell to expand, increasing the radius of the star overall.

Two contending factors are now at work. On the one hand, the star has become more luminous. On the other hand, it is now larger. The latter factor outweighs the former, entailing a fall in the photosphere effective temperature. (Total luminous output from the photosphere is determined both by the attained photosphere effective temperature and by the attained photosphere radius, i.e. by the extent of stellar bloat. If the overall radius increase is large, then a reasonable modest increase in total luminous output has to be accompanied by a temperature decrease.)

In MK observational terms, the star, now defined to have departed the theoretical MS and simultaneously arrived on the theoretical SGB, has on the one hand moved some modest distance upward out of luminosity class V, and has on the other hand advanced redward, i.e. has evolved in the sense OBAFGKM.

(*B.b*) For stars of mass above  $\sim 1.25 M_{\odot}$  (and nevertheless not, we repeat, attaining the  $\sim 8 M_{\odot}$  or  $\sim 10 M_{\odot}$  threshold that makes an eventual supernova possible), the star is found under computer modelling to undergo a more radical internal reorganization. On this more radical scenario, not just the inactive helium-rich core, but the entire star, suffers a rapid contraction. It is this spectrograph-detectable event that is in the “(*B.b*)” case taken to define the end of the theoretical MS phase.

As in the less radical “(*B.a*)” scenario, the star increases in luminosity, with some of the liberated pre-contraction gravitational potential energy once again translated into an increase in temperature (with, once again, a consequent speeding up of hydrogen fusion in the shell). In contrast with the “(*B.a*)” scenario, however, the star is of a reduced radius overall. Under the unavoidable correlation of overall luminous output with both attained photosphere effective temperature and attained photosphere radius, the now shrunken, and yet now brightened, photosphere must now be of a higher temperature. In MK observational terms, the star therefore now quite abruptly not only advances upward in the V–IV–III–II–I sense, but also advances blueward, i.e. evolves in the sense MKGFABO.

Whereas in scenario “(*B.a*)” the star is said to arrive on the theoretical SGB simultaneously with its departing the theoretical MS, in the “(*B.b*)” scenario now under consideration arrival on the theoretical SGB is defined as occurring just a little later than departure from the theoretical MS, with a further episode of core contraction following the overall contraction that under “(*B.b*)” defines departure from the theoretical MS. This further episode of core contraction yields a cooling of the photosphere, and consequently a spectrograph-observable change in the sense OBAFGKM.

In scenario “(*B.a*)” i.e. for stars exceeding  $\sim 1.25 M_{\odot}$ , movement through the SGB is rapid, making the detection of such stars statistically improbable, and generating the so-called “Hertzsprung Gap” in HR-diagram plots of same-age stars when the subject population is so selected as to be duly rich in masses exceeding  $\sim 1.25 M_{\odot}$ , and duly rich both in observational-MS stars and in observational-RGB stars. (Many open clusters meet this sampling requirement.) The statistical improbability notwithstanding, our 325-star set does succeed in capturing several fleeting residents of the Hertzsprung Gap, at any rate (in order of increasing RA)  $\alpha$  Aur Ab (the close Capella companion),  $\epsilon$  Leo,  $\zeta$  Leo A (Adhafera),  $\omicron$  UMa A (Muscida), and  $\zeta$  Her A.

From this point onward, it is no longer necessary to distinguish scenarios “(*B.a*)” and “(*B.b*)” Under both scenarios, residency on the SGB (admittedly started, as we have just said, in one way in the “(*B.a*)” scenario, in a different way in “(*B.b*)” with residency in the former case brief) in due course yields a cooling of the photosphere. With this cooling, the photosphere opacity rises, causing not only the photosphere-proximate layers but even much of the deeper interior to convect. Since, however, convection is a markedly efficient mode of energy transport, the star becomes progressively more luminous and larger, while keeping its photosphere effective temperature roughly constant. As this observationally dramatic increase in luminosity starts, the star is defined as leaving the theoretical SGB and (simultaneously) arriving on the theoretical RGB.

As in the late phases of theoretical MS life, and as in the theoretical SGB, so also here on the theoretical RGB, the star is fusing hydrogen in a shell overlying an increasingly massive, although still inactive, central ball of helium. Now, however, luminosity is much higher than in the MS and SGB phases. As the still-inactive

central helium ball increases in mass, it gradually contracts under its own weight. Some of the gravitational potential energy thus liberated once again becomes thermal energy in the ball, as dictated by the Virial Theorem. With the helium ball now getting gradually hotter, the overlying hydrogen-fusing shell becomes gradually hotter also, producing in turn a gradual speeding-up of its hydrogen fusion, and therefore a gradual increase in the star's (already high) luminosity.

RGB life comes to an end with one of two possible kinds of transition to core helium fusion, both entailing a decrease in overall luminosity and yet without much change in photosphere temperature. The transition is violent in the case of the less-massive stars in our set, less violent in the case of the more-massive stars in our set: we again omit details. The core-helium-fusion phase is analogous to, and yet is briefer than, the core-hydrogen fusion that characterizes the earlier part of the theoretical MS. The exact destination of this transition depends on whether the star was at the time of its birth (its arrival on the theoretical MS) metal-poor or metal-rich.

For a star born as metal-poor, exit from the RGB takes it rapidly to the “theoretical HB.” This region of the theoretical luminosity-vs-photosphere-effective-temperature plot corresponds to a long, roughly horizontal, roughly straight locus of points, which we might term the “observational HB,” on the MK surface. Since globular clusters are metal poor, the observational HB becomes prominent when a globular is (at least partly) resolved into its constituent stars, for which spectroscopy then yields individual MK types. Different metal-poor stars switching on their core-helium fusion are found to arrive at different points on the observational HB, i.e. to attain different photosphere effective temperatures. The particular attained photosphere effective temperature is found in computer modelling to depend chiefly not on the mass of the newly ignited helium core (this proves on modelling to be rather constant across the metal-poor population), but on the mass of the outer, non-helium, layers.

However, with just two or three or so known exceptions—the most celebrated of these being  $\alpha$  Boo (Arcturus)—our 325 MK-classified bright stars are metal-rich. Moreover, the exceptions in our set of 325 are perhaps all at phases of evolution either preceding or following residency on the theoretical HB and observational HB. We will therefore not discuss the HB further.

For a star born as metal rich, exit from the RGB, i.e. the switching on of core-helium fusion, involves a rapid transition to the theoretical and observational “Red Clump” (in effect the redmost rump of the grander theoretical and observational HB), as further discussed at, e.g. [en.wikipedia.org/wiki/Red\\_clump](https://en.wikipedia.org/wiki/Red_clump). Since the Red Clump is the helium-fusion analogue of theoretical-MS core-hydrogen fusion, it is unsurprising that it is followed by an evolutionary phase analogous to the observational RGB and theoretical RGB, namely the observational AGB and theoretical AGB.

On the AGB, helium-core fusion has come to an end, with the star at this late stage in its life harbouring an inert core rich in carbon and oxygen. Fusion now proceeds, simultaneously or alternately, in an inner shell of helium and an overlying (and in terms of overall luminous output, for most of the AGB lifetime dominant) shell of hydrogen. With more than one shell in play, evolution becomes rather elaborate. In particular, it is possible for the helium shell to be temporarily inactive, simply accreting mass from the helium ash being dumped on it by the overlying hydrogen shell. Once the helium shell becomes sufficiently massive, it turns on helium fusion, causing the overlying hydrogen shell to expand and briefly switch off. The net result of this is a temporary drop in the luminosity of the star, until the helium burning in turn subsides and the hydrogen burning resumes. In its overall evolution along the AGB, and in its post-AGB transition to the quiet, dead state of a white dwarf, a star can undergo even many tens of such “helium shell flash” episodes. Additionally characteristic of evolution on the AGB are pulsation and mass loss. The possibility is dramatically illustrated in our 325-star set by  $\alpha$  Her Aa (Rasalgethi), and with a still higher mass loss by  $\omicron$  (omicron) Cet Aa (Mira).

We will skip over the further details of stellar evolution toward the white-dwarf corpse phase, remarking here only that in the case of a star nearly, but not quite, massive enough to die as a supernova, even carbon may be fused before all thermonuclear activity finally ceases.

Two concluding remarks are now in order.

(1) Mention has already been made of “Dredge-Up” as a process affecting the elemental composition of the spectroscopically observed photosphere. In terms of the concepts now laid out, it can be added that “Dredge-Up” may occur in the violent and deep convection of the RGB, as “First Dredge-Up” (FDU), or after the RGB, as “Second Dredge-Up” (SDU) and “Third Dredge-Up” (TDU). A highly evolved star may experience more than one episode of TDU, and it is also possible for FDU and TDU to occur without SDU. Our table cites

$\alpha$  Tau A (Aldebaran) as a star that has undergone FDU. On the other hand, our table in its present state of development does not cite instances of SDU or TDU.

(2) The deducing of a star's evolutionary stage from its observed MK type, as it makes its way off the MS toward, eventually, the AGB, is not always straightforward. In the case of the most massive stars (with masses greater than  $\sim 8 M_{\odot}$  or  $\sim 10 M_{\odot}$ , and with death-by-supernova therefore impending, and with temperature evolution late in life at one or more stages proceeding in the sense OBAFGKM and at one or more stages proceeding in the contrary sense MKGFABO), temporary observed residence, as a Cepheid variable, on the Instability Strip (IS) raises the question (not always easy to answer) "Is this star making a first, a second, or a third crossing of the IS?" As pointed out in the table "Remarks," this problem complicates, in particular, the analysis of that rather untidy Cepheid variable that is  $\alpha$  UMi Aa (Polaris). For those stars massive enough to achieve core-helium fusion at some point in their lives, and not so massive as to die a supernova death (a condition met in our 325-star set by all the stars below  $\sim 8 M_{\odot}$  or  $\sim 10 M_{\odot}$ ), it sometimes proves difficult to distinguish residency on the theoretical RGB, residency in the theoretical Red Clump, and residency in the theoretical AGB from the available spectroscopy. Indeed the theoretical "Asymptotic Giant Branch" is so named because it corresponds in observational terms to a locus of MK-surface points running perilously close to, so-to-speak, asymptotically approaching, that just slightly redder locus of points that is the observational RGB.

### **5.9: Supplementary remarks on rotation with "Be phenomenon" and "shell" (in MK types O, B, A)**

Some of our 325 bright MK-classified B stars have an "e" flag, for emission lines in spectroscopy. Some, and yet not all, such cases involve the important, and not yet well understood, "Be phenomenon." Strictly speaking, the presently known "Be-phenomenon" stars in our set of 325 are at least the following 19 (in order of increasing RA):  $\gamma$  Cas A,  $\alpha$  Eri (Achernar),  $\varepsilon$  Cas,  $\eta$  Tau Aa (Alcyone),  $\eta$  Ori Aa,  $\zeta$  Tau (Tianguan),  $\alpha$  Col A (Phact),  $\kappa$  CMa,  $\beta$  CMi A (Gomeisa),  $\omega$  Car,  $\rho$  Car (HR4140),  $\gamma$  UMa A (Phecda),  $\delta$  Cen Aa,  $\mu$  Cen Aa,  $\eta$  Cen,  $\delta$  Sco A (Dschubba),  $\alpha$  Ara A,  $\zeta$  Oph, and  $\beta$  Cep Aa (Alfirk). As we discuss again below,  $\beta$  Lyr Aa1 (Sheliak) may or may not constitute a twentieth case, and some doubt hangs additionally over  $\gamma$  Ara A (in our treatment, not a "Be phenomenon" star, because too evolved; but perhaps we are wrong). Further, closely related to the Be phenomenon is the spectroscopic (predominantly B-star) "shell" phenomenon. The amateur-spectroscopy essay in the Handbook current printed editions notes that the spectroscopic-shell phenomenon, and by implication the Be phenomenon, is a potentially fertile field for amateur spectroscopy. We accordingly supply here a general briefing on the Be phenomenon and its "shell" associate, highlighting the connection of both Be and shell with the often-troubling topic of rotation.

Although many of the most tempting amateur targets in the Be-phenomenon and "shell" fields are members of our 325-star set, we nevertheless discuss the Be and shell phenomena for the most part in general terms, without restriction to the set of 325. We hope thereby to maximize the value of our discussion, and in particular to stimulate an interest in Pleione, as a Be and sometimes "shell spectrum" star not much fainter than our mag.  $\sim 3.55$  cutoff.

Of all the non-cluster B stars in the galaxy, about 17% at some point in their lives present the "Be phenomenon," with the phenomenon more prevalent at the hotter (near-O) than at the cooler (near-A) end of the B range. Within the overall set of galactic stars, the exceedingly rare O stars are known to sometimes present the same phenomenon (with the term "Oe" star therefore used occasionally in the literature). In our set of 325,  $\zeta$  Oph, as an O star with a photosphere almost, and yet not quite, cool enough to entail classification as a hot B, is an instance. Also within the overall set of galaxy stars, some A stars are known to present the Be phenomenon. Again, our 325-star set furnishes an instance, namely  $\gamma$  UMa A (Phecda): this star is of MK temperature type A0, and so is just barely cool enough not to fall into the B classification bin. Nevertheless, since the phenomenon (which we will soon describe in proper detail) occurs predominantly in the B stars, the term "Be phenomenon" is standardly applied to stars in all three of the O, B, and A observational MK temperature types.

Several qualifying comments are now necessary.

The Be-phenomenon stars are not to be confused with the "Herbig Ae/Be 'stars'." The latter are not stars in the strict sense, but instead are contracting starlike bodies that have not yet achieved starbirth, i.e. have not yet started core hydrogen fusion. In their present stage of development, they are continuing to heat up under gravitational contraction, and are (unsurprisingly for objects condensing out of ISM clouds) embedded in

circumstellar dust.

A true “Be phenomenon star” need not currently have emission lines in its spectrum. It must, on the other hand, be known to have at some point in its past presented emission. In observational practice, the emission is always found to occur in at least one or more lines of the hydrogen Balmer series.

The condition of past-or-present emission, while necessary, is not in its turn sufficient. A supergiant in MK type B, with Balmer emission, is not a Be-phenomenon star. For a star to be Be-phenomenon, it must either lie on the theoretical MS or (as in the case of Be-phenomenon  $\zeta$  Tau (Tianguan) in our table, observationally in MK luminosity class III) be evolved only modestly beyond the theoretical MS.

Also not harbouring a Be-phenomenon star is a theoretical-MS or near-theoretical-MS member of a binary system with mass transfer, in which the observed hydrogen Balmer emission comes from an incandescent mass-transfer stream. In the table, this is perhaps the case for  $\beta$  Lyr Aa1 (Sheliak), which certainly has such a mass-transfer stream. Confusingly, however, a “shell” spectrum is observed for Sheliak, and “shell” in the case of a young B star (as we explain below) is generally, or even inevitably, associated with the Be phenomenon. Perhaps all that can be said here is that Sheliak is a confusing case. (It has certainly been notorious over the decades, in one way or another, as a challenge to modelling.) The conceptual point remains that if, hypothetically speaking, emission in a young B star were to come from no source other than a mass-transfer stream, thanks to that star’s membership in a tight binary system, then that star, while being obliged to show the observation-driven “e” flag in its MK type, would not count as an instance of the Be phenomenon.

This, then, concludes the qualifying comments. To recapitulate: the true Be-phenomenon stars are theoretical-MS or near-theoretical-MS stars with presently observed or historically observed emission lines, where the emission is not due to a mere mass-transfer process attributable to membership in a tight binary system.

The astrophysical task is now to determine what produces the emission. Emission must mean that the star has somehow managed to shed significant quantities of incandescent gas. Copious shedding cannot be attributed to stellar winds, since winds play only a minor role in mass-shedding for stars within or near the MS (except, perhaps, for the case of stars at the hottest end of the O range, where even the concept of time-spent-on-theoretical-MS is, as noted above, problematic). Our own Sun, for instance, as an MS star, sheds a mere tenth-of-a-trillionth of its mass per year.

The cause of the copious shedding has not yet been determined with confidence. It is possible that all Be stars are rapid rotators (although, as we remarked in Subsection 4.4, spectroscopy, with its incorporation of “n” or “nn,” as occasionally appropriate, in an MK type, cannot by itself detect rotation when the star is oriented pole-on to Earth). On the other hand, there are many rapidly rotating theoretical-MS or near-theoretical-MS O, B, and A stars that do not present the Be phenomenon.

The following picture therefore suggests itself: if the star is a rapid rotator, and in addition possesses some mechanism “X” for launching photosphere gas from near its equator into its equatorial plane, then an incandescent disk forms, girdling the star, and registering as emission at the spectrograph. With the star a rapid rotator, it will not be a sphere but a rotationally somewhat flattened object, with local gravity in the photosphere somewhat lower at the equator than at the poles, and with launching into an equator-plane orbit consequently favoured. The observed hydrogen Balmer emission is in this picture a signature of hydrogen ionization in the disk, under a violent barrage of UV from the (hot, as O-or-B-or-A) photosphere: Balmer-lines hydrogen light is emitted as part of the process in which free electrons and hydrogen nuclei recombine, where a captured electron falls to the penultimate energy level from some higher level.

The equatorial-disk picture was first proposed in 1931. Now quite widely accepted is a “Viscous Decretion Disk” elaboration of this idea, introduced in [1991MNRAS.250.432L](#). “Decretion” proves a useful contrived astronomical term, created as an antonym for “accretion.” Accretion disks figure in various astrophysics contexts, for instance in such black-hole binaries as Cyg X-1 (material shed by the readily amateur-visible member of this binary falls first onto an accretion disk around the black-hole event horizon), and again in the case of starbirth, where material from the gestating ISM cloud forms an accretion disk around the protostar, in a process that might see the disk eventually transform itself into a bevy of exoplanets, with perhaps also a belt of small rocky asteroid-like bodies, and with some analogue of our Solar System’s zodiacal dust, all orbiting an infant star. Correspondingly, a “decretion disk” forms when an astronomical object (in our case the Be-phenomenon star) for one reason or another releases matter into orbit in its neighbourhood.

Although the dimensions of the hypothesized disk are not easily investigated, emission in the Balmer

hydrogen- $\alpha$  line in the cases so far studied has been found to come from a disk on the order of 0.3 au to 0.6 au in radius. We seem to have here, in other words, one of the grandest of all theoretical-MS or near-theoretical-MS stellar spectacles.

Unfortunately, it is a spectacle that at best can be imaged only fuzzily, even with the most capable current optical interferometers. Let the Jupiter disk, of diameter  $\sim 50''$ , familiar from the small telescope, become a circular tea-tray 50 cm in diameter. The binaries resolvable in good seeing by the small telescope, at a separation of  $\sim 1''$ , thereby become a pair of points on that tray lying 1 cm apart. The most celebrated of the Be-phenomenon stars,  $\gamma$  Cas A, already noted as spectroscopically peculiar by the first stellar spectroscopist, Fr Angelo Secchi, in or shortly before 1866, lies at a distance of 600 ly from Earth. A disk of incandescent gas on the order of 0.5 au in radius, or 1 au in diameter, is seen at this distance as an object a mere 5 mas across. In terms of the tea-tray, this corresponds to an object around 50 microns wide, in other words to an object having the approximate width of a human hair. Consistent with the picture of gases launched by “Mechanism X” into circumstellar orbit is the [2007A&A...464...59M](#) discovery that the gas in Be-phenomenon star  $\alpha$  Ara A is in a normal central-gravitational-field (i.e. Keplerian) orbit, moving unconstrained by any such nongravitational forces as magnetism, and not possessing the kinetics of a mere stellar wind.

What, then, can “Mechanism X” be? It is possible that different Be-phenomenon stars have different gas-launching mechanisms. Outflows from the poles are not currently considered relevant. Nonradial pulsation, on the other hand, may play a role in at least some cases, as may also local magnetic phenomena at the low latitudes. (There is perhaps no known case of a Be-phenomenon star with a strong global magnetic field.) Helpfully, all hitherto scrutinized Be-phenomenon stars have been found to be pulsating variables, although in some cases the pulsation-produced luminosity variation is at the millimagnitude level or below, eluding detection by ground-based photometry. (In addition to facing possible very-low-amplitude variations, photometric monitoring of the stellar pulsation is confronted by the complication that the disk itself may vary photometrically (possibly with high amplitude).)

Nonradial pulsation aside, it is possible that in some cases, where the Be star is a member of a binary with tight orbit, or at any rate with an orbit possessing a tight periastron, the “X” role is played by the perturbing gravitational field of the companion.

Some Be-phenomenon stars have emission (from, on the currently accepted modelling, equatorial disks) which is, so far as the existing multidecade observational record goes, stable. Other Be-phenomenon stars, however, present emission lines only intermittently, in their years or decades of “outburst.” Two prominent instances of outburst-and-quiescence in our 325-star set are the already-cited  $\gamma$  Cas A and the recently active  $\delta$  Sco A (Dschubba). Another well-known instance, although a little too dim for inclusion in the 325-star set, and sharing the notoriety of bright  $\gamma$  Cas A, is Pleione. This star, easy in binoculars as the northern neighbour of Atlas at the eastern extremity of the Pleiades, presented an emission-line outburst of uncertain commencement extending to 1903, and presented additional emission-line outbursts in the periods 1955–1972 and 1989–2005.

Where the disk is permanent, the “X” mechanism works steadily to launch fresh consignments of photospheric gas into orbit, i.e. to perpetuate the decretion. The ongoing launch compensates for the ongoing accretion of matter from at least the inner part of the disk back onto the photosphere. If the mechanism should for some reason cease to operate, decretion ceases, and yet accretion continues. This has the consequence that the disk vanishes (with, however, some of the outlying parts of the disk lost not to accretion onto the photosphere but to outflow, into the embedding ISM).

On some current modelling, a typical Be disk increases in thickness rather gently as one progresses outward (with radially directed tangents to the disk, as taken at the points where disk meets photosphere, yielding a tight “full-opening angle” of  $\sim 10^\circ$ ). A further geometrical detail from some current modelling may also be noted: if, as is often the case, the Be-phenomenon star is a member of a binary not tight enough to produce mass transfer, and yet tight enough to produce a gravitational perturbation from the companion star, and if the Be-phenomenon star equatorial plane diverges somewhat from the orbital plane of the binary system, then the disk is warped.

We may now turn from Be to the related “shell spectrum” phenomenon. The term is somewhat unfortunate, being perhaps a relic from discussions in the early 20th century, when it was perhaps thought that an O or B or A star in or near luminosity class V could, under the right circumstances, surround itself not with an equatorial disk of gaseous ejecta (as on the currently accepted modelling) but with a literal “shell” of gaseous ejecta, in other words with an enclosing blanket. For better or worse, the term has stuck, surviving the acceptance of the

disk morphology (and has nothing to do with thermonuclear-fusion shells in stellar interiors, as discussed in subsections 5.6, 5.7, and 5.8 of this essay). A “shell” spectrum in a rapid rotator, oriented equator-on to Earth, occurs when some lines are seen not in the expected broadened absorption typical of an equator-on rapid photosphere, but in, or also in, narrow absorption. Typically, though not inevitably, the unexpected narrow absorption lines occur as narrow absorption cores within Balmer emission.

On the current understanding, “shell” in this sense typically results when a Be star not only generates its (perhaps temporary) disk of equatorial ejecta but happens to be oriented more or less equator-on in relation to the spectrograph. Under these circumstances, part of the disk lies between photosphere and spectrograph, yielding the absorption. Since this part of the disk is moving more or less orthogonally to the line of sight, i.e. is neither approaching the spectrograph nor receding, its absorption lines escape the rotational broadening characteristic of absorption lines from the photosphere.

Although a Be-phenomenon star with equator-on orientation can, as just noted, be simultaneously in emission-line outburst and in “shell,” it sometimes happens that shell absorption is present in a Be-phenomenon star even after its emission has for the time being subsided. The Be-phenomenon star Pleione, in particular, had a shell spectrum without emission in the period 1938–1954, and then again for some years after 1973.

Rapid rotators fitting the definition of “shell spectrum” occur even somewhat outside our present domain of interest, the Be-phenomenon stars, with instances known even in type F, right down to the F5 “rotation break.” It remains the case, however, that “shell” is most prominently connected with the Be, as a phenomenon contemporaneous with a Be outburst or present in a star that at some earlier or later time is observed to be in Be outburst.

What, in this general Be-cum-“shell” field, are the possible lines of activity for the amateur spectroscopist?

On the humblest level (even with a visual spectroscope and no camera, as in the case of 1860’s Fr Angelo Secchi), it is possible to monitor theoretical-MS or near-theoretical-MS rapid rotators, to see whether emission is currently present or currently absent. The sudden onset of emission would be newsworthy of communication to AAVSO, to the LESIA laboratory at Paris-Meudon (as mentioned again below), or to other appropriate program authorities.

On a less humble level, where spectrograms are taken, and are converted into intensity-against-wavelength plots, or “extracted one-dimensional spectra,” with such professional astrophysical tools as IRAF, the evolution of emission-line and shell-absorption-line profiles could be tracked. In particular, where shell absorption is present simultaneously with emission, as in the (conveniently strong) hydrogen Balmer lines, duly equipped amateurs could examine from month to month whether emission is currently stronger on the violet, or on the contrary red side of the partitioning absorption.

Finally, we suggest in a speculative spirit that it might prove possible to keep a month-upon-month polarimetry log (although we do not ourselves know whether any amateurs in any country have attempted polarimetry, whether in a Be-phenomenon context or in other contexts): if the Be-phenomenon star is not seen pole-on, then some light from its photosphere will be scattered toward the polarimeter by free electrons in the disk and will therefore be linearly polarized.

The recent literature includes a long review article, [2013A&ARv..21...69R](#), on the Be phenomenon. The IAU Working Group on Active B Stars (a group whose domain of interest includes, and yet is not confined to, the Be and shell phenomena) has a homepage at [activebestars.iag.usp.br/bstars](#), with a link to its newsletter materials, including a newsletter archive. The LESIA laboratory at the Observatoire de Paris-Meudon maintains the “BeSS Database” comprising Be-phenomenon stars, the Herbig Ae/Be “stars” briefly mentioned near the beginning of this subsection, and a “B[e]” category of supergiants, at [basebe.obspm.fr/basebe](#).

## **SECTION 6: Supplementary user guide, concerning the treatment of photometric variability and photometric non-variability in our “Remarks” column**

### **6.1: Preliminary remarks concerning photometric variability and photometric non-variability**

#### **6.1.1: Temporal and amplitudinal thresholds for photometric (V-band) variability:**

We confine the entirety of this “Section 6” photometry discussion to measurements (i) in the Johnson-Morgan V passband (the central portion of the UBV, or “ultraviolet-blue-visual,” system introduced in [1953ApJ...117..313J](#)), and (ii) in two reasonable approximations for V.



The 1953 three-band UBVR system (extended in later years in several ways, notably to a five-passband UBVR system) is in two respects a large concession to pragmatism.

First, the “U,” or near-ultraviolet, passband has in practice a short-wavelength cutoff determined by local atmospheric conditions. The air over different ground-based observatories absorbs ultraviolet to different degrees, and even the constancy of ultraviolet absorption at any one site is not guaranteed, at any rate not over spans of years and decades.

Second, the generous width of the passbands makes this photometric system a notably coarse approximation to the (admittedly unattainable) photometric ideal of spectrophotometry, in which the incoming flux from a star would be plotted, not even in mere magnitudes, but in laboratory units of joules-per-square-metre or ergs-per-square-centimetre. On this ideal, flux would be taken in exceedingly narrow wavelength bins, say of width 1 pm (0.01 Å), as a full “Spectral Energy Distribution” (SED) histogram all the way from the vanishingly faint gamma-ray tail to the vanishingly faint radio-wave tail. (If the ideal were by some miracle to be attained, astrophysical benefits would accrue: one could read off the effective temperature of a given star’s photosphere merely by finding the Wien peak of the black-body curve that best fits the histogram curve. Further, by comparing the flux, as the area under the full SED curve, with the flux that is the definite integral, from negligible-gamma to negligible-radio, of the black-body curve, one could directly measure the black-body-perturbing effect of spectral absorption lines and spectral emission lines.)

Narrow-band photometric systems, the best known of which is Strömgren, provide a somewhat better approximation to the SED ideal. This pair of pragmatic concessions notwithstanding, Johnson-Morgan UBVR possesses several features that have earned it a wide following from the 1950s up to the present.

Firstly, the readily measurable difference of U and B (the “U-minus-B colour”) allows one to predict the extent of the Balmer-discontinuity jump that would be found in the more laborious procedure of taking a spectrogram centred on the Balmer-limit wavelength (in the near ultraviolet) of 364.6 nm. The extent of the jump is in turn a useful indicator, at any rate for stars redder than (cooler than) MK temperature types O and B, of photosphere density, and so supplies for stars cooler than MK types O and B a usable estimate or indication of their MK V, IV, III, II, or I luminosity class.

Again, the readily measurable B-minus-V colour allows photometry to predict the MK temperature type that would be found if the star were to undergo the more laborious procedures of spectroscopy. UBVR photometry can at the present time be performed with a basic thermocouple-cooled astronomical CCD (as a sufficiently exact substitute for the RCA 1P21 photomultiplier tube presupposed by the UBVR definition in [1953ApJ...117..313J](#)) even on a budget of a few thousands of CAD (or USD, or EUR), with even such a modest telescope aperture as 0.3 m or 0.2 m.

Finally, the generous width of the passbands helps ensure that UBVR photometry at any given aperture, with a CCD camera of any given sensitivity and noise level at an observatory site of any given constant quality, can push its way to fainter stars than would be feasible for a spectrograph under the same conditions. The following points from within the present writer’s experience, although merely anecdotal, are nevertheless suggestive of the practical advantages of photometry, at any rate in a situation where only a coarse indication of MK temperature type and MK luminosity class is needed:

- When Canada’s David Dunlap Observatory (DDO) was in the final (2007-era) phases of its research-grade spectroscopy, a professional-grade liquid-nitrogen-cooled camera was operated under light-polluted suburban skies at the principal (1.88 m) DDO telescope. It was at this point considered feasible to obtain usable spectrograms from stars of mag. 13 or so. On the other hand, the present writer does not recall, from a couple of years of DDO warm-room operations, taking usable spectrograms from stars as faint as mag. 15.
- Of the three telescopes operated at Tartu Observatory (in northeastern Europe, under rural skies with two horizon-hugging urban light-pollution incursions), the smaller two are used principally for photometry. Both are considered capable of yielding useful UBVR data at even mag. ~18, with in both cases thermocouple-cooled (not liquid-nitrogen-cooled) CCDs. One has 0.6 m of aperture and is equipped with traditional glass UBVR-system filters. The other has just 0.3 m of aperture but compensates for its restricted light-grab by using interference filters in place of glass. This choice of filter technology is found to make about the same faint photometric targets feasible in its case as are

found feasible for the 0.6-m telescope. Here, then, is a mag.  $\sim 18$  photometric capability, supporting among other things the estimation of MK two-dimensional spectral luminosity-versus-temperature types, from an aperture so small as to procure just one stellar photon for every  $\sim 40$  procured at DDO.

From the wide realm of UBV, we are here—we repeat—confining ourselves to V and to two of its reasonable approximations. The pair of approximations is of importance, given the frequent need to examine the photometric behaviour of stars over several decades, including from decades prior to the 1950s-onward implementation of UBV. In examining, for example, the historical record for Mira, one might first input “Mira” at the “Pick a Star” interface of [www.aavso.org](http://www.aavso.org), selecting “Plot a light curve.” This yields a light curve covering just the last couple of years. Upon clicking, however, within the displayed plot, on “Plot Another Curve,” and now requesting a plot of Mira from 1850 January 1 to the present, one gets a vast historical record, the bulk of which antedates the 1950s. For the interpretation of what is displayed, it is useful to have some knowledge of the pair of V-passband approximations to which we now proceed.

(a) Visual-estimate approximations to the V passband, in the best case good to  $\pm 0.1$  mag. as measures within their own (roughly-V) passband, have been made with the unaided eye at the telescope eyepiece since the days of Friedrich Wilhelm August Argelander. His first *Bonner Durchmusterung* volume appeared in 1859. Perhaps at a similar accuracy was also the pioneering 1850’s work of von Seidel. Additionally, as an improvement on the mere eye-at-eyepiece, various types of comparative photometer, notably the Zöllner, contributed to 19th-century photometry between the advent of the *Bonner Durchmusterung* and the introduction of the first (“pg-magnitude”) photographic plates.

(b) The V passband was approximated before the 1953 Johnson-Morgan definition with the “photovisual” photographic emulsions, as 20th-century “pv” magnitudes. (More primitive emulsions, with useful sensitivity only at the blue end of the spectrum and brought into significant observatory use from the 1880s onward, instead yielded the “pg,” or “photographic,” magnitudes. These early measurements do not approximate today’s V passband as well as their photographic-plate successors, the pv magnitudes.)

With the V passband and its approximations duly highlighted, a conceptual question arises: how short an interval of V-passband or V-approximation passband photometric constancy suffices for a star to be considered “not a variable star”? Or equivalently: how rapid is a photometric change required to be for a star to count as (at least sluggishly) variable?

All stars evolve over the scale of at any rate gigayears, with evolution in some cases producing marked photometric changes already over the scale of megayears or kiloyears. Nobody, however, would want to take this as a reason for calling all stars “variables.”

One might here be tempted to draw a conceptual distinction, within the special realm of the “intrinsic” variables (the special realm is examined in sub-subsection 6.2.1 below), as follows: (A) On the one hand within this realm are the intrinsic-variable stars whose photometric variations correspond directly to an advance along an evolutionary track. Such a star would count as “intrinsically varying, and yet not momentarily a variable.” An instance of this would be a star that, even while remaining within the theoretical and MK-phenomenological MS, while stably burning core hydrogen, steadily brightens over a period of megayears, as its core becomes progressively more helium-rich, progressively denser, and progressively hotter. In this gradual evolutionary advance, the steady temperature rise favours a steadily increasing energy output, as the highly productive hydrogen-to-helium CNO process becomes progressively more favoured over the less productive hydrogen-to-helium “pp chain” process.

(B) On the other hand within this realm are the intrinsic variable stars whose photometric variations are rapid enough to be discernible even within one single stage of stellar evolution. Instances of “(B)” would be furnished by the Cepheids, whose photometric fluctuations occur rapidly, at well-demarcated particular stages in their career, specifically when helium-shell burning repeatedly takes them to “blueward excursions” that happen to cross the Instability Strip (the IS) in two-dimensional MK classification space. Over a period of just a few cycles, the evolutionary stage of a Cepheid is for practical purposes fixed (even though evolutionary effects can make themselves felt, through tiny changes in pulsation period, over tens or hundreds of cycles—as is again remarked in sub-subsections 6.1.4 and 6.2.4, below).

Again, instances of “(B)” would be furnished by intermittently flaring MS stars, such as  $\alpha$  Cen C, destined to remain on the MS for many gigayears, and so almost at an evolutionary standstill across even the whole probable span of human history.

Appealing though this distinction might seem, the temptation to draw it is best resisted. A solar-mass star late in life, suffering a helium shell flash episode, as nested shells of helium and hydrogen have their thermonuclear fusion turn on and off (this situation is discussed in subsection 5.8 above), would have to be placed under heading “(A).” Awkwardly, such a star would have to go under heading “(A)” even if one of its evolutionary-process photometric transitions were to prove dramatically swift, consuming a mere kiloyear, or a mere year. Moreover, it would be unclear what to make of Wolf-Rayets (among the massive stars) and Miras (among the solar-mass stars). Although these two stellar types are universally considered variables, it might be argued that their episodes of mass shedding, with their consequences for V-passband light curves, are processes of (rapid) advance along the evolutionary track. This would, awkwardly and counterintuitively, force Wolf-Rayets and Miras to go under heading “(A).”

Having drawn attention to the conceptual problem, we resist, as we say, the temptation to offer it this particular solution. We suggest instead that “variability” is a mere pragmatic matter, not to be defined formally, and to be governed merely by the exigencies of given concrete photometry programs. A team studying  $\delta$  Sct-type variability (a type of rapid pulsation, discussed in detail in sub-subsection 6.2.4 below) might well conduct differential photometry of some  $\delta$  Sct star at a single observatory over a continuous period of 10 hours, or in the more ambitious case of the “Whole Earth Telescope” over a continuous period of 500 hours, with a more westerly observatory in that globe-spanning consortium coming onto the duty roster as daybreak forces a less westerly observatory to await its next nightfall. In differential photometry, a field is imaged of the target star, and additionally of a “comparison star” and (for monitoring both the stability of the “comparison” and the stability of the observatory equipment) a “check star.” If, in this research program, the “comparison” and “check” are found not to vary against each other, beyond the level of mere stable-instrument noise, over the duration of the observing run, it would be reasonable to call the comparison and the check “non-variable.” This situation contrasts with the situation of a team applying the same target star/ comparison star/ check star method to a slow Cepheid, pulsating with a roughly 10-day period. In the case of a Cepheid, observations might be taken not continuously, but merely a few times each night, over a run of six months. It would now in a pragmatic way be convenient to call the comparison star and the check “non-variable” if they are found not to vary against each other, above the mere stable-instrument noise level, over the duration of the six-month campaign.

We end this timeframes discussion by raising three AAVSO-specific questions that we may possibly hope to answer in later years, as this Handbook supplement goes through revisions:

- Over what timeframe is photometric constancy asserted at the (for many purposes authoritative) AAVSO(VSX) database when a star is flagged as not variable? (This is the flagging performed with the grey “N” symbol, in the AAVSO(VSX) Web interface—with the grey “N,” as distinct from the red “S,” which marks a suspected variable, and as distinct from the green “V,” which marks a confirmed variable.)
- Over what timeframe is photometric constancy asserted at AAVSO(VSX) when a star not only receives the grey “N” status flag but additionally is given the “CST” variability classification symbol (as distinct, for instance, from a star receiving the green “V” status flag and the classical-Cepheid “DCEP” variability classification symbol, or again as distinct from a star receiving the green “V” status flag and the eruptive-variable (UV Cet-type) “UV” variability classification symbol)?
- Over what timeframe is photometric constancy asserted at AAVSO (outside the specific ambit of the AAVSO database that is AAVSO(VSX), when a star is placed into one of the non-variable “standard fields,” from the work of Landolt and Henden, announced at [app.aavso.org/vsd/stdfields](http://app.aavso.org/vsd/stdfields)? (An initial glance at a foundational Landolt paper, [1992AJ....104..340L](#), suggests to this writer that a reasonable answer might be “constant over ~10 nights,” as distinct from “constant over ~100 nights”: is this correct?)

A second, more obvious, conceptual question arises also. How small can the overall (V passband or approximate-V passband) photometric change be, over whatever temporal span has for whatever (perhaps rather pragmatic) reason been adopted, before the specimen under study is deemed to be “not a variable”? Here the question is not, so to speak, “How sluggish is the swing in a variable permitted to be?” but, rather, “How low is

the permitted swing amplitude?” Some perhaps as yet unknown, but large, percentage of stars would prove variable in the V passband over any reasonable chosen fixed temporal span if measurements could be made at the micromagnitude level. Spectroscopy of the Sun, with the slit directed at various different portions of the 31- or 32-arcminute apparent disk, reveals multiple localized oscillations with periods of a few minutes. It would therefore be unsurprising to find some corresponding V-passband fluctuation, with a similar short period, in the micromagnitude range, if someone were at each instant in the interval to integrate the total V-passband flux from the entire apparent disk. Further, a major development in 21st-century photometry has been the introduction of stellar photometry from space, at the micromagnitude level (pioneered by the Canadian 2003–2019 MOST nanosatellite mission; more recent work includes BRITe, and micromagnitude stellar photometry is a useful by-product also of exoplanet missions, such as CoRoT at ESA (2006–2013), and Kepler (2009–2018) at NASA and TESS (2018–present) at NASA). This Handbook supplement, however, will confine itself to variability as in practice handled in most of the AAVSO(VSX) database, in other words to variability at a level on the order of 10 millimagitudes.

Such variability is within the range of even modest amateur electronic equipment, notably of an off-the-shelf astronomical CCD camera set up to image a star field wide enough to contain the target star, a comparison star, and a check star.

With V-passband variability thus characterized, in particular possible to answer, in the affirmative, a question liable to arise at public-outreach events, for instance at events offered by RASC: “Is the Sun variable?” The Sun varies in AAVSO-relevant terms over its 22-year magnetic, or 11-year sunspot, cycle, quite apart from the possible (sub-AAVSO) micromagnitude variability previously mentioned. An observatory equipped with an off-the-shelf astronomical CCD and working some tens or hundreds of light-years outside the Solar System would detect tiny V-passband changes in the Sun from week to week, or at the very worst from quinquennium to quinquennium, at any rate outside such quiet-sun (low-sunspot-number) decades as the roughly 1645-through-1715 Maunder Minimum. This hypothetical observatory would note a clear regularity, although less strict than can be noted over ten days (even with the naked eye) in the case of pulsating  $\eta$  Aql or eclipsing  $\beta$  Lyr. The hypothetical observatory would detect V-passband maxima of approximately equal height, and V-passband minima of rather more varying depth, with an overall swing on the order of 10 millimag from maximum to minimum. So yes (one can say from the podium on RASC public-lecture occasions): V-passband variability, in the familiar AAVSO sense, is present within a not-too-protracted timeframe in even a star as placid as the Sun.

### **6.1.2: Stars versus multiple-star systems in the AAVSO(VSX) taxonomy of photometric variabilities:**

This Handbook supplement follows the usual [notably, the AAVSO(VSX)] practice of working in most cases with entire binary, or indeed with entire nested-binaries-hierarchy, systems, rather than with individual stars. Rare exceptions can arise where a binary features an angular separation so wide as to make it natural and easy, even at the level of binoculars, to work with individual components. It is also conceivable that some such exceptions, in other words some cases in which the literature has become accustomed to discussing individual components in a binary or a nested-binaries hierarchy, arise in this supplement for merely historical reasons. (Is it perhaps for merely historical reasons that photometry references discuss separately  $\alpha$  Cen A (Rigel Kentaurus) and  $\alpha$  Cen B (Toliman)?) This supplement consequently discusses such things as “ $\beta$  Per,” in place of such things as “Algol” (treating “ $\beta$  Per” as the entire hierarchically organized system within which the now rigorously IAU-named Algol, in other words the star that in rigorous WDS nomenclature becomes  $\beta$  Per Aa1, is a component: the gross, naked-eye, photometric  $\beta$  Per dips are due to eclipses of hot, small  $\beta$  Per Aa1 by large, cool  $\beta$  Per Aa2, and conversely the small CCD-detectable photometric dips are due to the transits of hot, small  $\beta$  Per Aa1 across the face of large, cool  $\beta$  Per Aa2).

The focus on systems, as opposed to individual stars, makes it easy not only to discuss cases in which the photometric variations are due to eclipses, but to discuss other binarity-involving cases also: for instance, where some difficult, perhaps only interferometrically resolved, binary, seen face-on and therefore not eclipsing (the two-star system “ $\omega$  F<sub>00</sub> Baris AB,” or “ $\omega$  FBr AB”) is found to harbour variability, and it is not clear whether the seat of the variability is  $\omega$  FBr A or  $\omega$  FBr B.

With all this said, we nevertheless allow ourselves the usual liberty of writing “variable star” in this Handbook supplement even where pedantry would require the phrasing “variability-harbours binary” or “variability-harbours hierarchical nested-binaries system.”

We also occasionally allow ourselves to write of variability in a “system” where what is likely the case really is variability on the part of what condensed from the gestating molecular cloud as a lone star, not as a binary.

Finally, we do not attempt to separate out any of the conceivable cases in which (i) what is traditionally entered into atlases merely as “ $\omega$  FBr” is found at high visual resolution to be the double “ $\omega$  FBr A” and “ $\omega$  FBr B,” and in which the AB pairing harbours variability (perhaps  $\omega$  FBr A is a flare star, and  $\omega$  FBr B is non-varying; perhaps, again,  $\omega$  FBr A is a flare star, and  $\omega$  FBr B is an unresolved eclipsing spectroscopic binary, destined some day to be resolved into “ $\omega$  FBr Ba” and “ $\omega$  FBr Bb”), and in which further (ii) as bad luck would have it, the pairing of  $\omega$  FBr A and  $\omega$  FBr B is a mere line-of-sight coincidence (meaning that there is in astrophysical reality no such thing as “the  $\omega$  FBr system,” even though there is such a thing as the (binary, rather than three-star) “ $\omega$  FBr B system.”

### **6.1.3: Taxonomy of photometric variables as less useful than taxonomy of photometric variabilities:**

It is possible for one and the same star, let alone for one and the same binary or nested-binaries system, to harbour more than one type of variability. A particularly striking instance of this possibility (although, admittedly, an instance falling outside Sample S) is some lone star, gravitationally unpaired with any companion, which simultaneously harbours BY Dra- and UV Cet-type variability. As noted in sub-subsection 6.1.7 below, a BY Dra variable has a severely inhomogeneous photosphere, and is a rotator, and is detected as variable when successive portions of the photosphere rotate into the view of the observatory, from one night to the next or from one week to the next. As noted in sub-subsection 6.1.9 below, on the other hand, a UV Cet-type variable has a violently active photosphere, with large flares. Since flares are driven by magnetism, and since magnetism in stars is a rotation-driven dynamo effect, it is not surprising to find the two types of variability co-occurring, at any rate in some (rather faint) stars outside Sample S.

A similar possibility, discussed again in sub-subsection 6.2.4 below, is the case of a star that at one and the same time presents both  $\delta$  Sct-type variability (in which pulsations are excited by the “kappa mechanism,” involving stellar-interior opacity changes) and  $\gamma$  Dor-type variability (in which pulsations are excited by convection).

We therefore have a situation reminiscent less of classic Linnean botanic taxonomy than of medicine. In botany, no plant that happens to be, say, a specimen of *Crocus vernus* is simultaneously considered a specimen of *Scilla sibirica*. Medicine, on the other hand, is obliged to accommodate “comorbidities”: one and the same patient may, *e.g.* simultaneously suffer both lung cancer and heart disease.

With all this said, we nevertheless do in this Handbook supplement allow ourselves the usual shorthand, in which one speaks of classifying “variables” (referring loosely, as already remarked in sub-subsection 6.1.2, to “variable stars” and “variable systems,” foregoing pedantry) when what is strictly meant is the classifying of “variabilities.”

### **6.1.4: Purely phenomenological taxonomy as less useful in photometry than in spectroscopy:**

In Section 5.1, MK spectral classification was discussed as a classification scheme at one and the same time based purely on observables and useful in astrophysics. In MK classification, the mere inspection of spectrograms, in the absence of astrophysical theorizing, is used to allocate a “temperature type” (typically some phenomenologically defined subtype of O, B, A, F, G, K, or M) and a “luminosity class” (typically one of the phenomenologically defined classifications V, IV, III, II, I). It is then found that stars occupying the same spot in the two-dimensional MK classification scheme are astrophysically similar, and that the more disparate two stars are in the scheme, the more astrophysically diverse they tend to be. In Section 5.1, a parallel was drawn with cardiac medicine, in which the diverse phenomenologies evident through the stethoscope are found to correspond well to diverse underlying physical conditions (the leaky valve, or alternatively the poor neuro-electrical signalling, or alternatively the aortic aneurysm).

With photometry, unfortunately, the situation is less favourable, calling to mind pre-modern efforts in fever medicine. One can, and physicians before the 19th century did, impose a classification based on observables, as when the three-day “tertian fevers” are distinguished from the four-day “quartan fevers.” However, it was noted in Section 5.1 that such a pure-phenomenology classification does not correspond in a useful way to the underlying physical realities of bacteria and viruses. In particular, cases that are in photometric terms rather similar can differ radically in their astrophysics. One might, for example, naively think that a regular alternation

of deep and shallow minima, in V-passband photometry, always has the same general underlying cause. Alternating deep and shallow minima are observed with many eclipsing systems (at the CCD-photometry level with  $\beta$  Per, as noted above, and at even the naked-eye level with that hierarchical system  $\beta$  Lyr—where the observed variation stems from the mutual motion of WDS-canonical  $\beta$  Lyr Aa1 (in IAU-canonical naming, Sheliak) and  $\beta$  Lyr Aa2, along with the motion of their connecting light-emitting mass stream). And yet this is not the exclusive preserve of eclipsing binaries: the light curves of RV Tau variables (outside Sample S), which are pulsators, and can occur outside a binary pairing, likewise feature alternating deep and shallow minima.

As a second example showing the inappropriateness of an MK-like phenomenological taxonomy in photometry, we note that two different underlying astrophysical causes can produce light curves with a clock-like regularity, each cycle lasting just a few days, and with the period changing by a few seconds over the span of many tens or a few hundreds of cycles. This light-curve phenomenology is generated on the one hand by some eclipsing binaries (with mass transfer driving a slight, and inexorably one-way, drift in period), and on the other hand by typical Cepheids (pulsators, in a subset of which a similarly inexorable tiny one-way year-upon-year change in period occurs: here the one-way drift is instead a consequence of changes in the stellar interior, as thermonuclear fuel reserves are progressively depleted). It is necessary, then, to forego the phenomenological approach successful with MK in spectroscopy, and to work instead on underlying causes. What is found to be important is not, so to speak, the number of days the fever lasts, but the nature of the underlying microbe.

### **6.1.5: Logical challenges in photometric taxonomy: (1) sets versus supersets; (2) essential (defining) features versus empirically salient features:**

We shall later, in our detailed examination of AAVSO(VSX) variable-star sub-subclasses, run up against two taxonomic problems arising repeatedly in natural science, even outside astronomy.

First, there is the danger of confusing sets with supersets (as would be comically the case if the curator of a taxidermy museum were to compile a catalogue with the entries “rodents, mammals, primates”). In variable-star work, it can be quite obvious where the intention is to demarcate sets and where the intention is to demarcate some superset. Clearly, for instance, for AAVSO(VSX) (and also for its leading input authority, Moscow-based GCVS) “E” denotes a superset, the general ensemble of eclipsing binaries, of several more narrowly defined groupings, such as the  $\beta$  Per-like (“Algol-like”) “EA” eclipsing binaries and the  $\beta$  Lyr-like “EB” eclipsing binaries. On occasion, however, the intention may not be obvious—as it will be necessary to remark in sub-subsection 6.2.4 below, in connection with the GCVS-and-AAVSO(VSX) Cepheid symbols “DCEP” and “DECEPS.”

Second, there is the danger of confusing features that are taken, at least provisionally, as pertaining with conceptual necessity to a group with features that are merely in empirical practice found salient (even invariably present) in the group. An instance, from outside astronomy, of this problem is furnished by zoology. How is one to define the domestic dog, in Linnean terminology *Canis familiaris*? In general, specimens of *Canis familiaris* are of a lighter bodily build than their interbreeding-capable genetic relatives the wolves, or *Canis lupus*, and are more amenable to life with humans than are wolves. These are not, however, conceptually essential features of the species. Feral dogs, perhaps even radically resistant to domestication, are observed in the field. Furthermore, *Canis familiaris* breeders can elicit massive, wolf-sized, body builds, as with what the American Kennel Club officially terms the “Irish Wolfhound.” Whether a given specimen belongs to *Canis familiaris* is ultimately a matter of underlying DNA-level causes, not necessarily well understood at this time. All that can be done is to start by picking out some paradigm specimens, decreed as clearly belonging to *Canis familiaris* (or as “clearly UV Cet-type flare stars,” or as “clearly  $\beta$  Cep pulsators”; or as “clearly basalts”; or again as “clearly cholera”; or whatever). One then has to seek for causes, as the “real essences” of the things being defined. In zoology, the “real essences” somehow involve genes. In geology, they involve chemistry, and additionally (an important consideration in the case of allomorphs) crystalline lattices. In the study of stars, they chiefly involve celestial mechanics and thermal physics.

It may well be that, as underlying causes become better studied, classification boundaries have to be shifted, with some provisional defining features even set aside, in a progressive distancing from mere “quartan versus tertian fever” taxonomy. What were once “fish,” formally “Pisces” (to the exclusion of whales) are now, with a better understanding of speciation (with Darwinian selection, over megayear and gigayear timespans), placed into a large so-called “clade” that does include the whales. This currently accepted clade contains no single Linnean grouping of “Pisces.” It does, on the other hand, contain multiple Linnean groupings for such things as

the cartilaginous fish, the armoured fish, and the bony fish, as well as various Linnean groupings of four-limbed land animals.

Although it is not necessary for Handbook supplement purposes to venture far into the conceptual analysis of taxonomy, a contextual remark at this stage may later help illuminate the various references later in this photometry section (in sub-subsections 6.2.2 and 6.2.4) to “real essence” problems. “Natural kinds,” or in more dramatic language “real essences,” repudiated in the 1930s by those parts of the philosophical community most closely engaged with natural science (notably by the “Logical Positivists”), are in favour once again, due to the 1970 “naming-and-necessity” analysis of philosophical logician Saul Kripke. His studies highlight the fact that taxonomic propositions emerge not as verbal deductions from clauses in a definition, but instead emerge in the course of empirical investigation, as truths that—surprisingly—succeed in being both a posteriori and conceptually necessary. Kripke remarks that while people have for millennia referred successfully (meaningfully) to gold, it is only in recent times that the real essence of gold has been identified, in other words that the “natural kind” that is gold has been adequately elucidated. A mere 18th-century definition, or quasi-definition, of gold might have specified that that substance is lustrous, yellow, malleable, and of high mass-per-unit-volume. Only in the 19th and 20th centuries did it become clear that gold is in its essence the 79th element in the periodic-table numbering, or still more adequately that gold is in its essence the element with 79 nuclear protons. The proposition that gold has 79 nuclear protons has, then, emerged as a truth that is a posteriori, and nevertheless is—surprisingly—necessary rather than contingent. If chemists or metallurgists were someday to find a way to prepare a brittle and charcoal-grey and low-density allotrope of the atomic-number-79 element, the product of their manipulations would still, as a matter of an only recently discovered definitional necessity, be gold.

#### **6.1.6: The “Silent Watchdog Problem” regarding photometric non-variability:**

The majority of stars in Sample S are either confirmed or suspected variables. It will, however, be seen toward the end of sub-subsection 6.2.7 that Sample S does contain a significant number of stars known to be in some reasonable sense not variable (as well as, frustratingly, a significant number of stars both not known to be in any reasonable sense variable and also not known to be in any reasonable sense not variable). Admittedly—but we will not examine this difficulty further—the problem of temporal thresholds for variability, inconclusively discussed in sub-subsection 6.1.1 above, becomes in a special way acute when a star or a binary system is classified as “known to be not variable.” (Has such a system been found V-passband constant over 1 year, or over 10 years, or over 50?) It is worth, then, keeping in view not only the logic-of-taxonomy challenges discussed in sub-subsection 6.1.5, but additionally what might be termed the “Silent Watchdog Problem.”

As with the pair of challenges from sub-subsection 6.1.5, here is in a sense a problem with a philosophical or logical aspect. The fact that a star fails to vary is significant, and consequently is a fact itself calling for a study of underlying causes. This is particularly the case with stars close in two-dimensional MK phenomenological space, and consequently also close in the two-dimensional total-energy-output-versus-effective-photosphere-temperature space of astrophysical theory, to known pulsators. Why, for instance, are there stars on or near that part of the IS which intersects with the phenomenological or theoretical MS, and which therefore ought to be  $\delta$  Sct-type pulsators (discussed in sub-subsection 6.2.4, below), which nevertheless have been found to be non-variable?

This problem is a special case of a more general problem, perhaps not yet solved: what determines the pulsation amplitude (large, small, or, as in the sub-subsection 6.2.4 concern just mentioned, zero) of a star residing on the IS?

This could be called, with reference to the Sherlock Holmes race-track tampering case, the “Silent Watchdog Problem.” In that Conan Doyle mystery story, a watchdog has failed to bark because the person approaching the stables in the night, intent on wounding a racehorse, was a trusted member of the dog’s own household:

- *‘Is there any point to which you would wish to draw my attention?’*

- *‘To the curious incident of the dog in the night-time.’*

- *‘The dog did nothing in the night-time.’*

-‘That was the curious incident,’ remarked Sherlock Holmes.

## **6.2: Overview of the AAVSO(VSX) photometric taxonomy**

### **6.2.1: Top-level classes, and their subclasses, in the AAVSO(VSX) taxonomy of V-band variabilities:**

In the AAVSO(VSX) taxonomy, variables are first divided into two broad classes. To these, and to their numerous subclasses, we within this particular Handbook supplement apply our own sequential letters, for enhanced readability. We reiterate now a point from sub-subsection 6.1.1 above, that we are confining ourselves to the V passband and its two good historical approximations (thereby skipping over some details at AAVSO(VSX), involving X-ray astronomy).

At the top of the AAVSO(VSX) taxonomic tree are the two broad classes of (A) extrinsic variables and (B) intrinsic variables, with the extrinsic class divided at AAVSO(VSX) into (A.a) variability-through-eclipsing and (A.b) variability-through-rotation. AAVSO(VSX) has also (A.c) variability through gravitational microlensing, even within the V passband. As might be expected, however, gravitational microlensing is absent from Sample S, since Sample S is drawn from the Sun’s own immediate galactic neighbourhood.

The guiding idea in extrinsic variability is that photometric fluctuations are due not to changes in a star, but to changes in the observatory’s view. Perhaps, for example, a binary system is seen nearly edge-on, with its stars consequently found to undergo mutual partial eclipses, or again total eclipses alternating with transits. Perhaps, again, a star that is for one or another reason photospherically inhomogeneous both (*a*) rotates and (*b*) is not seen strictly pole-on, and as a consequence of this pair of circumstances is found to present different longitude ranges to the observatory at different times. Such a star might resemble the Sun in being a spotted oblate spheroid. On the other hand, such a star might suffer an egg-like distortion, through being (*i*) in a binary system far enough from an edge-on orientation to Earth to present no eclipses or transits to the observatory, and on the other hand (*ii*) close enough to an edge-on orientation to ensure that different areas of its (far-from-spherical, even far-from-oblate-spheroid) photosphere present themselves to the observatory at different times. There is admittedly an element of arbitrariness here, with the possibility arising of an egg-shaped distortion that is not only presented to the observatory from different perspectives at different times (as could be the case even in a perfectly circular, adequately tight, orbit), but also (in a case of high orbital eccentricity) is intrinsically varying—negligible, perhaps, at apastron, and severe at periastron. Strict pedantry would in such a case require one to speak of something like “externally occasioned intrinsic variation.”

The intrinsic class of variables divides at AAVSO(VSX) into the subclasses of (B.a) pulsating, (B.b) eruptive, and (B.c) cataclysmic variables. In all these subclasses, it is the star, not the observatory perspective on some orbiting or spinning star, that undergoes the changes.

The pulsators do not require further examination at this stage. They are, however, examined at length, through a multitude of AAVSO(VSX) sub-subclasses of pulsational variability, in sub-subsection 6.2.4 below.

Eruptive variability involves gross photometric changes that stem not from pulsation (even though pulsation might also be present, as in medical terms a separately classifiable comorbidity), but instead stem from some single-star non-pulsational process. One possible such process is the formation of a bright equatorial “decretion disk,” from a hot star that is on or near the theoretical MS. (This is the “Be-phenomenon” case, discussed in subsection 5.9 above.) Another possible such process is a copious bright-matter outflow, from a star evolved far beyond the theoretical MS, with the formation of an outright sphere-like shroud (as with  $\eta$  Car, outside Sample S, and in a different way with the Wolf-Rayet (WR) stars, represented in Sample S by the WR component in the unresolved spectral binary that is  $\gamma$  Vel Aa). A third possible such process is flaring. A fourth is veiling by a dark carbon-rich outflow (as with a star much discussed in both amateur and professional astronomy, although too faint for inclusion in Sample S, R CrB).

The conceptual line between mere “eruptive” variability and outright “cataclysmic” variability, in other words between what are for readability in this Handbook supplement labelled “B.b” and “B.c,” perhaps defies rigorous characterization. In actual taxonomic practice, however, there would seem to be no ambiguous cases. Cataclysmic variability involves the recurrent novae (represented in Sample S by one system, T CrB), and additionally in AAVSO(VSX) work outside Sample S by various possibilities, of which the following four are particularly noteworthy: (*i*) the novae not yet observed to be recurrent; (*ii*) the “dwarf novae,” or U Gem-type variables; (*iii*) those exotic phenomena of recent professional study that are the contact-binary mergers; and (*iv*)



the supernovae. In all these instances of cataclysmic variability, some star is found to undergo a fundamental intrinsic change, more radical than the mere formation of a bright equatorial decretion disk or the formation of a bright or dark ejected-mass shroud.

We note, as a final comment on the subclasses, that the printed-edition RASC Handbook treatment of variability (constrained to be brief) diverges in two respects from the AAVSO(VSX) taxonomy that this Handbook supplement (unconstrained by page count) is following. First, the printed-edition treatment classifies rotational variability as intrinsic. This is to be defended as appropriate in a brief overview, in that a rotating star with photosphere patches, such as spots or large (Betelgeuse-type?) convection cells, is indeed in a simplified sense “doing something,” is indeed “in a process”: as time passes, there is at any rate a change in the flux reaching the observatory. Second, the printed treatment makes the cataclysmic variabilities a subclass of the extrinsic variabilities. This is to be defended as appropriate in a brief overview, in that the most common cases of cataclysmic variability are extrinsically occasioned instances of intrinsic variation: they occur in binaries, specifically in a scenario in which some white dwarf first accretes matter from outside—from its companion, through Roche-lobe overflow—and then reacts explosively to the intrusion.

### **6.2.2: The AAVSO(VSX) sub-subclasses of the eclipsing-variability subclass of the extrinsic variability class:**

It is now possible to proceed to the details of the AAVSO(VSX) taxonomy, examining its fine-grained levels of subdivision. Except where noted to the contrary, the AAVSO(VSX) symbols are used also as an authority that AAVSO(VSX) follows closely, although in a spirit of occasional correction, namely the venerable (1948 onward) “General Catalogue of Variable Stars,” or GCVS (administratively associated with the section of Lomonosov Moscow State University entitled the Sternberg Astronomical Institute, formally the Государственный астрономический институт имени Штернберга: particulars are at [www.sai.msu.ru/groups/cluster/gcvs/](http://www.sai.msu.ru/groups/cluster/gcvs/)).

It goes almost without saying that most of the individual detailed levels of subdivision are discussed not only in individual short paragraphs within the VSX introductory page ([www.aavso.org/vsx/index.php?view=about.vartypes](http://www.aavso.org/vsx/index.php?view=about.vartypes)) and (except in the few cases where VSX and GCVS diverge) its GCVS equivalent [www.sai.msu.ru/gcvs/gcvs/vartype.htm](http://www.sai.msu.ru/gcvs/gcvs/vartype.htm), but additionally, and in greater detail, with appropriate bibliographies, in readily locatable individual *Wikipedia* articles. In a few exceptional cases, however, *Wikipedia* details are not readily locatable. In these cases, we add some bibliographic discussion, with the necessary *Wikipedia* pointers. We also allow ourselves remarks on *Wikipedia* materials that, while readily locatable, make especially useful points, over and above what is available from the just-cited AAVSO(VSX) and GCVS introductory pages.

Represented in Sample S are the provisional or placeholder divisions “E” (for the entire eclipsing-variabilities subclass), “E/GS,” “EA,” and “EB” (for portions of the eclipsing-variabilities subclass not yet specified at the duly canonical sub-subclass level), along with the duly canonical sub-sub-classes denoted by the symbols “EA/DM,” “EA/GS,” “EA/SD,” “EB/GS,” and “EB/SD.”

- The **E** binaries are simply the eclipsing binaries (whether totally or merely partially eclipsing). The bare “E” symbol is a placeholder, in the sense of being appropriate for cases in which little is presently known about the system, making it not yet possible to assign a variability sub-subclass.

Within Sample S, this category is represented at AAVSO(VSX) (as consulted in 2022) by the following:  
-\*  $\theta$  Tau (a system that is considered by AAVSO(VSX) to exhibit not only confirmed  $\delta$  Sct-type intrinsic (pulsational) variability, but also possible eclipsing-variability-not-further-specified, yielding the compound VSX symbol, with a colon flagging a mere possibility, “DSCTC+E:”)  
-\*  $\epsilon$  Car (a system considered at AAVSO(VSX) to harbour possible, not confirmed, variability, and accordingly (a) flagged as a mere suspected variable in the VSX interface (with the red “S” status flag, not the green “V”), and (b) in terms of classification symbols given the uncertainty-marked “E:” for “possible eclipsing variability that, if present at all, cannot at the present time be further specified.”)

- The **E/GS** group consists of the eclipsing binaries in which at least one component is in MK phenomenology either a giant or a supergiant. Under this definition, a binary initially and provisionally

labelled merely “E” is to be classified as E/GS even if one component is either a giant or a subgiant, while the other is less luminous, for instance because still residing on the MK-phenomenological MS. While more specific than the bare “E,” this symbol, too, is a placeholder, in the sense of being appropriate for cases in which the available data do not yet make it possible to specify a duly canonical sub-subclass (Sample S features both “EA/GS” and “EB/GS”) of the eclipsing-variability subclass.

Within Sample S, this placeholder category is represented at AAVSO(VSX) (as consulted in 2022) by  $\zeta$  Tau (a system that is considered by AAVSO(VSX) to harbour not only E/GS variability, but additionally  $\gamma$  Cas-type variability, yielding the compound VSX symbol “E/GS+GCAS”).

- In the traditional GCVS notation, followed by AAVSO(VSX) but not universally adopted in the literature, an **EA** is an eclipsing system with a light curve making it possible to specify beginning time and ending time of the eclipses. The traditional definition is thus given in terms of photometric phenomenology, without an attempt to specify an underlying physical mechanism. In particular, (a) in an EA system a Roche lobe might possibly be filled, with a mass-transfer stream therefore possible. However, if there is a filled Roche lobe, with mass transfer, the transfer is on the definition of EA so slight as to make the light curve nearly constant between eclipses. Similarly, (b) “EA” on this definition allows very slight distortions of the mutually gravitating stars from oblate spheroids to ellipsoids or similar shapes, but with the departure from oblate-spheroid symmetry, if present at all, so slight as to keep the light curve nearly constant between eclipses. The bare use of “EA” as a GCVS-and-AAVSO(VSX) classification symbol, without a qualifier in the style “EA/x,” is appropriate in case of an inability to further characterize the evolutionary status (MS? or, rather, evolved beyond MS?) of the binary-system components. While we in this Handbook supplement follow the GCVS-and-AAVSO(VSX) tradition, the authoritative textbook [2007uvs..book....P](#) (in its Section 5.3, on p. 108) favours a not purely photometric-phenomenological definition, in other words favours a more astrophysical definition, on which an “EA” eclipsing binary is required to be detached. It must be admitted that if, as advocated in sub-subsection 6.1.5 above, photometric taxonomy is a search for “real essences,” then the phenomenological use of “EA” is a departure from photometric-taxonomy best practice—justifiable, however, on the basis that “EA” is applied in a mere placeholder spirit, before deeper astrophysical studies become available.

Within Sample S, this category is represented at AAVSO(VSX) (as consulted in 2022) by the following:

- \*  $\delta$  Cas (a system considered at AAVSO(VSX) to be a confirmed variable, but to feature possible-and-yet-not-confirmed EA-type variability: VSX accordingly assigns not the symbol “EA,” but the symbol “EA:”)
- \*  $\eta$  Ori (a system considered at AAVSO(VSX) to harbour not only confirmed EA-type variability (with the placeholder EA type not more closely specified) but also possible-and-yet-not-confirmed  $\beta$  Cep-type pulsational variability: VSX accordingly assigns the compound symbol “EA+BCEP:”)
- \*  $\delta$  Ori
- \*  $\delta$  Vel
- \*  $\lambda$  Sco (a system considered at AAVSO(VSX) to harbour not only EA-type variability (with the placeholder EA type not more closely specified) but also  $\beta$  Cep-type pulsational variability: VSX accordingly assigns the compound symbol “BCEP+EA”)
- \*  $\delta$  Cap (a system considered at AAVSO(VSX) to harbour not only EA-type variability (with the placeholder EA type not more closely specified) but also the  $\gamma$  Dor-type and the  $\delta$  Sct-type pulsational variabilities; VSX accordingly assigns the compound symbol “EA+GDOR+DSCT”; cf further the discussion, with reference to [2010ApJ...713L.192G](#), of GDOR-and-DSCT sub-subclass overlap (comorbidity) in sub-subsection 6.2.4 below)

- An **EB** is in the traditional GCVS notation, followed by AAVSO(VSX) but perhaps (the present writer’s knowledge in composing the current version of this Handbook supplement is not in the relevant way sufficient) not universally adopted in the literature, an eclipsing binary with a light curve (1) making it impossible to specify beginning time and ending time of the eclipses, and (2) satisfying the

additional requirement that the period be longer than in the case of the exceedingly rapid contact binaries, where periods of 1 d or less are typical. (The short periods are partly definitive of the EW UMa stars, in GCVS-and-AAVSO(VSX) notation the “GW” stars, absent from Sample S. The prototype, EW UMa itself, is far fainter than the Sample-S magnitude cutoff, varying in the V band between mag. 9.83 and mag. 11.08.) The present writer believes it is correct to characterize the traditional GCVS-and-AAVSO(VSX) “EB” definition in the same way as the traditional GCVS-and-AAVSO(VSX) “EA” definition is characterized above, namely as a definition in terms not of underlying astrophysics but in terms of photometric phenomenology. The present writer believes, subject to eventual correction by authorities such as AAVSO, that the GCVS and AAVSO(VSX) phrasing, which includes the clause “eclipsing systems having ellipsoidal components,” is offered as an explanation for the definitive phenomenology without being made part of the definition. The “B” in the notation “EB” was evidently intended historically as a mnemonic for  $\beta$  Lyr, notorious for the impossibility of assigning beginning times and ending times to its so-gradual eclipses. However, one should, even if adhering—as we here in this Handbook supplement do—to the GCVS-and-AAVSO(VSX) notational scheme with its phenomenological formal definition for “EB,” keep in view an admonition from the already-cited textbook (on p. 107 of its Section 5.3): “[ $\beta$  Lyr] is so bizarre that it should not be a prototype for any class.” The puzzling  $\beta$  Lyr, in other words, should be regarded as a star that, after the EB class is defined in general phenomenological terms without specific reference to  $\beta$  Lyr, is simply found to lie within the class, and is on close inspection found to present peculiarities not necessarily presented by other stars satisfying the EB definition. In any case, the EB systems, like the EA systems, are so classified merely in a placeholder spirit, given the unavailability of deeper physical studies: the canonical sub-subclasses of the eclipsing-variability subclass of the extrinsic-variability class are, rather, EB/GS and EB/SD, as explained below.

Within Sample S, this placeholder category is represented at AAVSO(VSX) (as consulted in 2022) by the following:

-\*  $\mu$  UMa (a system considered by AAVSO(VSX) to present possible-yet-not-confirmed EB variability, and additionally to present slow irregular cool-star pulsational variability, yielding the compound VSX symbol “EB:+LB”)

-\*  $\beta$  Lyr (a system also considered by AAVSO(VSX), at any rate on the present writer’s interpretation of a somewhat intricate situation, to present possible-yet-not-confirmed intrinsic, eruptive, variability of the DPV type: VSX assigns the symbol “DPV:/EB” [as is discussed in a little more detail in sub-subsection 6.2.5 below, in an examination of the “DPV” sub-subclass of the eruptive-variability sub class of the intrinsic-variability class])

- EA/DM: The **EA/DM** sub-subclass of the eclipsing-variability subclass of the extrinsic-variability class consists of the EA binaries that are fully detached, and in which both components are so unevolved as to be in MK phenomenology MS stars.

Within Sample S, this category is represented at AAVSO(VSX) (as consulted in 2022) by the following:

-\*  $\lambda$  Tau

-\*  $\beta$  Aur

-\*  $\alpha$  CrB

- The **EA/GS** sub-subclass of the eclipsing-variability subclass of the extrinsic-variability class consists of the EA binaries in which at least one component is so evolved as to be in MK phenomenology either a giant or a supergiant. (Under this definition, an EA binary is to be classified as EA/GS even if one component is in MK phenomenology either a giant or a supergiant, while the other is much less evolved, and so much less luminous, for instance through still residing on the MK-phenomenology MS.)

Within Sample S, this category is represented at AAVSO(VSX) (as consulted in 2022) by the following:

-\*  $\gamma$  Per

-\*  $\epsilon$  Aur

-\*  $\eta$  Gem (a system considered by AAVSO(VSX) to harbour both EA/GS and semiregular cool-star variability, yielding the compound VSX symbol “EA/GS+SRA”)

- The **EA/SD** sub-subclass of the binary-variability subclass of the extrinsic-variability class consists of the semi-detached EA binaries, *i.e.* the EA binaries in which one (and only one) star is overflowing its Roche lobe. (It now becomes a little tricky to sort out what were characterized in subsection 6.1.5 above as the “real essences.” So far as can be seen from the present writer’s perspective—this is written very much subject to eventual correction by photometry authorities, such as AAVSO authorities—(a) an EA/SD system would typically, although not as a matter of unavoidable astrophysical necessity, be an EA/GS system (overflow, from a filled Roche lobe, would be a typical scenario only in the case of a bloated, evolved-beyond-MS, star), while (b) an EA/GS system might well fail to be an EA/SD system (for a binary, even with both components supergiants, might well have an orbit wide enough to prevent Roche lobe overflow). “EA/GS” and “EA/SD” are on this provisional reading not intended to be disjoint (as in, for instance, zoology the sets of rodents and primates are intended to be disjoint, or as in geomorphology the classes of mountains and plateaux are intended to be disjoint).

Within Sample S, this category is represented at AAVSO(VSX) (as consulted in 2022) by  $\beta$  Per.

- The **EB/GS** sub-subclass of the eclipsing-variability subclass of the extrinsic-variability class consists of the EB binaries in which at least one component is so evolved as to be in MK phenomenology either a giant or a supergiant. This sub-subclass is thus the EB parallel to the EA/GS sub-subclass, discussed above.

Within Sample S, this category is represented at AAVSO(VSX) (as consulted in 2022) by  $\gamma$  Phe (a system considered by AAVSO(VSX) to also harbour slow irregular pulsation in a cool star, yielding the compound VSX symbol “EB/GS+LB”).

- The **EB/SD** sub-subclass of the binary-variability subclass of the extrinsic-variability class consists of the semi-detached EB binaries, *i.e.* the EB binaries in which one (and only one) star is overflowing its Roche lobe. This sub-subclass is thus the EB parallel to the EA/SD sub-subclass, discussed above (and is subject to the same real-essence questions as were raised above for the EA/SD sub-subclass).

Within Sample S, this category is represented at AAVSO(VSX) (as consulted in 2022) by  $\mu^1$  Sco.

### **6.2.3: The AAVSO(VSX) sub-subclasses of the rotating-variability subclass of the extrinsic variability class:**

Represented in Sample S are the provisional or placeholder division “ROT” (for the entire rotating-variabilities subclass), and additionally the duly canonical sub-subclasses denoted by the classification symbols “ACV,” “BY,” “ELL,” “LERI,” “R,” and “SXARI,” and finally a special situation (perhaps a sub-sub-subclass of the canonical ELL sub-subclass, formally a subset of “ELL”?) denoted by the symbol “HB.” Of these eight symbols, all but three have been taken over from GCVS—the placeholder “ROT,” and the canonical “LERI,” and the specialized “HB.” (AAVSO(VSX) has additionally found it necessary to go beyond GCVS in noting a peculiarity within SXARI, in other words within the SX Ari-type variabilities—namely, variability due not merely to rapid stellar rotation in the presence of a strong stellar magnetic field, but additionally featuring “eclipse-like dimmings probably caused by magnetospherically confined circumstellar disk material that occults the central star.” For this situation, AAVSO(VSX) introduces the symbol “SXARI/E.” Since, however, the refinement is not represented within Sample S, it will not be examined here.)

- The **ROT** cases of variabilities are simply the photospherically inhomogeneous (for instance, spotted) variable-because-rotating stars not otherwise classified. This AAVSO(VSX)-although-not-GCVS symbol would be appropriate for inhomogeneous-photosphere rotating stars awaiting more thorough study, in the same sense in which the GCVS-and-AAVSO(VSX) classification E would be appropriate for eclipsing binaries awaiting more thorough study.

Within Sample S, this category is represented at AAVSO(VSX) (as consulted in 2022) by the following:

- \*  $\alpha$  Cas (a system considered at AAVSO(VSX) to be a confirmed variable, but to feature possible-and-yet-not-confirmed rotational variability: VSX accordingly assigns not the symbol “ROT,” but the symbol “ROT:”)
- \*  $\eta$  Tau (a system considered at AAVSO(VSX) to harbour not only rotational variability, but additionally MS, hot-star, slow-pulsator variability, yielding the compound VSX symbol “ROT+SPB”)
- \*  $\zeta$  Pup (a lone star considered at AAVSO(VSX) to be a confirmed variable, but to feature possible-and-yet-not-confirmed rotational variability: VSX accordingly assigns not the symbol “ROT,” but the symbol “ROT:”)

- The **ACV** sub-subclass of the rotational-variability subclass of the extrinsic-variability class consists of the variabilities in which a star’s strong magnetic fields produce chemical inhomogeneities in the photosphere (with abnormally strong lines of Cr, Si, Sr, and the rare earths), and in which movement of the star in turn causes the brightness to vary as first one set, then another set, of localized anomalous-composition photosphere patches is presented to the observatory. Spectral types are required to fall within the range B8p through A7p (“p” for “chemically peculiar”). In evolutionary terms, the star is required to still lie on the MK-phenomenological MS. The prototype, a member of the unresolved  $\alpha$  CVn A (also known as the “ $\alpha^2$  CVn”) spectral binary, surely has its rotation affected by its close companion (perhaps with tidal locking?). Nevertheless, membership in a binary (tight and tidally locked, or otherwise) is not made part of the definition.

Within Sample S, this category is represented at AAVSO(VSX) (as consulted in 2022) by the following:

- \*  $\alpha$  And (a system considered at AAVSO(VSX) to be a confirmed variable, but to feature possible-and-yet-not-confirmed variability of the type whose paradigm is  $\alpha$  CVn A (“ $\alpha^2$  CVn”): VSX accordingly assigns not the symbol “ACV,” but the symbol “ACV:”)
- \*  $\alpha$  Dor
- \*  $\mu$  Lep
- \*  $\theta$  Aur
- \*  $\gamma$  Cen (a system considered at AAVSO(VSX) to be a confirmed variable, but to feature possible-and-yet-not-confirmed variability of the type whose paradigm is  $\alpha$  CVn A (“ $\alpha^2$  CVn”): VSX accordingly assigns not the symbol “ACV,” but the symbol “ACV:”)
- \*  $\epsilon$  UMa
- \*  $\alpha$  CVn (the system that serves as the ACV paradigm, thanks to the variability of its component  $\alpha$  CVn A, also known as  $\alpha^2$  CVn)
- \*  $\alpha$  Cir (a system that is also considered at AAVSO(VSX) to harbour variability of the character associated with rapidly oscillating chemically peculiar MK-type A stars, yielding the compound VSX symbol “roAp+ACV”)

- The **BY** sub-subclass of the rotational-variability subclass of the extrinsic-variability class consists of variability of the “BY Dra” type. The prototype BY Dra is, admittedly, fainter than the Sample S magnitude cutoff, varying in the V band between magnitude 8.04 and magnitude 8.48. In this grouping, as with the ACV-variability instances discussed above and the SXARI-variability instances discussed at the end of the present sub-subsection, the photometric variation is due to the rotation of an inhomogeneous photosphere, and the star is required not to be evolved beyond the MK-phenomenology MS. In this case, however, the effective temperature is required to be low, corresponding to the two

coolest types in the MK-spectroscopy OBAFGKM sequence. The inhomogeneities typically or always involve localized chromosphere-perturbing activity, in some cases with flaring (causing some specimens of BY variability to be instances of comorbidity, classified not only as BY but additionally as instances of UV Ceti-type (GCVS-and-AAVSO(VSX) symbol “UV”) eruptive variability).

Within Sample S, this category is represented at AAVSO(VSX) (as consulted in 2022) by the following:

-\*  $\epsilon$  Hya (a system considered at AAVSO(VSX) to be a confirmed variable, but to feature possible-and-yet-not-confirmed BY Dra-type variability: VSX accordingly assigns not the symbol “BY,” but the symbol “BY:”)

-\*  $\alpha$  Cen A (considered at AAVSO to be a suspected variable, rather than a confirmed variable, and accordingly assigned not the symbol “BY” but the symbol “BY:”; here, in one of its infrequent departures from classifying entire binaries or entire nested-binaries systems, AASVO(VSX) treats the brighter component of the  $\alpha$  Cen binary as a specimen in its own right [retrievable in a VSX lookup under such things as its BSC designation “HR5459”])

- The **ELL** sub-class of the rotational-variability subclass of the extrinsic-variability class (with “E” for “ellipsoid”) consists in variability due to egg-shape-distorted stars that, through the geometry of their motion, present different photosphere areas to the observatory at different times. Such a situation would be impossible for a solitary star, whether slowly rotating (and so almost a perfect sphere) or rapidly rotating (and so an oblate spheroid with its polar radii less than its equatorial radii). Egg-shape distortion can, however, arise for one or both components of a sufficiently tight binary. AAVSO(VSX), following its senior authority GCVS, additionally requires that ELL variability not arise from an eclipsing binary. Without this additional requirement, EB binaries (subsection 6.2.2 above) in which the light variation is in part due to the varying orientations of ellipsoidal photospheres would become cross-classified as instances of ELL variability. But the present writer wonders whether the GCVS and AAVSO(VSX) decision to avoid cross-classification is appropriate. Might there be some point in distinguishing among (a) partially-or-totally eclipsing binaries in which the photometric fluctuation is due to eclipsing, with egg-shape-distortion, if present at all, making only a negligible contribution to the fluctuation, (b) partially eclipsing binaries in which the photometric fluctuation is due about as much to eclipsing as to the egg-shape distortion, and (c) non-eclipsing binaries in which the photometric fluctuation is solely due to the egg-shape distortion? The first of these three cases might be marked with the appropriate one of the eclipsing-pathology symbols, in the style “EA/x” or “EB/x,” the second with “EA/x+ELL” or “EB/x+ELL” (actually, one would not expect “EA/x+ELL,” but might well encounter “EB/x+ELL”), and the third with “ELL.” The second case, with its “+,” would then become duly highlighted as an instance of comorbidity. On the current practice at GCVS and AAVSO(VSX), provision for the possibility of this comorbidity is (unfortunately, on the present writer’s assessment) lacking.

Within Sample S, this category is represented at AAVSO(VSX) (as consulted in 2022) by the following:

-\*  $\alpha$  Tri

-\*  $\eta$  Aur (a system considered at AAVSO(VSX) to be a confirmed variable, but to feature possible-and-yet-not-confirmed ELL variability: VSX accordingly assigns not the symbol “ELL,” but the symbol “ELL:”)

-\*  $\sigma$  Pup (a system considered at AAVSO(VSX) to harbour not only ellipsoid-type variability, but additionally variability due to slow and irregular stellar pulsation of a cool star: VSX accordingly assigns the compound symbol “ELL+LB”)

-\*  $\alpha$  Vir (a system considered at AAVSO(VSX) to harbour not only ellipsoid-type variability, but additionally pulsational variability of the  $\beta$  Cep type; VSX accordingly assigns the compound symbol “ELL+BCEP”)

-\*  $\pi$  Sco

-\* T CrB (a system considered at AAVSO(VSX) to harbour ellipsoid-type variability, over and above the recurrent-nova feature for which this system is perhaps most celebrated; VSX accordingly assigns the compound symbol “NR+ELL”)

- For AAVSO(VSX), although not, as already noted, for GCVS, **LERI**, or  $\lambda$  Eri-type, variability, is in effect due to a subset of what we in this Handbook supplement call the Be-phenomenon stars (as explained in subsection 5.9 above, the MS or near-MS stars at some point in their known history exhibiting emission lines in spectroscopy); stars in this subset are additionally required by AAVSO(VSX) to exhibit periodic photometric variations, due to one or both of (1) non-radial pulsation and (2) rotation (whether of an inhomogeneous photosphere, or of an inhomogeneous overlying decretion disk, or both). In its introductory, “Conventions Used,” paragraphs at [www.aavso.org/vsx/index.php?view=about.vartypes](http://www.aavso.org/vsx/index.php?view=about.vartypes), VSX takes care to warn users that in the current state of research, the correct choice among “(1) only,” “(2) only,” and “both (1) and (2)” is not known. The present Handbook-supplement writer concurs with the [en.wikipedia.org/wiki/Lambda\\_Eridani\\_variable](http://en.wikipedia.org/wiki/Lambda_Eridani_variable) suggestion that GCVS errs in assigning to the BCEP class the AAVSO(VSX) LERI prototype,  $\lambda$  Eri (with a V-band range of mag. 4.17 to mag. 4.34, almost, but not quite, bright enough for inclusion in Sample S).

Within Sample S, this category is represented at AAVSO(VSX) (as consulted in 2022) by the following:

- \*  $\gamma$  Cas (a system considered at AAVSO(VSX) to harbour not only  $\lambda$  Eri-type variability, but additionally X-ray variability (an AAVSO(VSX) theme left unexamined in this Handbook supplement) and  $\gamma$  Cas-type variability, with the LERI variability considered possible-and-yet-not-confirmed; VSX accordingly assigns the compound symbol “GCAS+X+LERI.”; the authoritative Be-phenomenon database at [obspm.fr](http://obspm.fr) catalogues  $\gamma$  Cas as an instance of the Be phenomenon)
- \*  $\nu$  Pup (a little disconcertingly, the authoritative Be-phenomenon database at [obspm.fr](http://obspm.fr) does not catalogue  $\nu$  Pup as an instance of the Be phenomenon; it is, on the other hand, the case that a shell spectrum, consistent with the Be phenomenon, has been asserted in the literature, with a reference to a “central quasi-emission peak”: the present writer does not know if this differs in any meaningful way from an actual “central emission peak”)
- \*  $\eta$  Cen (a system considered at AAVSO(VSX) to harbour not only  $\lambda$  Eri-type variability, but additionally  $\gamma$  Cas-type variability; VSX accordingly assigns the compound symbol “GCAS+LERI” [and, reassuringly,  $\eta$  Cen is catalogued as an instance of the Be phenomenon by [obspm.fr](http://obspm.fr)])
- \*  $\alpha$  Ara (a system considered at AAVSO(VSX) to harbour not only  $\lambda$  Eri-type variability, but additionally  $\gamma$  Cas-type variability; VSX accordingly assigns the compound symbol “LERI+GCAS” [and, reassuringly,  $\alpha$  Ara is catalogued as an instance of the Be phenomenon by [obspm.fr](http://obspm.fr)])

- The **R** sub-subclass of the rotational-variability subclass of the extrinsic-variability class consists of the instances of variability in which a component of a binary shows “reflection” in its light curve. The term refers here not to reflection as from a mirror, but rather to absorption at one wavelength and re-radiation at some possibly different, longer, wavelength. “Reflection” in this sense may be expected in a sufficiently tight binary, with the wavelength change notable if the two stars are of notably differing photospheric temperatures. (The present writer conjectures that one and the same component in one and the same binary might present both “ELL” and “R” variability, as an instance of comorbidity.) Admittedly, the notion of “extrinsic” variability contains here a potential element of arbitrariness, paralleling the potential element of arbitrariness already noted in sub-subsection 6.2.1 in connection with the classification of shape-distortion, i.e. “ELL,” variation as “extrinsic.” The star that is in the sub-subclass R sense “reflecting” not only is placed into a special temperature regime through irradiation by its companion, if notably different in MK spectral type from its companion, but would actually be in a temporally varying (fluctuating) regime (and so might be accused of actual “externally occasioned intrinsic variation”) if its orbit were to be so eccentric as to make the apastron distance markedly smaller than the periastron distance.

Within Sample S, this category is represented at AAVSO(VSX) (as consulted in 2022) by the following:

- \*  $\nu$  Cen
- \*  $\gamma$  Lup

- The **SXARI** sub-subclass of the rotational-variability subclass of the extrinsic-variability class consists of the instances of SX Ari-type variability. Both GCVS and VSX remark that the SX Ari stars are “high-temperature analogues” of the  $\alpha$  CVn A (“ $\alpha^2$  CVn,” symbol “ACV”) class: strong magnetic fields produce chemical inhomogeneities in the photosphere (with abnormally strong lines of neutral helium and doubly ionized silicon), and rotation of the star in turn causes the brightness to vary as first one set, then another set, of localized anomalous-composition photosphere patches comes into view. Spectral types are required to fall into the range B0p through B9p (“p” for “chemically peculiar”). In evolutionary terms, the star is required to still lie on the phenomenological MS. The prototype, the solitary and rapidly rotating star SX Ari, is rather too faint for inclusion in Sample S (varying between mag. 5.75 and mag. 5.81), but is nevertheless bright enough to possess a Flamsteed number, as 56 Ari. *Wikipedia* consequently discusses this prototype at the mildly unexpected URL [en.wikipedia.org/wiki/56\\_Arietis](https://en.wikipedia.org/wiki/56_Arietis) (while discussing the variability type at the expected URL [en.wikipedia.org/wiki/SX\\_Arietis\\_variable](https://en.wikipedia.org/wiki/SX_Arietis_variable)).

Within Sample S, this category is represented at AAVSO(VSX) (as consulted in 2022) by  $\epsilon$  Cas (a system considered at AAVSO(VSX) to be a confirmed variable, but to feature possible-and-yet-not-confirmed SXARI variability: VSX accordingly assigns not the symbol “SXARI,” but the symbol “SXARI.”; the [obspm.fr](https://obspm.fr) database considers  $\epsilon$  Cas to be an instance of what we in the Handbook call the “Be phenomenon,” and yet, perhaps surprisingly, this system is not classified either as GCAS or as LERI by AAVSO(VSX)).

- The **HB** sub-subclass of the rotational-variability subclass of the extrinsic-variability class is characterized by AAVSO(VSX) (not by GCVS), in the following terms: “Heartbeat stars. A type of eccentric binary stars ( $e > 0.2$ ) whose light curves resemble a cardiogram. They are ellipsoidal variables that undergo extreme dynamic tidal forces. As the two stars pass through periastron, brightness variations occur as a consequence of tidal deformation and mutual irradiation. There may also be tidally induced pulsations present. The morphology of the photometric periastron variation (heartbeat) depends strongly on the eccentricity, inclination and argument of periastron. The amplitude of variations is very small, usually below 0.01 mag. but it may exceed 0.3 mag. in extreme cases.” It is stated at [en.wikipedia.org/wiki/Heartbeat\\_star](https://en.wikipedia.org/wiki/Heartbeat_star) that this variability type was demarcated in 2012, on the basis of OGLE, and more significantly of NASA Kepler-mission, photometry. The present writer thinks, subject to correction, that an HB binary, if non-eclipsing, would at AAVSO(VSX) be considered a special case of ELL variability, in the same set-and-subset sense as an EA system is considered both at AAVSO(VSX) and at GCVS to be a special case of an E system. It appears reasonable for AAVSO(VSX) to place its special HB situation somehow within the “rotating” subclass of extrinsic variabilities, rather than somehow within the “pulsating” subclass of intrinsic variabilities: under the just-quoted AAVSO(VSX) HB characterization, while a star in a highly eccentric HB binary may at periastron be driven into pulsation by the gradient in the gravitational field of its companion, it need not be driven into pulsation. (As stated or implied in the characterization, there may be, and yet not be, “tidally induced pulsations present.”) A background point of special interest is supplied by [en.wikipedia.org/wiki/Heartbeat\\_star](https://en.wikipedia.org/wiki/Heartbeat_star) with its remark that if pulsations are present, then they can take on a one-sided geometry, due to the tidally induced shape distortion of the pulsator.)

Within Sample S, this category is represented at AAVSO(VSX) (as consulted in 2022) by the following:

-\*  $\iota$  Ori

-\*  $\theta$  Car

-\*  $\epsilon$  Lup (a system considered at AAVSO(VSX) to harbour not only HB-type variability, but additionally— if the present writer understands AAVSO correctly—many-interior-nodal-surfaces, few-photospheric-nodal-lines g-mode pulsation on the part of a hot MS star: VSX accordingly assigns the compound symbol “HB+SPB”)

#### **6.2.4: The AAVSO(VSX) sub-subclasses of the pulsating-variability subclass of the intrinsic-variability class:**



(0) The following sequence of general points regarding stellar pulsation serves as a background briefing, before our (unavoidably complex and protracted) examination of the (unavoidably numerous) individual AAVSO(VSX) categories of pulsating variables:

- Stellar pulsation can involve pressure waves (“p-waves”) or gravity waves (“g-waves”; not to be confused with the more exotic, more elusive propagating disturbances, or “gravitational waves,” in local spacetime curvature detected from 2016 onward by LIGO, and from 2017 onward also by the Virgo team). A kind of p-and-g-mongrel stellar wave is additionally said to occur in some stellar cases. Pressure waves—whether in stars, or in planetary atmospheres and oceans, or for that matter in solids, as when sounds conduct through rock—are acoustic. In the known stellar cases, p-waves are from the realm of infrasound rather than of human hearing. In particular, pressure waves in the solar photosphere have been detected, through Doppler-shift measurements in stellar-disk spectroscopy, with periods on the order of 5 minutes. Sounds in the Sun are thus on the order of 16 octaves below the concert-hall Middle C. Gravity waves could also be called “buoyancy waves,” since buoyancy, rather than pressure, is in their case the local restoring force. In terrestrial experience, gravity waves are encountered not only at water-air interfaces, but also (in meteorology) where a higher and a lower layer of the atmosphere differ widely in temperature, and there is something—in notable cases a mountain ridge, disturbing the free movement of air masses—to excite oscillations in the thermal boundary. In the standing waves that form on a violin string or drumhead, there is something rather similar to a gravity wave, but with the restoring force supplied by the tension in the filament or membrane, rather than by buoyancy. (Admittedly, the standing wave for its part creates pressure waves, in other words acoustic waves, in the surrounding air, and it is these that actually get heard in the concert hall.)
- An oscillating mass, whether a violin string or an organ-pipe air column, or a bell, or a drumhead, or a musical acoustics-lab Chladni plate, or a star, typically oscillates simultaneously in many different “modes,” also termed “eigenmodes.” Single-mode oscillation, on the other hand, is encountered in the pure tone generated in the acoustics-lab headphones by an audio-frequency tank circuit (capacitor and inductor in series), yielding what an oscilloscope reveals to be a purely sinusoidal alternating current. To a good approximation, single-mode oscillation is encountered in the concert hall when a tuning fork is struck.
- In a violin string’s “fundamental mode,” the movement of each point on the sounding string, with the exception of the anchored endpoints at the peg-box and bridge, obeys a sinusoidal displacement-versus-time curve. The individual sinusoids, although of different amplitudes (larger toward the middle of the string, smaller toward its anchored ends), are identical in period and identical in phase. This period, adjustable by adjusting the string’s restoring force, in other words by tightening or loosening its peg, is the “natural” or “resonant” fundamental-mode frequency.
- In each of its various other modes of vibration, the violin string has a nonzero number of interior motionless points, or “nodal points”: one such point, dividing the string into equal vibrating halves, in the “first overtone”; two such points, dividing the string into equal vibrating thirds, in the “second overtone”; and so on. Each of these overtone modes has its own “natural” period (in the case of the first, second, ... overtone, half, one-third, ... as great as the natural fundamental-mode period). As with the fundamental mode, the individual displacement-versus-time graph of each moving point is a sinusoid, with all these sinusoids possessing the same period (a period that is to be thought of as “natural,” or as “resonant,” for the given overtone). In contrast with the fundamental mode, however, points separated by an interior motionless point now do not vibrate in phase, but with some fixed phase difference (180° in the case of the first overtone, 120° in the case of the second overtone, 90° in the case of the third overtone, and so on).
- Analogous points hold for a sounding plate or sounding membrane, such as a Chladni plate or a drumskin. Now, however, there are not motionless points, but motionless lines (in the case of a drum stretched over a circular hoop, circles interior to the hoop, and concentric with it, and additionally radial straight lines). These systems of motionless lines may be inspected by sprinkling the Chladni plate or drumhead with a fine powder, shaken aside where a given mode puts the metal or skin surface into motion, and left undisturbed where the mode leaves the surface motionless.

- Analogous points hold also for a three-dimensional vibrator, and in particular for a spherical rotating star devoid of significant global magnetism (in other words having at most localized magnetic poles, as in sunspots). In place of the drumhead's nodal lines, however, there are now nodal surfaces—zero, one, two, ... spherical surfaces in the stellar interior, each concentric with the star's photosphere, and additionally zero, one, two, ... planar surfaces in the stellar interior. The various curves formed where the planar surfaces meet the photosphere are circles of latitude and longitude, with the longitude circles intersecting at the rotational poles. (A curious refinement arises, however, in the case of the rapidly oscillating peculiar MK-type A stars, or AAVSO(VSX) “roAp” stars, as is noted again later in this sub-section: a global dipole magnetic field is present, similar to the Earth's dipole magnetic field; with such stars, as with Earth, the magnetic poles are in general offset from the rotational poles; and although the same system of nodal latitude and nodal longitude circles is present as in the non-magnetic case, this system is now anchored not on the rotational but on the magnetic poles.)
- The simplest of the possible spherical-vibrator cases is the sphere that oscillates purely radially, remaining at all times a sphere while changing in size. Such a purely radial oscillation (in stellar astronomy, notably present in the Cepheid variables; for which, however, non-radial modes can also be found) might be in the fundamental mode (the literature counts interior nodal spheres with “ $n$ ,” writing this as the case “ $n = 0$ ”), but could also be in, *e.g.* the first, second, ... overtone (written as “ $n = 1$ ,” “ $n = 2$ ,” ...). A gaseous first-overtone, *i.e.* “ $n = 1$ ,” spherical oscillator might have a non-travelling p-wave oscillation, with a spherical nodal surface deep in the interior. The surface is a place where the local pressure is constant. When the local pressure at points inside the nodal surface is rising (falling), pressure at points outside the nodal surface is falling (rising). A similar situation of purely radial oscillation is possible also with higher  $n$ -values. In more intricate spherical cases, on the other hand, the sphere is deformed, like a wobbling jelly ball, and one accordingly speaks of “non-radial pulsation.” In such cases, there are zero, one, or more interior nodal surfaces (so that one continues to speak of the fundamental, the first overtone, and the higher overtones, in other words of cases  $n = 0, n = 1, n = 2, \dots$ ) but additionally one, two, or more nodal lines at the surface, analogous to the lines seen on the powdered drumskin. With a rotating star, the analogy with the drumskin is in a notable respect imperfect: where in a drumskin the nodal lines do not migrate as the sound persists, in a star the entire ensemble of longitude circles does migrate, either in the same sense as the sphere's rotation (in “prograde movement”) or in the contrary sense (in “retrograde movement”). Pulsating stars with 0, 1, 2, 3, ... interior nodal planes, *i.e.* exhibiting 0, 1, 2, 3, ... nodal lines at the photosphere, are said to be of “degree”  $l = 0, 1, 2, 3, \dots$ . In addition, the symbol “ $m$ ” is used to document separately the photospheric nodal lines-of-longitude and photospheric lines-of-latitude, in what is called an accounting of “azimuthal order”: for each given  $l, m$  ranges over the set  $\{-l, -l+1, \dots, 0, \dots, l-1, l\}$ , with  $|m|$  the number of photospheric nodal lines that are lines of longitude, in other words the number of lines that correspond to azimuthal-rather-than-declination variation. The integer  $m$  is negative (positive) if the lines of longitude are migrating with respect to the stellar rotation in the retrograde (prograde) sense. So, for example, if a star were to have among its various pulsational modes the mode corresponding to  $n = 4, l = 17$ , and  $m = -6$ , then it would be vibrating in the fourth overtone (in other words with four spherical nodal surfaces in its interior), and with a grand total of 17 photospheric nodal lines. Of these 17, 11 would be lines of latitude, and 6 lines of longitude, and the ensemble of lines of longitude would be migrating opposite to the direction of stellar rotation. This classification scheme now makes it possible to ask, in the spirit of Sherlock Holmes in “Silver Blaze” (subsection 6.1.6 above), “Which of the various arithmetically possible ( $n, l, m$ ) combinations (arithmetically possible in the sense that  $|m| \leq l$ ) is absent from the given star's actually realized pulsation modes—or more generally, is of some unexpected, perhaps dramatically low, amplitude—and why?”
- Violin strings are celebrated for their musical quality, in that of all the arithmetically possible modes, only a small number are excited to more than a negligible amplitude, and the natural periods of these few stand to each other in aesthetically pleasing simple integer ratios. In drumheads, on the other hand (with the exception of the pleasingly musical tympani), many modes are excited, simple integer ratios of periods are no longer prominent to the ear, and the effect is in musical terms a dissonance. A ringing Chladni plate is perhaps less pleasant than a violin string, while more musical in quality than a bass

drum or snare drum. How musical or dissonant, then, are the stars—or, at any rate, how musical or dissonant is the star most extensively studied, the Sun? The disappointing answer is that in musical terms the Sun is merely in the category of drums-other-than-tympani. Nevertheless, at [www.konkoly.hu/staff/kollath/stellarmusic](http://www.konkoly.hu/staff/kollath/stellarmusic) Hungary's Konkoly Observatory documents various explorations of aesthetic possibilities, offering various tonal suggestions for avant-garde composers.

- Oscillations in stars can be excited in at least two different ways: through the “kappa mechanism,” and alternatively (as in the case of the Sun's observed p-waves) through short-lived episodic perturbations driven by the vagaries of convection. The present version of this Handbook supplement examines only the first of the two in detail.
- In the “kappa mechanism,” some significant percentage of the atoms of some relevant atomic species, at some appropriately deep level in the stellar interior, undergo a further degree of ionization as the pressure and density at that level temporarily increase. This process of ionization enhancement temporarily absorbs energy coming up from below, causing that layer to temporarily be a less efficient upward conduit of energy. The temperature below this increasingly ionized, temporarily transport-blocking, layer accordingly rises. When a sufficiently high temperature is reached, the star at last expands, with the blocking-layer opacity now falling in a release of its dammed-up energy, and with its ionization reduced. As the star expands, momentum carries the upward movement temporarily above the normal point of equilibrium between upward pressure and downward weight, in overshoot. As gravity finally wins out over upward pressure, the star starts to contract again, eventually with overshoot below the normal pressure-versus-weight equilibrium point. With the downward overshoot comes an increase of pressure and density in the partial-ionization layer, once again producing enhanced ionization in the relevant atomic species, causing the cycle to repeat. In the case of the Sample S IS stars, the relevant blocking-layer atomic species is singly ionized helium (doubly ionized when pressure and density increase, and the layer opacity rises). In the case of the  $\beta$  Cep-type pulsators, on the other hand, the relevant atomic species is (as is again soon noted in this sub-subsection) some ionization level of iron.
- Stellar pulsation is of astrophysical interest, in helioseismology (and in the twenty-first century, also in the emerging discipline of asteroseismology). A principal achievement of helioseismology has been the identification, in the solar interior, of the solar tachocline, or boundary surface between the Sun's radiative deep interior and its overlying convective zone. Perhaps the most contested question in helioseismology is this: Are g-waves (believed by both sides in the recent debate to be present at deep levels, right down to the centre of Sun) detectable by subtle traces present at the photosphere, and therefore amenable to Doppler-shift spectroscopy? We will not attempt to pronounce on the current state of the debate here. We do, on the other hand, briefly note the practical implications of the debate: a duly developed helioseismology might hope to enhance current capabilities for predicting solar-dynamo phenomena, including sunspots and coronal mass ejections (CMEs); success in predicting CMEs might in turn someday help engineers mitigate the global power-grid destruction entailed by a CME on the scale of the 1859 “Carrington Event.”

(1) We begin this examination of the AAVSO(VSX) pulsational sub-classes with the hot pulsating “BCEP,” “SPB,” and “SPBe” cases.

- The **BCEP** sub-subclass of the pulsating-variability subclass of the intrinsic-variability class consists of the “variables of the  $\beta$  Cep type” (also known in parts of the literature as “ $\beta$  CMa stars”). These are pulsating stars, lying within the hot (blue) MK spectral types O8 through B6, and not so evolved as to be (invariably unstable) supergiants. Although such stars are too hot to lie on the IS, where pulsation is driven by the ionization of singly ionized helium to doubly ionized helium, their pulsation mechanism has since the 1980s been known to be analogous to the IS mechanism, with iron ionization (deep in the stellar interiors) playing the role of IS helium ionization in the less deep layers of somewhat cooler (yellow, MK A,F,G) stars. The 1980s BCEP theoretical advance is discussed further by [2007uvs..book....P](#), in Section 6.5, on p. 142.

Within Sample S, this category is represented at AAVSO(VSX) (as consulted in 2022) by the following:

- \*  $\gamma$  Peg (a system that is considered at AAVSO(VSX) to harbour not only BCEP-type variability, but also another of the pulsational sub-subclasses of variability, yielding the compound symbol “BCEP+SPB”)
- \*  $\epsilon$  Per
- \*  $\eta$  Ori (a system in which the BCEP variability is considered at AAVSO(VSX) to be possible-and-yet-not-confirmed, and in which AAVSO(VSX) also finds confirmed eclipsing variability, not yet fully studied, of the placeholder EA type: this yields the compound symbol “EA+BCEP:”)
- \*  $\zeta$  CMa (a system considered at AAVSO(VSX) to harbour possible, not confirmed, variability, and accordingly (*a*) flagged as a mere suspected variable in the VSX interface (with the red “S,” not the green “V”), and (*b*) in terms of classification symbols given the uncertainty-marked “BCEP:,” for “possible BCEP variability”)
- \*  $\beta$  CMa
- \*  $\chi$  Car
- \*  $\delta$  Cru
- \*  $\alpha$  Mus
- \*  $\beta$  Cru
- \*  $\alpha$  Vir (a system considered at AAVSO(VSX) to harbour also ellipsoid-type variability, yielding the compound VSX symbol “ELL+BCEP”)
- \*  $\epsilon$  Cen
- \*  $\beta$  Cen
- \*  $\iota$  Lup (a case considered at AAVSO(VSX) to harbour confirmed variability, but to be a possible-and-yet-not-confirmed instance of BCEP variability, yielding the symbol “BCEP:”)
- \*  $\alpha$  Lup
- \*  $\kappa$  Cen
- \*  $\delta$  Lup
- \*  $\sigma$  Sco
- \*  $\theta$  Oph (a system considered at AAVSO(VSX) to harbour not only BCEP-type variability, but also another of the pulsational sub-subclasses of variability, yielding the compound symbol “BCEP+SPB”)
- \*  $\lambda$  Sco (a system considered at AAVSO(VSX) to also be an eclipser of the placeholder EA type, yielding the compound VSX symbol “BCEP+EA”)
- \*  $\kappa$  Sco
- \*  $\beta$  Cep (the paradigm for the sub-subclass)

- The **SPB** symbol, used at AAVSO(VSX) although not at GCVS, denotes a sub-subclass of the pulsating-variability subclass of the intrinsic-variability class characterized as “Main Sequence B2–B9 stars (3–9 solar masses) that pulsate in the high radial order low degree g-modes.” We take it here in this Handbook supplement (subject to possible eventual correction by AAVSO, or by others) that here “radial order” refers not to the symbol “*l*” as discussed earlier in this sub-subsection, but simply to “*n*” (since why otherwise could the phenomenon be “radial”?). On this reading of AAVSO(VSX), as the various pulsation modes of an SPB star are inventoried, modes are found in which there are many interior spherical nodal surfaces, and yet no modes are found having a large number of photospheric nodal lines: the pulsation of an SPB star, while non-radial, is just mildly non-radial.

Within Sample S, this category is represented at AAVSO(VSX) (as consulted in 2022) by the following:

- \*  $\gamma$  Peg (a system considered at AAVSO(VSX) to also harbour  $\beta$  Cep-type variability, yielding the compound symbol “BCEP+SPB”)
- \*  $\eta$  Tau (a system considered at AAVSO(VSX) to also harbour some type of rotational variability, yielding the compound VSX symbol “ROT+SPB”; although AAVSO(VSX) asserts neither  $\gamma$  Cas-type nor  $\lambda$  Eri-type variability, the system is classified by the [obsprm.fr](http://obsprm.fr) database as an instance of what is in Subsection 5.9 above called the “Be phenomenon”)
- \*  $\eta$  UMa (a system that might, in view of its rapid rotation and its line variability, be thought to exhibit the “Be phenomenon”; however, this system is absent from the authoritative Be-phenomenon database at [obsprm.fr](http://obsprm.fr))

- \*  $\epsilon$  Lup (a system considered at AAVSO(VSX) to also harbour the “Heartbeat Star” type of rotational extrinsic variability, yielding the compound VSX symbol “HB+SPB”)
- \*  $\theta$  Oph (a system considered at AAVSO(VSX) to also harbour  $\beta$  Cep-type variability, yielding the compound VSX symbol “BCEP+SPB”)
- \*  $\zeta$  Peg (a system that might, in view of its fast rotation, possibly be an instance of the “Be phenomenon,” especially if the MK luminosity class, namely III, assigned in this Handbook supplement (Table, “MK Type” column) should happen to be in error, with the correct class instead being V; however,  $\zeta$  Peg is absent from the authoritative Be-phenomenon database at [obspm.fr](http://obspm.fr))

- The **SPBe** symbol, used in VSX although not at GCVS, is explained by AAVSO(VSX) as signifying “rapidly rotating Be stars showing g-mode non-radial pulsations.” The present writer wonders if this sub-subclass of the pulsating-variability subclass of the intrinsic-variability class should be subdivided, or alternatively if its characterization should be tightened up, to exclude stars that have evolved far beyond the MS: a rapidly rotating Be star not evolved far beyond the MS, and with a history of at least temporary Be emission, is an instance of the Be phenomenon (with its emission due, on the current understanding of that phenomenon, to the formation of an equatorialcretion disk, as discussed in Subsection 5.9 above); and on the other hand, a highly evolved B star in emission (here, admittedly, rapid rotation is unlikely) will be liable to have its emission instead due to copious winds, and so will be liable to sit within an ejected bright gaseous aggregate not confined to the geometry of a disk. It is stressed in Subsection 5.9 above that a highly evolved star of MK type B, with emission, does not qualify as an instance of the Be phenomenon.

Within Sample S, this category is represented at AAVSO(VSX) (as consulted in 2022) by  $\beta$  CMi (classified as an instance of the Be phenomenon by the [obspm.fr](http://obspm.fr) database).

(2) We continue this examination of the AAVSO(VSX) pulsational sub-classes by taking a group of pulsators having no very striking affinity with any other pulsator group represented in Sample S (although arguably having some affinities with the GCVS-and-AAVSO(VSX) S Dor-type variables, or “SDOR stars”: a celebrated SDOR specimen, P Cyg, temporarily surmounted the Sample S brightness threshold around the year 1600, and perhaps again on one or two occasions later in the 1600s; another celebrated SDOR specimen,  $\eta$  Car, temporarily surmounted the Sample S brightness threshold in 1843, having, however, fallen far below the threshold by 1856; as explained in Section 1 above, the Handbook brightest-stars supplement chooses to omit these specimens from Sample S, since there is no concrete, timeframe-assignable, expectation of their at some point once again surmounting the threshold).

- The **ACYG** sub-subclass of the of the pulsating-variability subclass of the intrinsic-variability class consists of the “variables of the  $\alpha$  Cyg type”: rapidly, non-radially, pulsating MK-class supergiants in the hot MK types B and A. The group was first considered a distinctive class of pulsators in 1985, in a predecessor edition of the present-day GCVS. Both the present-day GCVS and AAVSO(VSX) mention Bep and Aep (“p” as the MK-qualifier flag for “chemically peculiar,” “e” as the MK-qualifier flag for “emission”). However, this is surely not definitive, since the unavoidable prototype,  $\alpha$  Cyg, MK-classified as A2 Ia, is in MK terms neither “p” nor “e.” (Here, then, is an instance of the situation discussed in sub-subsection 6.1.5 above, where defining features in a taxonomic group need to be distinguished from empirically noted features, widely present across the group.) Since multiple modes of rapid pulsation are on the GCVS-and-AAVSO(VSX) definition possible, beat effects, capable even of giving the impression of irregular pulsation, may be expected. The ACYG pulsators are notably hotter (bluer) than the IS and near-IS variables (a broad family represented in Sample S by the DCEP, DCEPS, DSCT, DSCTC, GDOR, and roAP pulsators; the IS and near-IS pulsators are in turn notably hotter (yellowish, less red) than the LB pulsators, the LC pulsators, the M pulsators, and the various SR-grouped pulsators).

Within Sample S, this category is represented at AAVSO(VSX) (as consulted in 2022) by the following:

- \*  $\beta$  Ori
- \*  $\epsilon$  Ori
- \*  $\kappa$  Ori
- \*  $\sigma^2$  CMa
- \*  $\eta$  CMa
- \*  $\alpha$  Cyg

(3) As a next step in this examination of the AAVSO(VSX) pulsational sub-subclasses, we take the fairly hot pulsators groups that are “DCEP”(these must be discussed at length, in view of their special role in establishing the intergalactic distance scale),”DCEPS,” “DSCT,” “DSCTC,” “GDOR,” and “roAp” (the “roAp” stars, too, must be discussed at length, in view of their special utility in modelling).

- The **DCEP** sub-subclass of the pulsating-variability subclass of the intrinsic-variability class consists of the “classical Cepheids,” a sub-subclass with prototype  $\delta$  Cep. These are yellow Population I (young, comparatively metals-rich) stars, in MK phenomenology bright giants and supergiants, in other words evolved Population I (evolved metals-rich) stars that have made a blueward excursion, temporarily residing on the IS. Multiple excursions toward the blue and returns toward the red are possible, with the consequence that a Cepheid may be making its first, but may also be (as temporarily yellowish) making even its fifth, crossing of the IS. The most common observed case is the case of a second crossing, where evolution is relatively slow. The pulsational character of the  $\delta$  Cep prototype has been known since 1894 (Belopolsky), and the fact that  $\delta$  Cep is a radial pulsator since 1914 (Shapley). The post-1912 efforts to establish the distance of nearby galaxies through a period-luminosity law (Leavitt, communicated by Pickering) were compromised by a failure to distinguish classical Cepheids from a less luminous Population II class of pulsators, with prototype W Vir. (The members of the latter class are not only intrinsically less luminous than the classical Cepheids, but also all happen to lie far from Earth: W Vir itself has an apparent V-magnitude range of 9.46 through 10.75, making it far fainter than the Sample S cutoff.) The failure caused distances of galaxies, very notably M31, to be underestimated by a factor of two. Baade rectified the situation, thereby helping to establish the correct distance scale for extragalactic astronomy, from his 1944 publishing onward. With the classical Cepheids duly segregated from the misleading W Vir stars, the determination of galactic distances, from observations of extragalactic Cepheids, still requires that some (nearby) classical Cepheids have their distances accurately determined. Until the launch of HST, with its fine-guidance sensor, Cepheids lay beyond the reach of the most reliable (because the most theory-neutral) method of distance determination, the purely astrometric measurement of parallax. From the 1950s until HST, Cepheid distances were determined in several rather indirect ways. In particular, some distances were determined from those Cepheids that could safely be assumed to be (not just coincident on the celestial sphere with, but actually resident in) those Population I assemblages that are the open clusters: open-cluster distances can be obtained from the intrinsic luminosities of cluster members, as deduced through spectroscopic “Main-Sequence fitting.” Again, some distances were determined from those Cepheids that could be established to have not merely optical, but actual binary companions, of spectroscopically ascertainable luminosities and readily known apparent magnitudes. The prospects for more direct classical-Cepheid distance determination have improved in recent years with HST, and additionally with the parallax-dedicated *HIPPARCOS* and *Gaia* missions. We touch on recent distance determinations again in our table, in our “Remarks” for  $\delta$  Cep. Distances aside, a topic of interest in Cepheids, and amenable to CCD investigation under a low financial budget, is the monitoring of period changes. Precise though the classical Cepheids are in their pulsation, tiny changes (perhaps on the order of a mere second in a hundred cycles) do occur. In the case of changes with a consistently increasing or consistently decreasing trend, the process becomes progressively more evident in the “O–C” (“observed minus calculated”) diagram, as the number of cycles plotted becomes progressively greater. Impossible though it may seem for the low-budget observer, working over just a few months or years, actually to observe stellar evolution, it is actually evolution that is postulated as a cause of those special-case Cepheid period changes that are found to have a single progressive trend.

Within Sample S, this category is represented at AAVSO(VSX) (as consulted in 2022) by the following:

- \*  $\beta$  Dor
- \*  $\ell$  (letter ell) Car
- \*  $\eta$  Aql
- \*  $\delta$  Cep

Another Cepheid, at maximum light shining just below the Sample S visibility threshold, is  $\zeta$  Gem (V-band range 3.62–4.18; the formally announced discovery of variability was made rather early, in 1844; further, [www.aavso.org/vsots\\_zetagem](http://www.aavso.org/vsots_zetagem) suggests that since the historical Arabic name “Mekbuda” means “pulled-in paw,” pre-modern Arab astronomers may have already noticed the variability).

- In GCVS-and-AAVSO(VSX) notation, the **DCEPS** symbol (“S” for “symmetrical”?) is used for cases like the DCEP prototype  $\delta$  Cep except in presenting low-amplitude brightness variations and symmetric light curves. For variability to count as DCEP-type, it is required to resemble the prototype  $\delta$  Cep: on this writer’s interpretation of that classification rule, resemblance to the  $\delta$  Cep DCEP prototype makes both a large amplitude and an asymmetric (rapid brightening, slow dimming) light curve mandatory. The present writer believes, subject to possible eventual correction, that the intention of both GCVS and AAVSO(VSX) is not to make the DCEPS a subclass of the DCEP class, but rather to make the two classes disjoint: if the light curve is of low amplitude and symmetrical, then the star is on this interpretation of the rules to be classified as DCEPS-and-not-DCEP. If the interpretation is correct, then the DCEPS variables are a sub-subclass of the pulsating-variability subclass of the intrinsic-variability class, logically coordinate with, and disjoint from, that sub-subclass that is the DCEP variables (in the same sense in which, e.g. in geomorphology the mountains and the plateaux are logically coordinate, disjoint, groupings).

Within Sample S, this category is represented at AAVSO(VSX) (as consulted in 2022) by  $\alpha$  UMi.

- The “DSCT” sub-subclass of the pulsating-variability subclass of the intrinsic-variability class consists of the “ $\delta$  Sct variables,” i.e. stars that reside on the IS, and additionally are in MK phenomenology residing on the MS. As members of the IS, part of their “real essence” is that they share in the helium-ionization pulsation-driving mechanism (that particular form of the “kappa mechanism” that is governed by the transition from singly ionized helium to doubly ionized helium) of the DCEP and DCEPS stars. Pulsation periods are found to be short, from 0.01 to 0.2 days.

Within Sample S, this category is represented at AAVSO(VSX) (as consulted in 2022) by the following:

- \*  $\rho$  Pup
- \*  $\zeta$  Vir
- \*  $\alpha$  Oph (a system considered at AAVSO(VSX) to also harbour  $\gamma$  Dor-type pulsational variability, yielding the composite VSX symbol “DSCT+GDOR”)
- \*  $\xi$  Ser (a system considered at AAVSO(VSX) to harbour possible, not confirmed, variability, and accordingly (a) flagged as a mere suspected variable in the VSX interface (with the red “S,” not the green “V”), and (b) in terms of classification symbols given the uncertainty-marked “DSCT:” for “possible DSCT variability”)
- \*  $\alpha$  Aql
- \*  $\alpha$  Cep
- \*  $\delta$  Cap (a system considered at AAVSO(VSX) to also harbour EA-type (eclipsing) variability and  $\gamma$  Dor-type pulsational variability, yielding the composite VSX symbol “EA+GDOR+DSCT”; for the “GDOR+DSCT” part of this composite, cf. the taxonomic-overlap discussion in [2010ApJ...713L.192G](https://doi.org/10.1007/978-1-4939-7131-2_192), examined in detail later in this sub-subsection, in the treatment of the “GDOR” sub-subclass)

- AAVSO(VSX) departs from GCVS in holding the **DSCTC** class (defined by GCVS as having low

amplitude, in contrast with the  $\delta$  Sct DSCT prototype) to lack physical relevance. The reasoning at AAVSO is that MK-phenomenology MS pulsators on the IS are very typically of low amplitude, and that that the only statistically helpful distinction is between (a) the moderate- and low-amplitude MS IS pulsators and (b) the MS IS pulsators presenting exceptionally high amplitudes, greater than 0.15 in V, labelled in AAVSO(VSX)-and-not-GCVS terms as “HADS.” Sample S lacks HADS stars. The usefulness of segregating even the high-amplitude  $\delta$  Sct pulsators is questioned in [2007uvs..book....P](#) (in Section 6.12, on p. 184), which suggests that the underlying determinant (we would here say, the relevant “real essence” feature) of pulsation amplitude is rotation, with the more rapid rotators pulsating more gently.

Its dissent from GCVS notwithstanding, AAVSO(VSX) does apply the deprecated symbol “DSCTC” to the following cases from Sample S, presumably for historical reasons:

- \*  $\beta$  Cas
- \*  $\theta$  Tau (a system that is considered at AAVSO(VSX) to also harbour possible-and-yet-not-certain eclipsing variability of some as yet undetermined eclipsing type, yielding the composite VSX symbol “DSCTC+E:”)
- \*  $\beta$  Leo
- \*  $\gamma$  Boo (a system considered at AAVSO(VSX) to be a confirmed variable, but only a suspected  $\delta$  Sct variable, classifiable as DSCTC; this yields the VSX symbol, with a colon flagging a mere possibility, “DSCTC:”)
- \*  $\gamma$  UMi
- \*  $\alpha$  Lyr (unfortunately used as a photometric standard from the pioneering 1850s visual work of von Seidel onward, and assessed at AAVSO(VSX) as variable in the V passband over the range  $-0.02$  through  $-0.07$ ; [2007ASPC..364..305G](#) discusses the unfortunate character of this historical error)

- The **GDOR** (“ $\gamma$  Dor”) sub-subclass of the pulsating-variability subclass of the intrinsic-variability class consists (in the characterization at AAVSO(VSX), which is a little tighter than the characterization at GCVS), of “high order g-mode non-radial pulsators, dwarfs (luminosity classes IV and V) from spectral types A7 to F7 showing one or multiple frequencies of variability.” In gross observational terms, periods are longer than for the  $\delta$  Sct variables, extending from 0.25 days to 4 days. Fig. 6.20 (p. 185) of [2007uvs..book....P](#) shows the GDOR group as adjacent to, and slightly cooler than, the IS. The characterizations of AAVSO(VSX), GCVS, and [2007uvs..book....P](#) notwithstanding, however, the key point regarding GDOR variability is that it is due not to the kappa mechanism as in the case of DCEPS variability, but to envelope convection-driven excitations. This leaves open the possibility that one and the same star should, in a display of co-morbidity, be both a  $\delta$  Sct variable (in its various short-period modes of pulsation) and a GDOR variable (in its various longer-period modes of pulsation). [2007uvs..book....P](#) remarks (in sub-subsection 6.12.2, on p. 187) that at least one star realizes this possibility. [2010ApJ...713L.192G](#) takes this theme of category overlap further, drawing on recent NASA *Kepler Mission* photometry, and making a fresh taxonomic proposal: “analysis of /.../data for hundreds of variable stars shows that the frequency spectra are so rich that there are practically no pure  $\delta$  Sct or  $\gamma$  Dor pulsators, in *Wikipedia* essentially all the stars show frequencies in both the  $\delta$  Sct and the  $\gamma$  Dor frequency range. A new observational classification scheme is proposed that takes into account the amplitude as well as the frequency and is applied to categorize 234 stars as  $\delta$  Sct,  $\gamma$  Dor,  $\delta$  Sct/ $\gamma$  Dor or  $\gamma$ Dor/ $\delta$  Sct hybrids.”

Within Sample S, the GDOR category is represented at AAVSO(VSX) (as consulted in 2022) by the following:

- \*  $\alpha$  Oph (a system that is considered at AAVSO(VSX) to harbour also  $\delta$  Sct-type pulsational variability (cf. the remark on  $\delta$  Cap, immediately above), yielding the compound VSX symbol “DSCT+GDOR”)
- \*  $\delta$  Aql (a system considered at AAVSO(VSX) to be a confirmed variable, but to feature possible-and-yet-not-confirmed GDOR variability; VSX accordingly assigns not the symbol “GDOR,” but the symbol “GDOR:”)



-\*  $\delta$  Cap (a system that is also considered at AAVSO(VSX) to harbour EA-type eclipsing variability and  $\delta$  Sct-type pulsational variability (is the latter fact perhaps a consequence of the taxonomic overlap (the comorbidity) discussed in the just-cited [2010ApJ...713L.192G?](#)), yielding the compound VSX symbol “EA+GDOR+DSCT”)

- The **roAp**, or rapidly oscillating stars that are in MK temperature-sequence OBAFGKM terms peculiar A, were noted in the 1970s as a distinctive species of pulsator by Donald Kurtz. The symbol “roAp,” used by AAVSO(VSX) for a sub-subclass of the pulsating-variability subclass of the intrinsic-variability class, is, however, absent from GCVS, which instead has a symbol “ACVO.” This is further explained at [cdsarc.u-strasbg.fr/ftp/cats/B/gcvs/vartype.txt](#): GCVS chooses to emphasize not the pulsational, but instead the rotational, aspect of the grouping, in its choice stressing the similarities between the peculiar-A rapid pulsators and the  $\alpha$  CVn A (“ $\alpha^2$  CVn”) inhomogeneous-photosphere rotators. As pulsators, the roAp stars reside within the  $\delta$  Sct portion of the IS, as an enclave of stars that are not  $\delta$  Sct-like, even while lying entirely in  $\delta$  Sct territory (in the same topological sense as infants born in the sovereign micro-nation of San Marino constitute an enclave, lying entirely within Italian territory, of persons-who-are-not-Italian-citizens). The AAVSO(VSX) definition is as follows: “Rapidly oscillating Ap variables. These are pulsating variables oscillating in high-overtone [i.e. with many interior nodal surfaces-of-constant-radius], low-degree [i.e. with only a low number of photospheric longitude-circle and latitude-circle nodal lines] non-radial pressure modes. Pulsation periods are in the range of 0.003–0.015 days (4–21 min.), while amplitudes of light variation caused by the pulsation are about 0.01 mag. in V. The pulsational variations are superposed on those caused by rotation.” To this definition it might be added, in the spirit of an “underlying real-essence” clarification, that the roAp stars are magnetic, in a sense that differs from the Sun and instead resembles Earth: the magnetism is dominated by a dipole field, whose poles do not in general coincide with the poles of rotation. It is the magnetic poles, not the rotational poles, that define the latitudinal and longitudinal nodal lines of the roAp stars. This unusual geometry presents a unique observational opportunity, in that as the star rotates, different portions of the overall latitude-longitude pulsational grid swing into the view of the observatory. (In the more usual case of a rotating not-globally-magnetic star, not significantly perturbed by any system companion, the poles in the latitude-longitude pulsational grid never change their orientation with respect to the observatory.) The position of the roAp stars within the IS notwithstanding, attempts to explain the pulsations in terms of the singly-ionized-helium kappa mechanism fail. Pulsation is believed to be governed by the dipole magnetic field, which at the magnetic poles inhibits convection. A further observationally convenient circumstance is that spectroscopy reveals the pulsation to involve some atomic species, yet not all. In particular, radial-velocity variations, as the spectroscopic signature of pulsation, are absent in the case of iron. This is explained by the hypothesis that pulsation occurs only at higher levels around the star, involving only those atomic species that are subject to radiational lofting: in a duly quiet stellar atmosphere, iron suffers not lofting, but gravitational settling. Spectroscopy thus conveniently supports inferences regarding the altitude dependence of the photometrically studied pulsation. A table in [en.wikipedia.org/wiki/Rapidly\\_oscillating\\_Ap\\_star](#) lists roAp stars down to V-band mag.  $\sim$ 10.3, serving as a useful reminder that this category contains  $\beta$  CrB A, at V magnitude  $\sim$ 3.6 or  $\sim$ 3.7 almost bright enough to qualify for inclusion in Sample S, and additionally what is perhaps the most chemically bizarre star known, HD 101065 (“Przybylski’s star,” at V-mag  $\sim$ 8.0; even the possibility, from some process of dredge-up, of photospheric plutonium, americium, curium, berkelium, californium, and einsteinium has been asserted).

Within Sample S, this category is represented at AAVSO(VSX) (as consulted in 2022) by  $\alpha$  Cir (a system that is also considered at AAVSO(VSX) to harbour (extrinsic) variability of the rotational type associated with  $\alpha$  CVn A (“ $\alpha^2$  CVn”), yielding the compound VSX symbol “roAp+ACV”).

(4) As the final step in this examination of the AAVSO(VSX) pulsational sub-subclasses, we take the cool (red) pulsators—the Miras (with symbol “M”), then the semiregular “SR” (a placeholder grouping, appropriate where detailed studies are lacking), “SRA,” “SRB,” “SRC,” and “SRD” stars, and finally the irregular “LB” and

“LC” pulsators.

- The **M** sub-subclass of the pulsating-variability subclass of the intrinsic-variability class consists of variables of the Mira (o Cet) type. It is a definitional requirement that the V-band brightness variation be at least 2.5 magnitudes. In actual photometric practice, however, much greater V-band variations can be found, even to the extent of 11 magnitudes. The infrared variation is smaller: as a Mira dims in visible light, it not only cools, but more importantly forms atmospheric titanium oxide, which absorbs visible radiation and re-radiates in infrared. It is additionally a definitional requirement that the MK luminosity class be “giant” (i.e. III), rather than “subgiant,” “bright giant,” or “supergiant” (or indeed, as is sometimes stated, “hypergiant”). In “real essence” terms, this evidently means that Miras are stars of a mass moderate enough to yield an eventual white dwarf, rather than a supernova. Again in “real essence” terms, the Miras are found to lie not on the RGB or in the Red Clump or the HB, but on the AGB, and indeed to be in the final million or so years of their existence as stars. A Mira is, other words, found to be only a million or so years away from evolving into a white-dwarf stellar corpse embedded in a planetary nebula. If a star is a human being, then a Mira is thus not a human in the last year or even the last month of life, but rather a human due to die in a day or two. Interferometry has established that a Mira can be far from spherically symmetric (as one might already conjecture from the observable variety in shapes of planetary nebulae, with the neat smoke-ring symmetry of planetary nebula M57 often absent). In contrast with the “irregular” LB and LC stars, periodicity is by definition discernible, even while not in practice found to be as strict as for the IS pulsators. It is in fact possible that pulsation deep within a Mira is more regular than at the observable level of the convection-perturbed photosphere. Mira pulsation has now been found to involve the fundamental mode, rather than an overtone. But on the other hand, in contrast with IS pulsators, there cannot be an expectation of constant energy generation from the stellar interior, or even of a nearly constant stellar structure from one cycle to the next: since a Mira is on the AGB, it has an inert core, with thermonuclear fusion confined to a hydrogen or helium shell, and these shells can turn on and off rapidly, in “flash” episodes. Emission is present (here the sub-subsection 6.1.5 concern arises once again: in typical observational practice? or, rather, as a matter of formal definition, ultimately stemming from underlying “real essence”?)—making the expected MK temperature types Me, Ce, and Se, rather than M, C, and S. Accompanying the emission is copious mass loss, as a precursor of planetary-nebula formation. Mass loss occurs when a shock wave (visible in spectroscopy as emission) propagating through the atmosphere causes local density to rise and dust grains to consequently condense. These grains, flung out into the ISM, carry some of the stellar gas with them. The topic of Miras is handled in a perhaps unexpectedly thorough way by *Wikipedia*, which offers not only the expected survey article [en.wikipedia.org/wiki/Mira\\_variable](https://en.wikipedia.org/wiki/Mira_variable), but additionally a multiple-article portal, with links to individual articles for over 60 individual Miras, [en.wikipedia.org/wiki/Category:Mira\\_variables](https://en.wikipedia.org/wiki/Category:Mira_variables).

Within Sample S, this category is represented at AAVSO(VSX) (as consulted in 2022 or 2023) by the following:

- \* o Cet
- \*  $\chi$  Cyg

- The **SR** symbol is to be thought of as a placeholder, appropriate where a cool pulsator is not yet well studied, being known simply to fit one of the five canonical pulsational sub-subclasses denoted by “SRA,” “SRB,” “SRC,” “SRD,” and “SRS.” (“SRS,” the group of red-giant semiregular rapid pulsators, however, happens not to be represented in Sample S.) The bare “SR” thus plays the same role in pulsational-variability taxonomy as the bare “EA” (and still more austere, the bare “E”) does in eclipsing-variability taxonomy: the SR grouping is a so-to-speak genus embracing various (in Sample S, four, and yet in full generality five) so-to-speak species. It is not the genus, but the species, that are the canonical sub-subclasses of the pulsational subclass of the intrinsic-variabilities class. The genus consists of all the semiregular significantly evolved moderately cool or very cool pulsators, whether giants or supergiants. Although many of the SR class variables have a rather modest V-band variation,

strong variation is possible also, as instanced by VX Sgr (in the SR “genus,” in the SRC “species,” with exceedingly strong variation in V between mag. 6.52 and mag. 12.4; the mag. 6.52 peak makes VX Sgr a bit too faint for inclusion in Sample S) and strongly-but-not-exceedingly-strongly variable  $\mu$  Cep (again, in the SR “genus,” in the SRC “species,” with variation in V between mag. 3.43 and mag. 5.1: this means that  $\mu$  Cep qualifies, although just barely, for inclusion in Sample S). *Wikipedia* has a convenient tabulation of what is here being called the “genus,” with its five “species,” and with some useful history, at [en.wikipedia.org/wiki/Semiregular\\_variable\\_star](https://en.wikipedia.org/wiki/Semiregular_variable_star). The article notably remarks that SRA, SRB, SRC, and SRD were formalized long ago, in 1958, at the tenth IAU General Assembly, with the fifth so-to-speak species (in AAVSO(VSX) terms, duly canonical pulsational sub-subclass) SRS, by contrast, a recent addition.

Within Sample S, this placeholder category is represented at AAVSO(VSX) (as consulted in 2022) by the following:

- \*  $\mu$  Gem
- \* N Vel
- \*  $\gamma$  Cru
- \*  $\delta$  Vir
- \*  $\alpha$  Ser (a system considered at AAVSO(VSX) to harbour possible, not confirmed, variability, and accordingly (*a*) flagged as a mere suspected variable in the VSX interface (with the red “S,” not the green “V”), and (*b*) in terms of classification symbols given the uncertainty-marked “SR:” for “possible SR variability”)
- \*  $\beta$  Peg

- The **SRA** sub-subclass of the pulsating-variability subclass of the intrinsic-variability class consists of the slow pulsators that are very cool giants (in the sense of falling into MK types M, C, S, Me, Ce, or Se) displaying “persistent periodicity.” Both GCVS and VSX remark that “many of these stars differ from Miras only by showing smaller light amplitudes.” However, to the present writer the remark is puzzling: all the various SR species are by definition semi-regular, and yet Mira for its part is better than merely semi-regular (even while not presenting the clock-like regularity of the IS stars). Perhaps in future editions of this Handbook supplement it will be possible to revisit the problem, resolving the puzzlement. Additionally, it is advisable to keep in view the admonition of [2007uvs..book....P](#) (in section 6.16.1, on page 205) that “the distinction between Miras and semi-regulars (an amplitude of 2.5 magnitudes) is rather arbitrary; there is a smooth gradation of properties between them, and there are stars whose amplitudes [since maxima are not the same from cycle to cycle] may vary from more than 2.5 magnitudes to less.”

Within Sample S, this category is represented at AAVSO(VSX) (as consulted in 2022) by  $\eta$  Gem (a system considered at AAVSO(VSX) to harbour not only SRA-type pulsational variability, but also EA/GS-type eclipsing variability: VSX accordingly assigns a symbol compounded from the pair of symbols “EA/GS” and “SRA,” namely “EA/GS+SRA”).

- The **SRB** sub-subclass of the pulsating-variability subclass of the intrinsic-variability class consists of the slow SR pulsators that are very cool MK-phenomenology giants (in the sense of falling into MK red types M, C, S, Me, Ce, or Se) displaying “poorly defined periodicity,” in other words pulsators like the SRA stars, except still less regular.

Within Sample S, this category is represented at AAVSO(VSX) (as consulted in 2022) by the following:

- \*  $\rho$  Per
- \*  $\gamma$  Hyi
- \* L<sub>2</sub> Pup
- \*  $\sigma$  Lib
- \*  $\alpha$  Her

-\*  $\beta$  Gru

- The **SRC** sub-subclass of the pulsating-variability subclass of the intrinsic-variability class consists of the slow SR pulsators (whether—the present writer presumes—displaying “persistent periodicity” or “poorly defined periodicity”) that are very cool stars (in the sense of falling into red MK types M, C, S, Me, Ce, or Se), and also are in MK-phenomenology terms more tenuous (more sharp-lined, more “tending to hypergiant”) than mere giants.

Within Sample S, this category is represented at AAVSO(VSX) (as consulted in 2022) by the following:

-\*  $\alpha$  Ori  
-\*  $\alpha$  Sco  
-\*  $\mu$  Cep

- The **SRD** sub-subclass of the pulsating-variability subclass of the intrinsic-variability class consists of the slow SR pulsators (whether—the present writer presumes—displaying “persistent periodicity” or “poorly defined periodicity,” and whether in MK-phenomenology terms giants or still-more-sharp-lined-than-giants), that are moderately cool (yellowish) stars, in the sense of falling into the MK yellow-temperatures range FGK.

Within Sample S, this category is represented at AAVSO(VSX) (as consulted in 2022) by the following:

-\*  $\pi$  Pup  
-\*  $\xi$  Pup

In both cases, the system is considered at AAVSO(VSX) to be a confirmed variable, but to feature possible-and-yet-not-confirmed SRD variability. VSX accordingly assigns in both cases not the symbol “SRD,” but the symbol “SRD:”.

- The **LB** group (for AAVSO(VSX), a sub-subclass of the pulsating-variability subclass of the intrinsic-variability class) is explained by GCVS and AAVSO(VSX) in the same words, with the same caveat regarding some special usage at GCVS, absent from AAVSO(VSX): “low irregular variables of late spectral types (K, M, C, S [C and S are chemically anomalous very-cool MK types, parallel to the cool (red) end of the classic-MK OBAFGKM sequence]); as a rule, they are giants. This type is also ascribed, in the GCVS, to slow red irregular variables in the case of unknown spectral types and luminosities.”

Within Sample S, this category is represented at AAVSO(VSX) (as consulted in 2022) by the following:

-\*  $\beta$  And  
-\*  $\gamma$  Phe (a system considered at AAVSO(VSX) to harbour not only LB-type pulsational variability, but also EB/GS-type eclipsing variability: VSX accordingly assigns a symbol compounded from the pair of symbols “EB/GS” and “LB,” namely “EB/GS+LB”)  
-\*  $\gamma$  And (a system considered at AAVSO(VSX) to harbour possible, not confirmed, variability, and accordingly (*a*) flagged as a mere suspected variable in the VSX interface (with the red “S,” not the green “V”), and (*b*) in terms of classification symbols given the uncertainty-marked “LB:” for “possible LB variability”)  
-\*  $\alpha$  Ari  
-\*  $\alpha$  Cet (a lone star (i.e. not in a binary) considered at AAVSO(VSX) to be a confirmed variable, but to exhibit possible-and-yet-not-confirmed LB-type pulsational variability; VSX accordingly assigns not the symbol “LB,” but the symbol “LB:”)  
-\*  $\gamma$  Eri (a system considered at AAVSO(VSX) to be a confirmed variable, but to harbour possible-and-yet-not-confirmed LB-type pulsational variability: VSX accordingly assigns not the symbol “LB,” but the symbol “LB:”)  
-\*  $\alpha$  Tau (a system considered at AAVSO(VSX) to be a confirmed variable, but to harbour possible-and-yet-

not-confirmed LB-type pulsational variability: VSX accordingly assigns not the symbol “LB,” but the symbol “LB:”)

-\*  $\sigma$  Pup (a system considered at AAVSO(VSX) to harbour not only LB-type pulsational variability, but also extrinsic variability in the ellipsoid-star class: VSX accordingly assigns the compound symbol “ELL+LB”)

-\*  $\mu$  UMa (a system considered at AAVSO(VSX) to harbour not only confirmed LB-type pulsational variability, but additionally possible-and-yet-not-confirmed eclipsing variability in the placeholder EB group; VSX accordingly assigns the compound symbol “EB:+LB”)

-\*  $\alpha$  Boo

-\*  $\gamma$  Dra (a system considered at AAVSO(VSX) to be a confirmed variable, but to harbour possible-and-yet-not-confirmed LB-type pulsational variability: VSX accordingly assigns not the symbol “LB,” but the symbol “LB:”)

-\*  $\eta$  Sgr (a system considered at AAVSO(VSX) to be a confirmed variable, but to harbour possible-and-yet-not-confirmed LB-type pulsational variability: VSX accordingly assigns not the symbol “LB,” but the symbol “LB:”)

- The **LC** sub-subclass of the pulsational-variability subclass of the intrinsic-variability class is similar to the “LB” sub-subclass, except that the pulsating stars are stipulated to be in MK-phenomenology terms higher-than-mere-giant. The present writer presumes, the “as a rule” clause in the above-quoted LB characterization notwithstanding, that LB and LC are intended to be disjoint classes. Regarding both LB and LC, it is necessary to keep in view the [2007uvs..book....P](#) admonition (in sub-subsection 6.16.1, on page 205) that it is “rather arbitrary” to make a division between (a) outright irregularity and (b) the semi-regularity that at GCVS-and-AAVSO(VSX) demarcates the SR so-to-speak genus and its five so-to-speak species.

Within Sample S, this category is represented at AAVSO(VSX) (as consulted in 2022) by the following:

-\*  $\epsilon$  Gem

-\*  $\sigma$  CMa

-\*  $\lambda$  Vel

-\*  $\eta$  Car

-\*  $\epsilon$  Peg

-\*  $\zeta$  Cep

(5) We note as a postscript that AAVSO(VSX) (although not GCVS) has a placeholder, **PULS**, applied to stars or systems that have been found in some large-scale survey to harbour pulsation, but whose pulsational character is not at present more precisely known. “PULS” thus serves the same purpose among the pulsators, at AAVSO(VSX), as is served by “E,” both at GCVS and at AAVSO(VSX), among the eclipsing binaries. Within Sample S, this category is represented at AAVSO(VSX) (as consulted in 2022) by  $\epsilon$  Oph.

### **6.2.5: The AAVSO(VSX) sub-subclasses of the eruptive-variability subclass of the intrinsic-variability class:**

- The **BE** symbol is used at GCVS to mark those stars that show what this Handbook supplement (subsection 5.9, above) calls the “Be phenomenon,” and in that additionally there is some photometric variability, but perhaps, in practice, photometric variability is never altogether absent from the overall history of a star when a “Be phenomenon” star is found at the spectrograph to have at some point in its history gone into emission), and in which (a third GCVS definitional requirement) there is no history of an outburst with the photometric extreme of the  $\gamma$  Cas variables. (The “Be phenomenon,” when identified in the spectroscopy of a hot MS star by one or more episodes of emission lines, is commonly found to involve also, in the decades-long photometry record, one or more episodes of photometric outburst. Common though this photometric accompaniment is, it is not invariable. It is where it is

missing—where the photometric variability, so far as historical records go, is always modest, never rising to the level of an outburst—that the “BE” symbol is used at GCVS.) AAVSO(VSX) for its part marks some systems with this same symbol, while noting that the symbol is formally and officially GCVS-not-VSX, and while remarking that “most [GCVS-classified BE stars] may be LERI variables.” It was remarked above that the LERI symbol is used at VSX without being used at GCVS. Perhaps AAVSO(VSX) applies the symbol “BE” for merely historical reasons, provisionally and in advance of eventual deeper study?

Within Sample S, this category is represented at AAVSO(VSX) (as consulted in 2022) by the following:

- \*  $\alpha$  Eri (classified as an instance of the Be phenomenon in the [obsmpm.fr](https://obsmpm.fr) database)
- \*  $\alpha$  Car (a system considered at AAVSO(VSX) to be a confirmed variable, but to harbour possible-and-yet-not-confirmed BE-type eruptive variability; VSX accordingly assigns not the symbol “BE,” but the symbol “BE:”)
- \*  $\omega$  Car (classified as an instance of the Be phenomenon in the [obsmpm.fr](https://obsmpm.fr) database)
- \*  $\gamma$  UMa (a system considered at AAVSO(VSX) to harbour not only BE-type eruptive variability, but also eruptive variability of the UV Cet type; VSX accordingly assigns the compound symbol “BE+UV”; the system is classified as an instance of the Be phenomenon in the [obsmpm.fr](https://obsmpm.fr) database)
- \*  $\zeta$  Oph (classified as an instance of the Be phenomenon in the [obsmpm.fr](https://obsmpm.fr) database)

- The symbol **DPV** is used at AAVSO(VSX), but not at GCVS, for a rather specialized sub-subclass of the eruptive-variability subclass of the intrinsic-variability class, the “Double Periodic Variables,” further divided in the case of some systems outside Sample S into “DPV/ELL” and “DPV/E,” where “E” simply means “eclipsing.” The eruptive episodes are required to have a rather specialized cause, involving binarity: DPV systems are characterized as “semi-detached interacting binaries (with a B-type component) with optically thick disks around the gainer, that experience regular cycles of mass loss into the interstellar medium and are characterized by orbital photometric variability (ellipsoidal, DPV/ELL or eclipsing, DPV/E) in time scales of few days and a long photometric cycle lasting roughly 33 times the orbital period.”

Within Sample S, this category is represented at AAVSO(VSX) (as consulted in 2022) by just one system, the notoriously hard-to-model  $\beta$  Lyr, with its DPV status considered by AAVSO(VSX) as possible-and-yet-not-certain. In a slight variation on its usual syntax (in the usual syntax, the uncertainty-marking colon follows a classification symbol, occurring either as the final character in a (compound or simple) symbol or as the character immediately preceding the compound-symbol marker “+”), AAVSO(VSX) assigns the symbol “DPV:/EB.” We in this Handbook supplement take this somewhat variant symbol to be shorthand for “definitely in the EB eclipsing-variability placeholder grouping within the eclipsing-variabilities subclass of the extrinsic-variabilities class, but also possibly-yet-not-certainly a system in the DPV sub-subclass of the eruptive-variability subclass of the intrinsic-variability class.”

- The **GCAS** group (for AAVSO(VSX), a sub-subclass of the eruptive-variability subclass of the intrinsic-variability class) denotes the “ $\gamma$  Cas variables,” in other words the stars that not only present what is in Subsection 5.9 above called the “Be phenomenon,” but in addition present that prevalent-and-yet-not-universal feature of the Be phenomenon that is the strong photometric outburst. A complication arises for the prototype (as we note again in the “Remarks” column of our table below, in the  $\gamma$  Cas entry): the prevalence of violent outbursts notwithstanding,  $\gamma$  Cas unfortunately cannot itself (in view of its atypical X-ray behaviour) be considered a fully typical Be-phenomenon star.

Within Sample S, this category is represented at AAVSO(VSX) (as consulted in 2022) by the following:

- \*  $\gamma$  Cas (considered at AAVSO(VSX) to harbour not only GCAS-type eruptive variability, but also X-ray variability, and possible-yet-not-confirmed LERI (extrinsic)  $\lambda$  Eri rotational-type variability; VSX accordingly assigns the compound symbol “GCAS+X+LERI:”; classified as an instance of the Be phenomenon in the [obsmpm.fr](https://obsmpm.fr) database)

- \*  $\delta$  Per (considered at AAVSO(VSX) to be a confirmed variable, but to harbour possible-and-yet-not-confirmed GCAS-type variability; VSX accordingly assigns not the symbol “GCAS,” but the symbol “GCAS:”)
- \*  $\zeta$  Tau (considered at AAVSO(VSX) to harbour not only GCAS-type eruptive variability, but also E/GS-type eclipsing variability; VSX accordingly assigns a symbol compounded from the pair of symbols “E/GS” and “GCAS,” namely “E/GS+GCAS”; classified as an instance of the Be phenomenon in the [obsprm.fr](https://obsprm.fr) database)
- \*  $\alpha$  Col (a system considered at AAVSO(VSX) to harbour possible, not confirmed, variability, and accordingly (a) flagged as a mere suspected variable in the VSX interface (with the red “S,” not the green “V”), and (b) in terms of classification symbols given the uncertainty-marked “GCAS:,” for “possible GCAS variability”; classified as an instance of the Be phenomenon in the [obsprm.fr](https://obsprm.fr) database)
- \*  $\kappa$  CMa (classified as an instance of the Be phenomenon in the [obsprm.fr](https://obsprm.fr) database)
- \*  $\rho$  Car (classified as an instance of the Be phenomenon in the [obsprm.fr](https://obsprm.fr) database)
- \*  $\delta$  Cen (classified as an instance of the Be phenomenon in the [obsprm.fr](https://obsprm.fr) database)
- \*  $\mu$  Cen (classified as an instance of the Be phenomenon in the [obsprm.fr](https://obsprm.fr) database)
- \*  $\eta$  Cen (considered at AAVSO(VSX) to harbour not only GCAS-type eruptive variability, but also (extrinsic)  $\lambda$  Eri rotational-type variability; VSX accordingly assigns the compound symbol “GCAS+LERI”; classified as an instance of the Be phenomenon in the [obsprm.fr](https://obsprm.fr) database)
- \*  $\delta$  Sco (classified as an instance of the Be phenomenon in the [obsprm.fr](https://obsprm.fr) database)
- \*  $\alpha$  Ara (considered at AAVSO(VSX) to harbour not only GCAS-type eruptive variability, but also (extrinsic)  $\lambda$  Eri rotational-type variability; VSX accordingly assigns the compound symbol “LERI+GCAS”; classified as an instance of the Be phenomenon in the [obsprm.fr](https://obsprm.fr) database)

- The **UV** sub-subclass (at AAVSO(VSX)) of the eruptive-variability subclass of the intrinsic-variability class consists of the eruptive variables of the UV Cet type, characterized as “K Ve–M Ve stars sometimes displaying flare activity with amplitudes from several tenths of a magnitude up to 6 mag. in V.” Both GCVS and AAVSO(VSX) remark that stars presenting BY Dra variability (symbol “BY,” discussed above), as MS rotators with photospheric inhomogeneities, can also present UV variability (since such rotation can be accompanied by flaring). UV Cet is itself far too faint, even in eruption, to pass the V-band mag.  $\sim 3.55$  threshold for Sample S: AAVSO(VSX), as consulted on 2022 Dec. 27, gives the V-band range 6.8–12.95. The *Wikipedia* writeup dedicated to the prototype UV Cet is, a little unexpectedly, [en.wikipedia.org/wiki/Luyten\\_726-8](https://en.wikipedia.org/wiki/Luyten_726-8). This is a consequence of the fact that UV Cet is the variable-star name given to a star originally catalogued as the “B” component in a tight binary, Luyten 726, and only subsequently discovered to be variable: “Although UV Ceti was not the first flare star discovered, it is the most prominent example of such a star, so similar flare stars are now classified as UV Ceti type variable stars. This star goes through fairly extreme changes of brightness: for instance, in 1952, its brightness increased by 75 times in only 20 seconds.” Also useful for background and context is [en.wikipedia.org/wiki/Flare\\_star](https://en.wikipedia.org/wiki/Flare_star) (as the closest approach in *Wikipedia* to an article explicitly dedicated to UV Cet-type variability).

Within Sample S, this category is represented at AAVSO(VSX) (as consulted in 2022) by  $\gamma$  UMa (considered at AAVSO(VSX) to harbour not only UV-type eruptive variability, but also BE-type eruptive variability; VSX accordingly assigns the compound symbol “BE+UV”; the system is classified as an instance of the Be phenomenon in the [obsprm.fr](https://obsprm.fr) database).

- The **WR** sub-subclass of the eruptive-variability subclass of the intrinsic-variability class involves photometric variability presented by a star that in spectroscopic terms is a Wolf-Rayet, and more particularly is a spectroscopic Wolf-Rayet not yet appearing as the nucleus of a planetary nebula. The photometric variability is in practice not found to be extreme: AAVSO(VSX) comments that what is encountered when photometry is directed to a star known from spectroscopy to be a Wolf-Rayet is “irregular light changes with amplitudes up to 0.1 mag. in V.” AAVSO(VSX) adds that such

fluctuations “are probably caused by /.../ non-stable mass outflow.” While the photometric WR variability is modest, the spectroscopic Wolf-Rayet phenomenon is so dramatic as to be observable by the eye, with a mere spectroscope, in other words with a spectrally dispersive eyepiece-attachment viewing instrument lacking the capability of creating a photographic record. A Wolf-Rayet star is seen at the spectroscope to be in emission, like a Mira (or indeed like a “Be phenomenon” star in an emission episode), and to have emission bands, rather than (as with the Miras and the “Be phenomenon” stars) mere emission peaks. The underlying astrophysical status of Wolf-Rayet stars not yet in planetary nebulae is the following: masses are high, in contrast both with the Miras and with some of the planetary-nebula nuclei; thermonuclear fusion is nearing its end (as with Miras; in the case of actual planetary-nebula nuclei, on the other hand, fusion has already ended); mass loss is still more copious than in the case of Miras; the mass outflow, driven by radiation pressure, causes emission peaks to be Doppler-broadened into the observed bands.

Within Sample S, this category is represented at AAVSO(VSX) (as consulted in 2022) by  $\gamma$  Vel.

#### **6.2.6: The sole Sample S AAVSO(VSX) sub-subclass of the cataclysmic-variability subclass of the intrinsic-variability class:**

The recurrent novae are designated **NR**. While the overall AAVSO(VSX) scheme recognizes various sub-subclasses of the cataclysmic-variability subclass of the intrinsic-variability class, the recurrent novae are the sole such sub-subclass represented in Sample S. It may well be that all observed novae would prove recurrent if observatory records were to extend over some tens of millennia. AAVSO(VSX), however, following GCVS, applies the symbol “NR” only to those novae that the available historical record shows to be recurrent. That record is brief. Spectacular historical novae, like historical supernovae, might in principle some day turn up as archaeo-astronomers continue their inspections of pre-modern chronicles. On the other hand, a duly careful photometric record, at an accuracy of  $\pm 0.1$  mag. in roughly the V band, down to around mag. 9.5, and surveying more than just 200 or 300 stars, perhaps goes no further back than the *Bonner Durchmusterung* (the observations for which began in 1852; [1999JRASC..93...17C](#) supplies some details.)

Within Sample S, this NR category is represented at AAVSO(VSX) (as consulted in 2022) by the T CrB system (actually assigned the compound symbol “NR+ELL,” since one component in this binary presents ellipsoidal-type extrinsic variability).

#### **6.2.7: Other Sample-S-relevant aspects of the AAVSO(VSX) taxonomy:**

(a) AAVSO(VSX) uses **VAR** as a placeholder for variables of unspecified type, explained as “used for suspected variables lacking deeper studies.” Despite the “suspected,” however, all the instances present in Sample S are flagged by VSX as confirmed variables (with the green “V” flag, not with the red suspected-variable “S” flag; in the case of the elaborate  $\beta$  Cap system, with its two most prominent members separated by fully 3.5 arcminutes, lookup in VSX must be made not from “ $\beta$  Cap” but from some such input as “ $\beta$  Cap 1”). The green “V” flag for confirmed variability is assigned by AAVSO(VSX) despite the presence, for all of these systems except possibly  $\zeta$  Cen and  $\beta$  Dra, of an entry in the suspected-variables catalogue NSV (a companion to GCVS, maintained by the same Lomonosov Moscow State University authorities as maintain GCVS).

Within Sample S, the “VAR” category is represented at AAVSO(VSX)(as consulted in 2022) by the following:

- \*  $\beta$  Cet
- \*  $\zeta$  Ori
- \*  $\delta$  CMa
- \*  $\beta$  Cnc
- \*  $\delta$  UMa
- \*  $\zeta$  Cen
- \*  $\beta$  Boo
- \*  $\beta$  Lib



- \*  $\beta$  Dra A
- \*  $\gamma$  Sge
- \*  $\beta$  Cap

(b) AAVSO(VSX) uses its red flag, for suspected variability, in a comparatively small number of cases, in many or all instances from NSV, but a mere subset of the Sample S selection that would be red-flagged as “suspected variables” if NSV were taken as the final arbiter. In a few instances (already discussed in the appropriate places earlier in this photometric “Section 6”), AAVSO(VSX) is able not only to conjecture variability, but to assign a conjectured variability type (using its colon notation, as, e.g. with “GCAS:” for conjectured  $\gamma$  Cas-type eruptive (intrinsic) variability). We order these various cases, as usual, by increasing RA, adding a few comments where necessary:

- \*  $\delta$  And
- \*  $\beta$  Phe
- \*  $\tau$  Cet
- \*  $\beta$  Ari
- \*  $\gamma$  And (with the conjectural variable-type assignment “LB:”)
- \*  $\beta$  Tri
- \*  $\alpha$  Per
- \*  $\zeta$  Per
- \*  $\pi^3$  Ori
- \*  $\iota$  Aur
- \*  $\epsilon$  Lep
- \*  $\beta$  Eri
- \*  $\alpha$  Aur
- \*  $\gamma$  Ori
- \*  $\alpha$  Lep
- \*  $\lambda$  Ori
- \*  $\alpha$  Col (with the conjectural variable-type assignment “GCAS:”; the system is classified as an instance of the “Be phenomenon” in the [obsprm.fr](http://obsprm.fr) database)
- \*  $\zeta$  CMa (with the conjectural variable-type assignment “BCEP:”)
- \*  $\beta$  Gem (with the conjectural variable-type assignment “VAR:”; why is this assignment made, given that  $\beta$  Gem already is red-flagged in VSX as a mere suspected variable?)
- \*  $\epsilon$  Car (with the conjectural (and placeholder) variable type assignment “E:”)
- \* o UMa
- \*  $\iota$  Uma (a suspected variable assigned, at any rate as of 2022 Dec. 29, the symbol “S” by VSX, for “unstudied variable stars with rapid light changes”: did VSX perhaps instead mean to write the colon-qualified symbol “S:”?)
- \*  $\iota$  Car
- \*  $\alpha$  Lyn
- \*  $\alpha$  Hya
- \*  $\theta$  UMa
- \*  $\epsilon$  Leo
- \*  $\eta$  Leo
- \*  $\alpha$  Leo
- \*  $\zeta$  Leo
- \*  $\gamma$  Leo
- \*  $\beta$  UMa
- \*  $\delta$  Leo
- \*  $\theta$  Leo
- \*  $\nu$  UMa
- \*  $\xi$  Hya
- \*  $\epsilon$  Crv
- \*  $\gamma$  Crv

- \*  $\delta$  Crv
- \*  $\beta$  Crv
- \*  $\varepsilon$  Vir
- \*  $\gamma$  Hya
- \*  $\eta$  Boo
- \*  $\beta$  UMi
- \*  $\alpha$  Lib
- \*  $\delta$  Boo
- \*  $\iota$  Dra
- \*  $\eta$  Lup (AAVSO(VSX) assigns to the  $\eta$  Lup system the symbol “CST:,” is perhaps the meaning here that constant light is more likely than the competing possibility of variability?)
- \*  $\beta$  Sco
- \*  $\delta$  Oph
- \*  $\eta$  Dra
- \*  $\beta$  Her
- \*  $\zeta$  Her
- \*  $\eta$  Her
- \*  $\varepsilon$  Sco
- \*  $\eta$  Sco
- \*  $\pi$  Her
- \*  $\xi$  Ser (with the conjectural variable-type assignment “DSCT:”)
- \*  $\theta$  Sco
- \*  $\beta$  Oph
- \*  $\nu$  Oph
- \*  $\delta$  Sgr (with the conjectural variable-type assignment “VAR:”; why is this assignment made, given that  $\delta$  Sgr already is red-flagged as a mere suspected variable?)
- \*  $\eta$  Ser
- \*  $\gamma$  Lyr
- \*  $\zeta$  Aql
- \*  $\pi$  Sgr
- \*  $\beta$  Cyg
- \*  $\gamma$  Aql (with the conjectural variable-type assignment “VAR:”; why is this assignment made, given that  $\gamma$  Aql already is red-flagged as a mere suspected variable?)
- \*  $\gamma$  Cyg
- \*  $\alpha$  Pav
- \*  $\alpha$  Gru
- \*  $\theta$  Peg
- \*  $\eta$  Peg
- \*  $\delta$  Aqr
- \*  $\alpha$  PsA
- \*  $\varepsilon$  Gru
- \*  $\gamma$  Cep

(c) AAVSO(VSX), unlike GCVS, has a special grouping for stars known to be non-variable in the V passband within some reasonable timeframe, and beyond at any rate the tiny level of around 1 millimag. in this same passband. In the VSX interface, these are the stars with the grey “N” flag (as distinct from the green “V” flag, for known variables, and the red “S” flag, for suspected variables). Within this grouping, some, but not all, stars are marked with the symbol “CST” (“constant”), for reasons not clear to this writer. Sample S is represented in the special classification as follows:

- \*  $\beta$  Hyi (with symbol “CST”)
- \*  $\eta$  Cas (with symbol “CST”)
- \*  $\theta$  Eri
- \*  $\delta$  Eri (with symbol “CST”)

- \*  $\alpha$  CMa (with symbol “CST”)
- \*  $\xi$  Gem (with symbol “CST”)
- \*  $\alpha$  CMi (with symbol “CST”)
- \*  $\alpha$  UMa (with symbol “CST”)
- \*  $\gamma$  Vir (with symbol “CST”)
- \*  $\zeta$  UMa
- \*  $\gamma$  TrA
- \*  $\kappa$  Oph (with symbol “CST”)
- \*  $\delta$  Her
- \*  $\gamma^2$  Sgr
- \*  $\delta$  Cyg (with symbol “CST”)
- \*  $\theta$  Aql (with symbol “CST”)

(d) A small minority of the items in Sample S are not mentioned in AAVSO(VSX) at all, whether as known variables, as suspected variables, or as known non-variables. They are listed here (following our usual practice) in order of increasing RA:

- \*  $\alpha$  Phe
- \*  $\eta$  Cet
- \*  $\alpha$  Hyi
- \*  $\gamma$  Cet
- \*  $\alpha$  Ret
- \*  $\epsilon$  Tau
- \*  $\beta$  Tau
- \*  $\beta$  Lep
- \*  $\zeta$  Lep
- \*  $\beta$  Col
- \*  $\alpha$  Car
- \*  $\gamma$  Gem
- \*  $\alpha$  Pic
- \*  $\tau$  Pup
- \*  $\epsilon$  CMa
- \*  $\delta$  Gem
- \*  $\alpha$  Gem
- \*  $\zeta$  Hya
- \*  $\beta$  Car
- \*  $\kappa$  Vel
- \*  $\circ$  Leo
- \*  $\upsilon$  Car
- \*  $\phi$  Vel
- \*  $\lambda$  UMa
- \*  $\mu$  Vel
- \*  $\nu$  Hya
- \*  $\psi$  UMa
- \*  $\lambda$  Cen
- \*  $\alpha$  Cru
- \*  $\beta$  Mus
- \*  $\iota$  Cen
- \*  $\pi$  Hya
- \*  $\theta$  Cen
- \*  $\alpha$  Cen
- \*  $\epsilon$  Boo
- \*  $\beta$  Lup
- \*  $\zeta$  Lup

- \*  $\mu$  Ser
- \*  $\beta$  TrA
- \*  $\tau$  Sco
- \*  $\alpha$  TrA
- \*  $\zeta$  Ara
- \*  $\zeta$  Dra
- \*  $\eta$  Oph
- \*  $\beta$  Ara
- \*  $\gamma$  Ara
- \*  $\upsilon$  Sco
- \*  $\mu$  Her
- \*  $\iota^1$  Sco
- \* G Sco
- \*  $\epsilon$  Sgr
- \*  $\alpha$  Tel
- \*  $\lambda$  Sgr
- \*  $\phi$  Sgr
- \*  $\sigma$  Sgr
- \*  $\xi^2$  Sgr
- \*  $\zeta$  Sgr
- \*  $\lambda$  Aql
- \*  $\tau$  Sgr
- \*  $\delta$  Dra
- \*  $\alpha$  Ind
- \*  $\eta$  Cep
- \*  $\beta$  Pav
- \*  $\epsilon$  Cyg
- \*  $\zeta$  Cyg
- \*  $\beta$  Aqr
- \*  $\gamma$  Gru
- \*  $\alpha$  Aqr
- \*  $\alpha$  Tuc
- \*  $\iota$  Cep
- \*  $\mu$  Peg
- \*  $\alpha$  Peg

### **6.3: Particulars regarding photometry-relevant portions of the table “Remarks” column**

(1) The treatment of individual stars in “Remarks,” in the table below, is best read in parallel with sub-sections 6.2.2 through 6.2.7 above, as follows: note the particular AAVSO(VSX) symbol (for instance, “ACV” or “DCEP” or “E/GS” or “ELL,” or indeed some compound symbol such as “ELL+BCEP” or “EA/GS+GCAS”) assigned in VSX to the case of interest; then examine not only (*i*) whatever “Remarks” may be able to say about the photometric features of the given case (for instance, if coded as a “DCEP” star, its particular pulsational period, and additionally whatever information we may have been able to communicate regarding known long-term period changes), but also (*ii*) sub-sections 6.2.2 through 6.2.7, for the detailed discussion of that particular symbol (where there is also a full inventory of the Sample S stars to which that particular symbol is assigned in VSX; it can be helpful to note what other brightest-stars cases parallel the given case, or alternatively to note that the given case is unique in Sample S).

As an illustration of the recommended procedure, we take  $\eta$  Gem.

This is a hierarchical system, at present only partly resolved, and named “ $\eta$  Gem A” and “ $\eta$  Gem B” in the authority, WDS, which is normative for the leftmost (“Star Name”) column in the brightest-stars table. Since  $\eta$  Gem A is for its part an as-yet-unresolved spectral binary, WDS writes “ $\eta$  Gem A” and “ $\eta$  Gem B” but is not as yet able to write “ $\eta$  Gem Aa,” “ $\eta$  Gem Ab,” and “ $\eta$  Gem B.” Since most of the light from the hierarchical

system is from the unresolved binary  $\eta$  Gem A, this Handbook supplement is constrained, following WDS, to place into the “Star Name” column the entry “ $\eta$  Gem A.” On the other hand, AAVSO(VSX) operates (as remarked in sub-subsection 6.1.2 above) in most cases with the photometry of entire systems, rather than with the photometry of system components. It is consequently necessary to use “ $\eta$  Gem” (not “ $\eta$  Gem A”) as the AAVSO(VSX) lookup term, if one desires to perform a lookup rather than to rely on our own (possibly out-of-date) table-entry report of the recent AAVSO(VSX) assessment.

One performs the (perhaps-desired) lookup in AAVSO(VSX) by first invoking menu item “Search” in the interface at [www.aavso.org/vsxl/](http://www.aavso.org/vsxl/), then entering the seven-character string [eta Gem](http://www.aavso.org/vsxl/). VSX returns much information, including not only the number of photometric observations currently available in the AAVSO database (these can be studied, if desired, from the interface at [www.aavso.org/vsots\\_zetagem](http://www.aavso.org/vsots_zetagem), as distinct from the interface at [www.aavso.org/vsxl/](http://www.aavso.org/vsxl/), starting from the “Pick a Star” field), but also some AAVSO(VSX) variability classification symbol. In the particular case of  $\eta$  Gem, the symbol is compound, “EA/GS+SRA” (as is rather unsurprising, for a three-star hierarchical system). The “EA/GS” portion indicates variability from an eclipsing pair with an Algol-like light curve, in which bottoms and tops are nearly flat, and in which the dimmings can be assigned precise starting and ending times. The “SRA” portion, on the other hand, indicates semi-regular pulsation from a star from the giant-yet-but-not-supergiant part of the MK-phenomenology luminosity-classes “V, IV, III, II, I” sequence, and at the cool end of the MK-phenomenology “OBAFGKM” temperature-types sequence.

Examining (i) what it has at this stage in the development of the bright-stars supplement been found possible to write, regarding photometry, in the  $\eta$  Gem “Remarks,” one additionally (ii) obtains context by looking up “EA/GS” in sub-subsection 6.2.2 and “SRA” in sub-subsection 6.2.4 (examining not only the general descriptions of the EA/GS and SRA sub-subclasses, but also the sub-subsection 6.2.2 list of Sample S stars that resemble  $\eta$  Gem in being assigned the AAVSO(VSX) symbol “EA/GS,” and the 6.2.4 list of Sample S stars that resemble  $\eta$  Gem in being assigned the AAVSO(VSX) symbol “SRA”: Sample S is found to contain several “EA/GS” cases, but no “SRA” case other than  $\eta$  Gem itself).

(2) We have attempted with our own special coding (as we soon explain) to give some indication, in “Remarks,” of Sample S objects for which further research—in principle of any kind, but in current practice in this Handbook supplement particularly in photometry—may be appropriate, even under a low budget of money and person-hours. With the coding we have in view less (a) the amateur equipped with binoculars or a Dobsonian, able to invest no more than ~2000 USD (or CAD, or EUR), or at the other extreme (b) the internationally prominent professional observatory, with a capital investment far into the millions, than (c) the amateur or institution (for instance, the small college) able to make an initial five-figure outlay. This capital investment is to be envisaged as procuring telescope, mounting, wind shelter (rolling-roof shed? small fibreglass dome?) and off-the-shelf mass-market astronomical CCD. In the target constituency “(c),” we envisage an individual or club or small institution—for instance, in the USA a four-year privately endowed college, with small endowment—additionally able to invest a few hundred person-hours for mastering some kind of relevant image-processing software suite, to turn FITS-format files from the astronomical CCD into publishable data. One such suite is the IRAF freeware (for decades dominant in North America, even within the more formidable constituency “(b)”).

To signal what should be signalled, in an unobtrusive way, with a particular view to the needs of the specially targeted “(c)” constituency, we accordingly incorporate in “Remarks” the text-searchable word “advisable” (marking that for such-and-such a sparsely monitored star, such-and-such further observations, of such-and-such a character might be “advisable”). In the “Remarks,” we try not to use this operationally relevant word “advisable” for any other purpose.

Our signalling of research-advisability, both at present and in foreseeable future versions of “Remarks” in future years, downplays those traditionally higher-budget, constituency “(b),” branches of observation that are precision astrometry, interferometry, and precision spectroscopy, concentrating instead (we reiterate) on that potential constituency “(c)” strength that is CCD photometry.

#### **6.4: Further photometry reading**

In the present stage of developing this Handbook supplement, we do not feel able to prepare even a list of the half-dozen most important amateur-appropriate photometry resources published since the year 2000. We do, however, make several miscellaneous, in part randomly ordered, bibliographic remarks, to supplement the

occasional detailed bibliographic remarks appearing in earlier “Section 6” subsections.

- As the second edition of Hearnshaw’s *Analysis of Starlight* ([2014anst.book.....H](#), mentioned in Subsection 5.1 above) is an authoritative history of spectroscopy, so in a parallel way is Hearnshaw’s *Measurement of Starlight* ([1996mest.book.....H](#), [2005mest.book.....H](#)) an authoritative history of photometry.
- A small, yet notable, supplement to Hearnshaw is [1949MNSSA...8...95C](#)—a public-outreach lecture by one of the leading 20th-century photometrists, indicating the state of observatory technique as the photographic plate was just beginning to give way to that single-pixel precursor of the CCD that was the photomultiplier tube.
- The already much-cited Percy book [2007uvs..book.....P](#), will be found suitable for readers at all levels, from the binoculars-equipped observer up to and beyond the professional enrolled in a Ph.D. program.
- The David Levy amateur handbook, originally published in 1989 under the title *Observing Variable Stars: A Guide for the Beginner*, went into a second edition perhaps variously dated to 2003 or 2005, under the new title *David Levy’s Guide to Variable Stars* ([2005dlgv.book.....L](#)). David Levy’s most significant upgrade is the addition of material on CCD methods.
- Helpful reading on the IS includes not only [en.wikipedia.org/wiki/Instability\\_strip](http://en.wikipedia.org/wiki/Instability_strip) and (from a slightly more general perspective) [en.wikipedia.org/wiki/Kappa%E2%80%93mechanism](http://en.wikipedia.org/wiki/Kappa%E2%80%93mechanism), but also (with terse, user-friendly insertions of relevant physics; this is part of an online course offering) [astronomy.swin.edu.au/sao/downloads/HET611-M17A01.pdf](http://astronomy.swin.edu.au/sao/downloads/HET611-M17A01.pdf).
- AAVSO advertises online photometry courses (of which, however, the present writer does not have personal experience) at [www.aavso.org/courses](http://www.aavso.org/courses).
- The troublesome topic of overtones, “orders,” and “degrees,” in the “ $n, l, m$ ” numerical classification of pulsating-star modes (sub-subsection 6.2.4 above) is discussed in user-friendly terms at [2006ASPC..349..10IK](#). Particularly useful is the set of examples in Figure 4, illustrating various possibilities for the subcase-rich case  $l = 3$ . As a preliminary to overtones, orders, and degrees in the 3-dimensional case of a spherical oscillator, it is useful to examine discussions of 2-dimensional cases, as with drumskins, Chladni plates, and bells, including [www.youtube.com/watch?v=IRFysSAxWxl](http://www.youtube.com/watch?v=IRFysSAxWxl) and [blogs.sw.siemens.com/simcenter/sound-and-vibration-of-carillon-bells](http://blogs.sw.siemens.com/simcenter/sound-and-vibration-of-carillon-bells).

#### **APPENDIX: Glossary of acronyms and similar designation**

The following is a glossary of the acronyms and similar designations used in the essay and table. We omit, as sufficiently obvious, a small handful of universally known acronyms (e.g. NASA), designations of chemical elements and chemical compounds (e.g. CO, for carbon monoxide), and the like. We do include a few designations of particular satellites or similar space missions (e.g. *BRITE*, *MOST*).

- AAT: Anglo-Australian Telescope (3.9 m, Siding Spring Mountain, New South Wales, Australia)
- AAVSO: American Association of Variable Star Observers
- AAVSO(VSX): AAVSO International Variable Star Index ([www.aavso.org/vsx](http://www.aavso.org/vsx))
- ALMA: internationally funded Chile-based radio interferometer (“Atacama Large Millimetre/submillimetre Array”)
- AMBER: spectro-interferometer at VLT (“Astronomical Multi-BEam combineR”)
- AGB: asymptotic giant branch (as a region in the two-dimensional MK luminosity-versus-temperature stellar classification space)
- Astron. Alm.: *The Astronomical Almanac*, as the joint annual publication, in print and to a reduced extent online, of the United States Naval Observatory and HM Nautical Almanac Office; “Section H” (not necessarily always up to date in the online version) provides V magnitudes, B–V and V–I colours, and MK types for several hundred bright stars; Astron. Alm. particulars can be had from [asa.hnnao.com](http://asa.hnnao.com) and [aa.usno.navy.mil/publications/asa](http://aa.usno.navy.mil/publications/asa)

- au: astronomical unit (the formal 2012 IAU definition is in effect a precisification, in the (SI) laboratory unit of metres, of the earlier epoch-of-Kepler au concept; before 2012 the concept was defined in astronomical, as distinct from laboratory, terms—before 1976 as the half the sum of the Earth-to-Sun distance at perihelion and the Earth-to-Sun distance at aphelion, and from 1976 onward with a gravitation-theory precisification of that half-of-sum concept)
- *BRITE*: BRITe Target Explorer, a.k.a. Canadian Advanced Nanospace eXperiment 3 (CanX-3: constellation of precision-astrometry satellites (6 attempted, 5 successfully deployed), as a Canada-Austria-Poland collaboration; first launch was in 2013)
- BSC5: Yale Bright Star Catalog, Version 5
- BSG: blue supergiant
- BTA-6: Bolshoi Teleskop Alt-azimutalnyi-6 (Большой Телескоп Альт-азимутальный-6, “Large Alt-Azimuth Telescope 6”): 6-m telescope on north side of Caucasus Mountains, Russia)
- CADARS: Catalogue of Absolute Diameters and Apparent Radii of Stars ([doi.org/10.1051/0004-6361:20000451](https://doi.org/10.1051/0004-6361:20000451))
- CHARA: the Mount Wilson optical interferometer (Center for High Angular Resolution Astronomy)
- CME: coronal mass ejection
- CNO cycle: the carbon-nitrogen-oxygen-catalyzed cycle under which the hotter stars fuse hydrogen into helium
- COAST: the Cambridge optical interferometer (Cambridge Optical Aperture Synthesis Telescope)
- CODEX: a series of computer codes for the numerical simulation of stellar atmospheres (Cool Opacity-sampling Dynamic EXtended)
- *CORIOLIS*: USA-but-not-NASA satellite launched 2003; mission involves not only instrumentation for Earth ocean-environs monitoring, but also solar-wind monitor SMEI (Solar Mass Ejection Imager)
- DAO: Dominion Astrophysical Observatory (on Vancouver Island, British Columbia)
- DR2: Data Release 2 (at *Gaia*)
- ESA: European Space Agency
- ESO: European Southern Observatory (multiple sites, in northern Chile)
- FDU: First Dredge-Up (as a stage in stellar evolution, soon after a star evolves out of the MS)
- FUV: far ultraviolet
- GCPD: General Catalogue of Photometric Data (University of Lausanne, Switzerland)
- GCVS: General Catalogue of Variable Stars (Sternberg Astronomical Institute, Moscow)
- GTR: general theory of relativity
- Hp: a visible-light passband used for photometry at *HIPPARCOS*
- HM Nautical: “His Majesty’s Nautical” (for UK publications and UK agencies)
- HR diagram, HR plot: two-dimensional luminosity-versus-temperature plot for the members of some given population of stars; it is useful to distinguish the “observational” (phenomenological, MK-classification) and the “theoretical” HR diagrams
- HST: *Hubble Space Telescope*
- IAU: International Astronomical Union (Paris)
- IR: infrared
- IRAF: Image Reduction and Analysis Facility: a suite of software tools, for astronomical tasks including aperture photometry and the “extraction of one-dimensional spectra” from raw spectrograms, available free of charge from the National Optical Astronomy Observatory (USA); very

widely used at North American professional observatories, and quite widely also, but in competition with MIDAS, at professional observatories outside North America: [ast.noao.edu/data/software](http://ast.noao.edu/data/software)

- IS: Instability Strip (as a region in the two-dimensional luminosity-versus-temperature stellar-classification space)
- ISM: interstellar medium
- *IUE*: International Ultraviolet Explorer (space telescope: NASA, ESA, and United Kingdom; 1978–1996)
- LESIA: Laboratoire d'Études Spatiales et d'Instrumentation en Astrophysique (physically at Paris-Meudon): [lesia.obspm.fr](http://lesia.obspm.fr)
- LPV: long-period variable
- LSR: Local Standard of Rest (as reference frame for kinematics of bodies in our own galaxy)
- $M_{\odot}$ : solar mass
- mas: milliarcsecond
- MK: Morgan-Keenan (two-dimensional phenomenological, non-theoretical, stellar classification scheme, with “MK luminosity classes” and “MK temperature types”)
- MOST (*Microvariability and Oscillations of Stars/Microvariabilité et Oscillations STellaire*): Canadian space telescope for precision photometry; launched in 2003, deactivated in 2019
- MS: Main Sequence (as a region in the two-dimensional luminosity-versus-temperature stellar classification space; it is useful to distinguish the “observational MS,” in other words the empirical MK luminosity class V, from the “theoretical MS”)
- My: megayears
- NCP: North Celestial Pole
- NPOI: a US Naval Observatory facility (Navy Precision Optical Interferometer)
- NSV: New Catalogue of Suspected Variable Stars (Sternberg Astronomical Institute, Moscow)
- OBAFGKMLTY: the temperature-ordered sequence of MK types, with O the hottest and Y the coolest; until the discovery of brown dwarfs, in types L, T, and (very recently) Y, the sequence was simply OBAFGKM, recalled by 20th-century students with the unfortunate mnemonic “Oh Be A Fine Girl Kiss Me” (implementing gender-neutrality, and allowing for the three progressively cooler brown-dwarf types, one might instead propose “Oh Be A Fine Gymnast, Kiss Me Like This, Yowee”); outside this sequence are the special MK labels (marking gross chemical anomalies) W (for the Wolf-Rayet stars; these turn out to be hot, like O stars), C (for stars whose photospheres are rich in carbon; these turn out to be cool, like K or M) and S (for stars with chemically anomalous photospheres, these are in terms of spectral phenomenology intermediate between M and C, and turn out to be cool); C is the current label for a group that was in earlier decades divided into R and N: additionally, the special “D” and “P” flags are used, in a more colloquial MK spirit, for planetary nebulae hosts and white dwarf “stars”
- OGLE: Optical Gravitational Lensing Experiment: a large long-term (1992-present) variability survey conducted largely from Las Campanas in Chile, under the leadership of Warsaw University, noted both for stellar-variability results and for exoplanet results
- OHP: Observatoire de Haute-Provence (France)
- PA: position angle
- PTI: Palomar Testbed Interferometer
- $R_{\odot}$ : solar radius
- $R^*$ : stellar radius (with reference to some given, reasonably spherical, star)
- $R_{\text{eq}}$ : equatorial radius (with reference to some given rotationally flattened star)



- RGB: red-giant branch (as a region in the two-dimensional luminosity-versus-temperature stellar classification space)
- $R_{\text{pol}}$ : polar radius (with reference to some given rotationally flattened star)
- RSG: red supergiant
- SAAO: South African Astronomical Observatory
- SB: spectral binary, whether double-lined or single-lined
- SB2: double-lined spectral binary
- SETI: search for extraterrestrial intelligence
- SI: *Système International d'Unités*; the internationally agreed system of second-metre-kilogram-ampere-kelvin-mole-candela laboratory units, at one time implemented with recourse to some physical artefacts (including most notoriously the “standard kilogram,” in a vault at the Bureau International des Poids et Mesures in France), but since a 2019 decision defined in a way that can be reproduced by any duly equipped laboratory, independently of artefacts; SI, in its various iterations through the decades, is a 1960 precisification of the earlier internationally agreed “MKS system,” from 1889
- SGB: sub-giant branch (as a region in the two-dimensional luminosity-versus-temperature stellar classification space)
- SMEI: Solar Mass Ejection Imager, as an instrument on the *CORIOLIS* satellite
- SN: supernova
- SNR: supernova remnant
- STIS: Space Telescope Imaging Spectrograph System (an instrument on *HST*)
- SWB: stellar-wind bubble
- TESS: *Transiting Exoplanet Survey Satellite* (NASA)
- UV: ultraviolet
- V: the visible-light passband in the UBVRI photometric passband system that best approximates the response of the human eye, as lying between the blue (“B”) and red (“R”) visible-light passbands
- VLT: a Chile-based facility of the European Southern Observatory (Very Large Telescope)
- VLTI: the interferometer at VLT
- VSX: AAVSO International Variable Star Index ([www.aavso.org/vsx](http://www.aavso.org/vsx))
- WFC3: Wide Field Camera 3 (as an instrument on *HST*)
- WFPC2: Wide Field and Planetary Camera 2 (as an instrument on *HST*)
- WD: white dwarf
- WIRE: *Wide-field Infrared Explorer* (a.k.a. *Explorer 75*, a.k.a. SMEX-5); a NASA space telescope, 1999–2000
- WR: Wolf-Rayet (as a type of star)
- WDS: Washington Double Star Catalog: [www.usno.navy.mil/USNO/astrometry/optical-IR-prod/wds/WDS](http://www.usno.navy.mil/USNO/astrometry/optical-IR-prod/wds/WDS)
- ZAMS: zero-age Main Sequence (the subregion of the MS comprising stars that have just begun stable core-hydrogen fusion)

### **History of recent revisions to both essay and table**

Recent revisions are tracked with UTC YYYYMMDDThhmmssZ timestamping, in the “major.minor.patch” version-numbering scheme common in software development.

- [20230214T144000Z/9.1.1](#): all hyperlinks blue-underlined and active; created bookmarks for sections and subsections; removed the word “Subsection” in headings.
- [20230208T183000Z/9.1.0](#): Changed AU to the abbreviation “au”; changed en dash to em dash in running text; changed all single and double quotes to “curly quotes”; changed all hyperlinks to blue-underlined. [Even though so marked, not all the identified hyperlinks actually link to anything. That will be a chore for later on.]
- [20230202T210000Z/9.0.0](#): Added to version 8.0.0 two hitherto overlooked, but occasionally bright, stars,  $\chi$  Cyg A and  $\mu$  Cep, and in part as a consequence of this addition checked and adjusted the numbers of stars asserted to be present in Sample S and some salient supersets of Sample S. Added long photometry section (“Section 6”) to essay. Added to the Remarks cell for each table entry a summary of AAVSO(VSX) situation (whether the given lone star or binary system or nested-binaries system is flagged at AAVSO(VSX) as a confirmed variable, as a suspected variable, or as a confirmed non-variable; and if one of these three flags is present, then also what variability classification symbol, if any, is assigned at AAVSO(VSX)). This version 9.0.0 was sufficiently polished to support uploading to the online-version server (but was prepared in the realization that the RASC editorial team might also choose to make small adjustments, to be documented here, before their upload, in which case what would be uploaded by that team would instead be version 9.x.y, with  $x$  most likely = 0 and  $y$  most likely = some small positive integer).
- [20220816T235901Z/8.0.0](#): Precessed RA, DEC to epoch 2023.5. Overhauled photometry, (*a*) taking  $m_v$  and  $B-V$  values from GCPD where possible, and otherwise from the various post-1990 sources used by SIMBAD, (*b*) indicating in  $m_v$  and Remarks columns all cases of confirmed variability, suspected variability, and confirmed non-variability documented at AAVSO(VSX). This version was sufficiently polished to support the print edition of the 2023 Handbook, but not yet sufficiently polished to support uploading to the online-version server.
- [20220311T033032Z/7.0.1](#): Significant editing and formatting, particularly tabs and fitting long remarks into available space; corrected a few spelling errors. Removed all instances of “http://” as these are redundant in a URL; also removed italics on HST (only use italics on full text, i.e. *Hubble Space Telescope*)
- [20220303T210237Z/7.0.0](#): Performed sufficient updating of the 5.x.x version series to support not only the print edition of the 2022 Handbook, but also to support uploading to the online-version server. With the now-noted binarity of  $\zeta$  CMa Aa,Ab, the count of Sample S (in the essay) was increased from 322 stars to 323 stars. Also in the essay, the former Section 4 was reassigned as Section 5, and a rather long Section 4 was inserted, discussing the astrophysics of binaries. Concomitantly with this addition of this essay section, astrometric detail was added, under “Remarks,” to perhaps roughly one out of every four table entries, in a general review of the treatment of binaries, and underlining was added in the leftmost column of the table to flag cases in which a binary system possesses a published orbital solution. Apart from many small revisions in the table, extended “Remarks” treatments were inserted for two stars of special interest,  $\delta$  Sco A (amateur photometry and amateur spectroscopy is particularly needed during and around the time of periastron passage, in 2022 May) and  $\alpha$  Her Aa.
- [20210811T201642Z/6.0.0](#): Performed sufficient updating of the 5.x.x version series to support the print edition of the 2022 Handbook, but without sufficient updating to support uploading to the online-version server.
- [20210807T203107Z/5.2.0](#): Made various copy-edit corrections (such as insertion of missing punctuation, correction of a few spelling errors), and additionally on the side of scientific substance made a few corrections or amplifications (chiefly as follows: amplified the essay elucidation of “n,” “nn” in MK types; improved an essay remark on rotation in stellar evolution; made essay correction regarding protracted-versus-brief membership of SGB; corrected essay list of Be-phenomenon stars (the phenomenon is not

observed in Adhara); added “SGB” to glossary of acronyms; improved table discussion of exoplanet status for  $\alpha$  Tau A (Aldebaran); corrected table magnitude range for  $\alpha$  Ori Aa (Betelgeuse); corrected table typo for angular distance in  $\alpha$  Cru AB (Acrux and companion; correct value is 3.5”, not 35”); corrected table typo for magnitude of  $\eta$  Oph B (correct value is 3.3, not 7.3); updated  $\alpha$  PsA Aa (Fomalhaut) table entry to reflect the fact that *HST*-imaged “exoplanet” Dagon (2008) has now faded below the imaging threshold, and is therefore now believed to be an expanding, and therefore an increasingly tenuous, debris cloud rather than a true exoplanet); this version is a supplement to the 2021 Handbook, with the upcoming 6.x.x series intended to support instead the 2022 Handbook

- [20210217T042710Z/5.1.1](#): Made minor adjustments to tabs and spacing for paragraphs before creating online PDF.
- [20210216T161213Z/5.1.0](#): Made minor adjustments (small points of syntax, spelling, punctuation, or similar, with much bibcode error correction). Added a long paragraph with five methods for retrieving a full-text, all-illustrations PDF from a typical astronomical bibcode citation. Corrected a mistake of astrophysical substance, in the subsection 4.8 discussion of onset-of-helium-core-fusion (violence in the onset of core-helium fusion is characteristic of the less massive, not of the more massive, incipient fusers-of-core-helium) This yielded a work sufficiently updated to support uploading to the online-version server.
- [20210128T145046Z/5.0.0](#): Made major revisions of the 4.0.0 version series, by adding several thousand words to the introductory online essay, with stellar-evolution background and a detailed briefing on the amateur-relevant “Be phenomenon” and “shell spectra” (and to a lesser extent by expanding “Remarks,” most notably for  $\alpha$  Eri (Achernar),  $\zeta$  Tau (Tianguan), and  $\alpha$  Aql A (Altair); other work on “Remarks” included routine updates for such things as binary position angles and celestial-sphere distances, and also comparison of our MK types against MK types as assigned by Astron. Alm. for epoch 2021.5, with the MK discrepancies logged). The work was not yet sufficiently polished to support uploading to the online-version server.
- [20200815T190800Z/4.0.0](#): Performed sufficient updating of the 3.x.x version series to support the print edition of the 2021 Handbook, but without sufficient updating to support uploading to the online-version server.
- [20191231T235959Z~/3.x.x series](#): Supplemented previous editions of this online publication in various ways, most notably by adding the (rather prolix) results of (rather detailed) primary-literature inspections for  $\alpha$  Cet Aa (Mira),  $\alpha$  Umi Aa (Polaris),  $\beta$  Per Aa1 (Algol),  $\alpha$  Tau A (Aldebaran),  $\epsilon$  Aur A (Almaaz),  $\alpha$  Ori Aa (Betelgeuse),  $\gamma$  Vel Aa,  $\alpha$  Leo A (Regulus),  $\alpha$  Vir Aa (Spica),  $\zeta$  Oph, and  $\alpha$  Lyr A (Vega).
- [20181231T235959Z~/2.x.x series](#): Supplemented the 1.x.x version series with some (rather detailed) primary-literature inspections for selected familiar bright stars, thereby expanding “Remarks.”

Star Name	RA (2023.5) Dec	$m_V$	$B-V$	MK Type	$\pi$ mas	$D$ ly	$\mu$ "/y	PA $^\circ$	$V_{rad}$ km/s	Remarks
$\alpha$ Sun And Aa	0 09.6 +29 13	-26.75 2.07	0.63 -0.11	G2 V B9p IV: (HgMn)	34	4.8 8 lm	0.214	140-12	SB2 <sup>†</sup>	Alpheratz  the SB2 components $\alpha$ And Aa, $\alpha$ And Ab (period 96.7 d) are now interferometrically measured, yielding orbital value $e = 0.5$ or $0.6$ ¶ slight variability, possibly of $\alpha$ CVn A (“ $\alpha^2$ CVn”) type, with range of mag. 0.04 in V, period 23.19 h (AAVSO(VSX) assessment in 2022: status flag = confirmed variable; variability classification symbol = “ACV:”)
$\beta$ Cas A	0 10.4 +59 17	2.27 <sup>†</sup>	0.34	F2 III	60	1.2 55	0.554	109 +12	SB <sup>†</sup>	Caph  slight var.: 2.25-2.29 in V, 0.1010 d second-brightest of the $\delta$ Sct variables (the brightest is $\alpha$ Aql A [Altair]) (AAVSO(VSX) assessment in 2022: status flag = confirmed variable;

$\gamma$	Peg A	0 14.4 +15 19	2.83 <sup>†</sup> -0.22	B2 IV	8	-2.6 400	0.009 168	+4 SB	<p>variability classification symbol = “DSCTC”</p> <p>¶ rapid rotator; <a href="#">2011ApJ...732...68C</a> finds the rotation to be &gt; 90% of breakup rate, and radius at poles to be ~24% less than radius at equator, with <math>\beta</math> Cas A of mass ~2 <math>M_{\odot}</math>, seen nearly pole-on; <math>\beta</math> Cas A is notable for being cooler than typical rapid rotators, lying just barely on the rapid side of the F5 “rotation break,” and additionally is notable for being old enough to have evolved off the MS, having in its MS career been an A star rather than an F star (generally, rotation slows as an aging star increases in radius: but our table of bright stars does harbour at least one other such evolved rapid rotator in type F, namely <math>\theta</math> Sco A); an envelope at this modest photospheric temperature is dominated by convection not only at the equator but even at the (~1000 K hotter) poles; consistently with this picture of an envelope everywhere convective, interferometry of <math>\beta</math> Cas A is found to yield results for low latitudes gravity darkening inconsistent with 1920’s von Zeipel law (the law is accurate only if an envelope is radiative); <a href="#">2011ApJ...732...68C</a> suggests that in its process of evolution off the MS (in which a core contracts, an envelope expands) <math>\beta</math> Cas A has been efficient in transferring angular momentum from core to envelope</p> <p>¶ <a href="#">2011ApJ...732...68C</a> Fig. 4 presents imaging (as a single star, not as a binary) of <math>\beta</math> Cas A, from CHARA interferometry</p> <p>¶ <math>\beta</math> Cas A is SB, of period 27 d, as yet unresolved, even in interferometry (so WDS is as yet unable to write “<math>\beta</math> Cas Ba,” “<math>\beta</math> Cas Bb”); NPOI interferometry has, however, succeeded in assigning a limb-darkened diameter, ~2.1 mas</p> <p><math>\gamma</math> Peg slight var., 2.82–2.86 in V, 3.64 h Algenib</p> <p><math>\gamma</math> Peg, as a multi-star assemblage, presents both <math>\beta</math> Cep-type and 53 Per-type variability: but we at the Handbook do not know how the combination of variabilities is distributed among the two stars that constitute the <math>\gamma</math> Peg A SB and the faint celestial-sphere neighbours <math>\gamma</math> Peg B (mag ~13) and <math>\gamma</math> Peg C (likewise mag. ~13)</p> <p>(AAVSO(VSX) assessment in 2022: status flag = confirmed variable; variability classification symbol = “BCEP+SPB”)</p>
$\beta$	Hyi <sup>†</sup>	0 26.9 -77 07	2.80 <sup>†</sup> 0.62	G1 IV	134.1	3.5 24.3	2.243 <sup>†</sup> 82	+23 <sup>†</sup>	<p>¶ E(B-V) = +0.01 possible exoplanet</p> <p>¶ included in NSV, but classified at AAVSO(VSX) as non-variable</p> <p>(AAVSO(VSX) assessment in 2022: status flag = confirmed non-variable; variability classification symbol = “CST”)</p> <p>¶ high space velocity (interloper from more remote galactic region?)</p>
$\alpha$	Phe	0 27.4 -42 11	2.39 <sup>†</sup> 1.09 <sup>†</sup>	K0 IIIb	38.5	0.3 ~85	0.426 147	+75 SB	<p><math>m_v</math>, B-V values are for the combined light of the SB system</p> <p>¶ AAVSO(VSX) situation in 2022: no status flag (so not confirmed variable, not suspected variable, not confirmed non-variable), a fortiori no variability classification symbol</p>
$\delta$	And <u>Aa</u>	0 40.6 +30 59	3.26 <sup>†</sup> 1.28 <sup>†</sup>	K3 III	~30.9	0.7 106	0.142 126	-7 SB <sup>†</sup>	<p>one of the rare instances in which a resolved SB has been resolved with direct imaging, rather than with interferometry</p> <p>(although angular diameter of <math>\delta</math> And Aa has been found through interferometry): <a href="#">2015ApJ...809...11B</a> reports a single 2014 observation via Palomar (5 m) Stellar Double Coronagraph, working in near IR with adaptive optics; angular separation was 0.4”, yielding Aa-to-Ab distance ~12 au, with <math>e</math>~0.5; since system is large, period is long (~20000 d?)</p> <p>¶ there seem (as of Sep. 2021) to be no reports of interferometric resolution of this SB</p> <p>¶ possible slight variability, range 0.1 in V:</p>

$\alpha$	Cas A	0 41.9 +56 40	2.22 <sup>†</sup>	1.17	K0 IIIa	~14.3	-2.0	230	0.060	122	-4 V?	Schedar
												<p>AAVSO(VSX) notes existence of NSV entry, but finds no records of AAVSO observations, and is not yet able either to assign a conjectural variability type or to deny variability (further photometric study advisable?) (AAVSO(VSX) assessment in 2022: status flag = suspected variable; no variability classification symbol)</p> <p>¶ <math>m_v</math>, B-V values are for the combined light of the <math>\delta</math> And Aa,Ab system</p> <p>¶ <math>\delta</math> And Ab (mag. 10.0) is asserted in <a href="#">2015ApJ...809...11BVVD</a> to be likely not a WD, as had been previously believed, but instead likely an MS star of MK type K (with the authors additionally noting, as one alternative possibility (in this case a remote alternative possibility), that <math>\delta</math> And Ab might be a pair of low-mass MS stars)</p> <p>AAVSO(VSX) millimag. variable (<i>TESS</i> mission; possibly rotational modulation, from starspots), period 2.41 d (AAVSO(VSX) assessment in 2022: status flag = confirmed variable; variability classification symbol = “<b>ROT</b>:”) ¶ limb darkening observed interferometrically (disk diameter 5.25 mas)</p> <p>¶ <math>\alpha</math> Cas D (mag. 9), PA 283°, 70" (2018), is mere line-of-sight coincidence, not gravitationally bound to <math>\alpha</math> Cas A</p>
$\beta$	Cet	0 44.8 -17 51	2.04 <sup>†</sup>	1.02	K0 III <sup>†</sup>	~33.9	-0.3	96	0.235	82	+13 V?	Diphda
												<p>anomalous in being X-ray-bright and yet a slow rotator</p> <p>¶ evolutionary status uncertain (helium core ignited already, or still contracting?)</p> <p>¶ slight variability (range 0.004 mag in V, period 167.79 d, with AAVSO(VSX) presently unable to assign a conjectural variability type (further photometric study advisable?) (AAVSO(VSX) assessment in 2022: status flag = confirmed variable; variability classification symbol = “<b>VAR</b>:”) ¶ Astron. Alm. (epoch 2021.5) assigns MK type G9 III CH-1 CN 0.5 Ca 1</p>
$\eta$	Cas <u>A</u> <sup>†</sup>	0 50.5 +57 56	3.44 <sup>†</sup>	0.57 <sup>†</sup>	G0 V <sup>†</sup>	168	4.6	19.4	1.222	117	+9 SB?	Achird
												<p>B:7.51, K4 Ve, 13.4", PA 62°→326°, 1779→2019 orbit 480 y; SB status has been asserted for <math>\eta</math> Cas A, and yet is said by WDS (as viewed 2021 Sept. 14) to be not confirmed; possibility of variability in <math>\eta</math> Cas system now discounted; our <math>m_v</math>, B-V values are for the combined light of <math>\eta</math> Cas AB (the corresponding values for <math>\eta</math> Cas B alone are 7.51, 1.39); overall field is crowded with dim stars, the brightest of which, apart from <math>\eta</math> Cas A and <math>\eta</math> Cas B, are the well-separated <math>\eta</math> Cas G (mag. 9.5; 420", PA 259° (2012); rectilinear-solution analysis of proper motions, 1852→2012, does not reveal any orbital motion) and the very widely separated <math>\eta</math> Cas H (mag. 8.5; 701", PA 355° (2012); analysis of proper motions covers only 1991→2012, and rectilinear-solution analysis of proper motions seems unavailable as of at any rate 2021 Sept. 14)</p> <p>¶ AAVSO(VSX) assessment in 2022: status flag = confirmed non-variable; variability classification symbol = “<b>CST</b>”</p> <p>¶ Astron. Alm. (epoch 2021.5) assigns MK type F9 V var.:1.6-3.0 (V); B:10.9, 2.1", PA 255°→259°, 1888→2002 orbit &gt;1500 y</p> <p>¶ first “Be phenomenon” discovery (Secchi, 1866); additionally the prototype for the <math>\gamma</math> Cas type of eruptive irregular variables; background on Be phenomena and <math>\gamma</math> Cas-type variability is given in <a href="#">www.aavso.org/vsots_gammacas:2002ASPC..279..22IH</a> summarizes the observational history, including major shell-spectrum phases in 1935-1936 and 1939-1940; despite its historical importance, however, Cas A cannot safely be taken as a typical “Be phenomenon” star, since it presents the peculiarity of hard thermal (and variable)</p>
$\gamma$	Cas A <sup>†</sup>	0 58.1 +60 51	2.39v <sup>†</sup> -0.13		B0 IVnpe (shell) <sup>†</sup>	5	-4.2	600	0.026	98	-7 SB	



$\alpha$	Eri	1 38.6	-57 07	0.45 <sup>†</sup>	-0.16	B3 Vnpe (shell?) <sup>†</sup>	23	-2.7	140	0.095	114	+16 V	<p>SB variability is of <math>\beta</math> Lyr type;  there is additionally  slow, irregular variability (unsurprising for a  binary containing a cool and evolved star)  (AAVSO(VSX) assessment in 2022:  status flag = confirmed variable;  variability classification symbol = “<b>EB/GS+LB</b>”)  ¶ Astron. Alm. (epoch 2021.5) assigns MK type  M0- IIIa</p> <p>ecl. slight var.: 0.40–0.46 in Hp, 1.263 d <b>Achernar</b>  AAVSO(VSX) does not assert ecl.,  but does assert the slight variability  typical of “Be phenomenon” stars  lacking outbursts (this is in many “Be phenomenon”  cases found to be <math>\lambda</math> Eri-type variability)  (AAVSO(VSX) assessment in 2022:  status flag = confirmed variable;  variability classification symbol = “<b>BE</b>”)  ¶ brightest of the “Be phenomenon” stars (but first  recognized as such only recently, ~1976); an active  Be phase that began between Dec. 2012 and early Jan.  2023 (after a period of inactivity, with <math>\alpha</math> Eri presumably  diskless, over the previous 7 years) was first noted in  amateur spectroscopy, in Brazil; Balmer  hydrogen-<math>\alpha</math> line indicates a slow, steady  buildup of the Be disk, over a period of  ~1.6 y, with polarization suggesting that disk was slightly  less dense in 2014 than it had been in 2013;  <a href="#">2017A&amp;A...601A.118D</a>, a case study of <math>\alpha</math> Eri, presents  for the first time in astrophysics images of a disk  forming around a Be-phenomenon star (with H-band (IR)  emission from the disk extending to an outer radius  of between 1.7 and 2.3 stellar equatorial radii, in good  agreement with current computations in the general  theory of the Be phenomenon); it is possible that plane  of <math>\alpha</math> Eri disk is inclined to plane of stellar equator; rapid  variations in polarization indicate that in addition to its  disk, <math>\alpha</math> Eri possesses rings, due to episodic ejections  of gas consignments from its photosphere  ¶ <math>\alpha</math> Eri is a notably rapid rotator (&lt; 2.1 d) within the  (currently small) population of stars interferometrically  resolved; period in or near the disk phases varies, either  because gas is injected (“decreted”) from photosphere  into Be disk or because Be-disk gas is  re-accreted onto photosphere;  interferometry as performed  when <math>\alpha</math> Eri is temporarily without its Be disk reveals  oblateness (cf e.g.  <a href="#">www.eso.org/public/unitedkingdom/news/eso0316</a>)  ¶ although <a href="#">2008A&amp;A...484L..13K</a> reports companion,  at Dec. 2007 angular separation &lt; 0.15”, WDS has  not, as of Nov. 2020, asserted binarity; the orbital  motion of this companion seems to not be correlated  with the repeated formation and disappearance  of the Be disk</p>	
$\tau$	Cet A+6P	1 45.2	-15 49	3.50	0.73	G8 V <sup>†</sup>		~274.0	5.7	11.9	1.921 <sup>†</sup>	296	-16 <sup>†</sup> V	<p>mass &lt; 1 <math>M_{\odot}</math> (unusual in Sample S, although  typical in Population P)  ¶ high space velocity, low metallicity: interloper from  thick galactic disk  ¶ AAVSO(VSX) notes existence of NSV entry,  notes record of many AAVSO observations,  states the possible mag. range as “3. 5-? V”;  AAVSO(VSX) assessment in 2022:  status flag = suspected variable;  no variability classification symbol assigned)  ¶ on original Frank Drake (1960) SETI target list  <b>[THIS STAR ONLY IN ONLINE VERSION OF TABLE]</b></p>
$\alpha$	Tri A	1 54.4	+29 42	3.42	0.49	F6 IV		52	2.0	63	0.234	177	-13 SB	<p>Mohtallah  slight variability (range 3.41–3.42 in V, period 1.74 d),  of the non-eclipsing, rotating-ellipsoid type  (AAVSO(VSX) assessment in 2022:  status flag = confirmed variable;  variability classification symbol = “<b>ELL</b>”)</p>
$\beta$	Ari $\underline{\Delta}$ <sup>†</sup>	1 55.9	+20 55	2.65v <sup>†</sup>	0.14	A4 or A5 Va <sup>†</sup>		56	1.4	59	0.148	138	-2 SB2 <sup>†</sup>	<p><math>\beta</math> Ari system possibly variable, 2.56–2.70 in V Sheratan  AAVSO(VSX) notes existence of NSV entry,  but does not find any record of  AAVSO observations, and is</p>

													not yet able either to assign a conjectural variability type or to deny variability (further photometric study advisable?) (AAVSO(VSX) assessment in 2022: status flag = suspected variable; no variability classification symbol assigned)
ε	Cas	1 56.1 +63 47	3.37 -0.16	B3 IV:pe (shell?) <sup>†</sup>	8	-2.2	400	0.037	121	-8	V		¶ β Ari B (mag. 5.2 in IR or Johnson R band or similar) is SB companion of β Ari A, and yet AB has also been resolved interferometrically as an exceedingly tight binary (15 measurements, 1988→2005, with two quite similar published orbital solutions [e is high in both solutions, at ~0.9; A-to-B distance is 0.08 au min, 1.2 au max; period is 107 d]) ¶ <a href="#">2003AJ...126.2048G</a> discusses MK type
													instance of “Be phenomenon”; but additionally, AAVSO(VSX) documents slight variability, possibly of the SX Ari type, with range 3.35–3.37 in V (AAVSO(VSX) assessment in 2022: status flag = confirmed variable; variability classification symbol = “ <b>SXARI:</b> ”) ¶ He-weak (cp α And, α Tel)
α	Hyi	1 59.4 -61 28	2.85 0.28	F0n III-IV <sup>†</sup>	45	1.1	72	0.265	84	+1	V		rapid rotator (< 30 h) ¶ metal-rich ¶ AAVSO(VSX) situation in 2022: no status flag (so not confirmed variable, not suspected variable, not confirmed non-variable), no variability classification symbol assigned
γ	And A	2 05.4 +42 26	2.26 <sup>†</sup> 1.37 <sup>†</sup>	K3 IIb <sup>†</sup>	9	-3.1	400	-0.065	-139	-12	SB		B: 5.0, B9 V, 9.6” (2019); C: 6.5, A0 V; BC 0.2” Almach BC orbit 63.7 y ¶ m <sub>v</sub> , B-V values are for γ And A; the combined-light values for γ And ABC are 2.11, 1.21 ¶ limb darkening observed interferometrically (disk 6.80 mas) ¶ Astron. Alm. (epoch 2021.5) assigns MK type K3- IIb ¶ AAVSO(VSX) assessment in 2022: status flag = suspected variable; variability classification symbol = “ <b>LB:</b> ”
α	Ari +IP	2 08.5 +23 34	2.01 <sup>†</sup> 1.15	K2 IIIab	~49.6	0.5	66	0.240	128	-14	SB		slight slow irregular variability, 2.00–2.03 in V <b>Hamal</b> (AAVSO(VSX) assessment in 2022: status flag = confirmed variable; variability classification symbol = “ <b>LB</b> ”) ¶ calcium weak
β	Tri	2 10.9 +35 06	3.00 <sup>†</sup> 0.14	A5 IV	26	0.1	130	0.154	105+10	SB2 <sup>†</sup>			period of this unresolved SB is 31.39 d, with orbit (at least two solutions published) rather elongated (e=0.4 or 0.5; inter-component distance possibly 0.17 au min, 0.42 au max) ¶ as of 2022 Jul. 29, AAVSO(VSX) notes existence of NSV entry, and states mag. range as “3.02-? V,” but finds no record of AAVSO observations, and is not yet able either to assign a conjectural variability type or to deny variability (further photometric study advisable?) (AAVSO(VSX) assessment in 2022: status flag = suspected variable; no variability classification symbol assigned) ¶ IR excess (circumstellar matter? possible signature of planetesimals)
o	Cet <u>Aa</u> <sup>†</sup>	2 20.5 -2 52	4.95v <sup>†</sup> 1.55	M5-10 IIIe <sup>†</sup>	11 <sup>†</sup>	1.7	300 <sup>†</sup>	0.238	178	+64	V		LPV, 2-10.1; next max expected 2022 mid-July Mira <sup>†</sup> recent o Cet Aa maxima Nov. 2019 (V~2.3), Sep.-Oct. 2020 (V~3.0), Aug. 2021 (V~2.6), AAVSO reports V band mag. 8.06 on 2022 Jan. 31: <a href="#">2009ApJ...691.1470T</a> discusses variability, including variation in dominant (333 d) pulsation period and the question of longer-period variations (AAVSO(VSX) assessment in 2022: status flag = confirmed variable; classification symbol = “ <b>M</b> ”) ¶ times of maxima are, and times of minima are not, independent of wavelength: minima are at least coarsely correlated with maximum diameter of o Cet Aa ¶ prototype of the o Ceti-type variables, mass ~1 M <sub>⊙</sub> : the first O-rich AGB star with a CI detection



(2018A&A...612L...11S)

¶ physical radius ~2 au in visual, ~4 au in IR, still greater upon taking instead the “radio photosphere,” which itself increases in radius as progressively longer radio wavelengths are selected: [2015ApJ...808...36M](#) draws parallels with  $\alpha$  Ori Aa, attributing radio inhomogeneities in both cases to convective cells (and cf also [2016A&A...592A...42K](#), which summarizes some recent radio work)

¶ Ab (VZ Cet) WD, 10.4, 0.5", orbit ~500–600 y; what is seen as o Cet Ab (with professional equipment), and indeed probably seen as itself variable (over and above the long-period variability o Cet Aa) may be not the WD itself, but an accretion disk around the WD, captured from the o Cet Aa wind

¶ nearest instance of (weak) symbiotic binarity, and the only symbiotic to be observed in all wavelength regimes from X-ray to (mm, also cm) radio; interferometry (in IR) is available from VLT, and *CASSINI* has yielded (via Saturn-ring occultations) tomographically recovered imaging ([2016MNRAS.457.1410S](#));

*GALEX* has found bow shock, tail (length 13 ly) in ISM: mass-loss rate ~2.5e-7  $M_{\odot}$ /y; asymmetric atmosphere is discussed in [2016MNRAS.457.1410S](#)

¶ [2018A&A...620A...75K](#) reports dust trail linking Aa,Ab (consistently with other reports of Aa-to-Ab mass transfer)

¶ [2016A&A...592A...42K](#) ([2017A&A...599A...59K](#)) discusses o Cet Aa dust nucleation generally, with reference to aluminum (resp. titanium) species: in o Cet Aa, it is silicates that dominate the spectrum (in contrast with less-evolved stars, in which alumina features are spectrally dominant);

[2016A&A...590A.127V](#) discusses SiO gas,

o Cet Aa inner dust shells: it seems still an open question whether o Cet Aa dust formation is cyclic, as part of the photometrically evident pulsation cycle, or proceeds independently of the pulsations

¶ X-ray emission from o Cet Aa was reported in 2005 ([2005ApJ...623L.137K](#), as the first X-ray detection from an AGB star), and OH, SiO maser emission has also been reported (cf, e.g. [2017MNRAS.468.1703E](#)); further,

[2015A&A...577L...4V](#) asserts a hot spot, proposing magnetic activity as the cause

¶ [2016A&A...590A.127V](#) summarizes history of modelling: models generally agree that near o Cet are alternating circumstellar layers of infall and outflow, and that at greater radii is an accelerating outflow, from dust-driven winds: recent observations have tended to agree with overall results from running CODEX (e.g. [2014A&A...565A.119S](#))

¶ [2016A&A...586A...69P](#) discusses discrepancies in distance determinations (350 ly, 380 ly, 340 ly, and (least reliable?) *HIPPARCOS* 300 ly)

¶ Aa,Ab orbit would, if better known, yield improved total mass of Aa,Ab system, thereby advancing the overall theory of AGB stars

¶ protoplanetary disk was detected around Ab in 2007

¶ Astron. Alm. (epoch 2021.5) assigns MK type M5.5–9e III

¶ Fabricius noted variability in 1956; Hevelius proposed the name Mira in 1642

¶ for entry-level briefing-with-bibliography, cf [www.aavso.org/vsots\\_mira2 Updating](http://www.aavso.org/vsots_mira2 Updating)

[www.aavso.org/vsots\\_mira](http://www.aavso.org/vsots_mira); and for summary of recent primary literature, cf first section of [2016MNRAS.460..673N](#)

B: 6.23, 2.0", PA 283°→299°, 1825→2015 Kaffaljidhma orbit  $\geq$  320 y

¶  $m_v$ , B–V values are for the combined light of  $\gamma$  Cet AB

¶ AAVSO(VSX) situation in 2022: no status flag (so not confirmed variable, not suspected variable, not confirmed non-variable), a fortiori no variability classification symbol assigned)

B: 4.35, A1 Va, 8.2", PA 82°→90°, 1835→2020 **Acamar**

¶  $m_v$ , B–V values are for the combined light of  $\theta$  Eri AB

¶ AAVSO(VSX) situation in 2022: no status flag (so not confirmed variable,

$\gamma$  Cet A<sup>†</sup> 2 44.5 +3 20 3.46<sup>†</sup> 0.09<sup>†</sup> A2 Va 41 1.5 80 0.207 225 –5 V

$\theta$  Eri A 2 59.2 –40 13 2.91<sup>†</sup> 0.12<sup>†</sup> A5 IV 30 0.5 100 0.057 293 +12 SB2

$\alpha$	UMi <u>Aa</u> <sup>†</sup>	3 02.2	+89 22	2.01 <sup>†</sup>	0.60 <sup>†</sup>	F5–8 Ib	7.5 <sup>†</sup>	–3.6	430 <sup>†</sup>	–0.046	–105?	–17 SB	<p>not suspected variable, not confirmed non-variable), a fortiori no variability classification symbol assigned) slight Cep. var. 4.0 d; B: 9.1, F3 V, 18.4" (2016) <b>Polaris</b> the brightest of the Cepheids, but not a classical Cepheid, matching instead the “s-Cepheid” light-curve phenomenology of <a href="#">1995A&amp;A...303..137B</a></p> <p>¶ AAVSO(VSX) as viewed 2022 Jul. 13, gives V-mag. range 1.97–2.00, period ~3.9696 d; period is increasing ~4.4–4.9s/y, with sudden change around 1963, and with <i>COROLIS</i> satellite suggesting a further recent change: period change is often in Cepheid theory linked to evolution, but this may not be the whole story here (in particular, pulsation-driven mass loss through stellar wind, as affirmed by some recent authors (denial also published) would increase the rate of period change)</p> <p>(AAVSO(VSX) assessment in 2022: status flag = confirmed variable; variability classification symbol = “<b>DECPS</b>”)</p> <p>¶ pulsation mode (1st overtone? 2nd? fundamental?), evolutionary history (1st crossing of IS? or 3rd crossing?), and distance are controverted by various 2010-through-2018 authors (we here use <i>Gaia</i> DR2 distance for <math>\alpha</math> UMi B as a proxy, assuming with the current literature that B is indeed gravitationally bound with Aa,Ab)</p> <p>¶ <math>\alpha</math> UMi Aa is first Cepheid with mass determined through purely dynamical means (via the Aa,Ab orbit: Aa is single-lined SB, and Aa, Ab have been resolved with HST, as first announced (0.17") in <a href="#">2008AJ...136..137E</a> [orbit ~30 y])</p> <p>¶ <a href="#">2000A&amp;A...360..399VV</a> warns that peculiarity of the <math>\alpha</math> UMi Aa light curve makes this particular Cepheid perhaps not a suitable anchor point for determining the overall Cepheid period-luminosity relation (essential though a determination of that relation is for calibrating the cosmic distance scale)</p> <p>¶ <math>\alpha</math> UMi Aa is important as a case study in the “Cepheid mass discrepancy” problem (Cepheid masses deduced from pulsation periods are found to be too low in comparison with masses from stellar-evolution modelling)</p> <p>¶ strictly a three-star system, UMi Aa+UMi Ab+UMi B; Aa,Ab has period 29.6 y, inter-component distance 6.7 au min, 27 au max, 17 au average (orbital solution has been published); B experiences Aa,Ab as essentially a point mass, with period <math>\geq 42,000</math> y, separation at least 2400 au; B is mag. 9.1, at angular separation 18" (no orbital solution published)</p> <p>¶ <math>m_v</math>, B–V values are for the combined light of UMi Aa,Ab,B; the corresponding values for <math>\alpha</math> UMi B are 8.20, 0.49</p> <p>¶ <math>\alpha</math> UMi Aa, Ab,B is approaching NCP: closest approach will be 14', in ~2105</p> <p>¶ B has <math>E(B-V)=0.0</math></p>
$\alpha$	Cet	3 03.5	+4 11	2.53 <sup>†</sup>	1.64	M2 III <sup>†</sup>	13	–1.9	250	0.078	188	–26 V	<p>slight var., 2.45–2.54 in V <b>Menkar</b></p> <p>variability is possibly of a slow-irregular type (AAVSO(VSX) assessment in 2022: status flag = confirmed variable; variability classification symbol = “<b>LB:</b>”)</p> <p>¶ radio source (due to stellar wind)</p> <p>¶ notably deficient in carbon</p> <p>¶ Astron. Alm. (epoch 2021.5) assigns MK type M1.5 IIIa</p>
$\gamma$	Per <u>Aa,Ab</u> <sup>†</sup>	3 06.5	+53 36	2.93v <sup>†</sup>	0.70	G8 III <sup>†</sup> + A2? V?	13	C1.5	240	0.006	175	+3 SB2 <sup>†</sup>	<p>composite spectrum, orbit 14.6 y, next eclipse 2035 eclipse duration &lt; 2 weeks; eclipse variation is significantly above threshold of naked-eye detection, with AAVSO(VSX) giving V-mag. range 2.91–3.21 (ranges in Johnson U and B bands are still larger); this is a system of the <math>\beta</math> Per type (AAVSO(VSX) assessment in 2022: status flag = confirmed variable; variability classification symbol = “<b>EA/GS</b>”)</p> <p>¶ angular separation of Aa,Ab is always <math>\leq 0.3''</math>, putting this binary, for part of its orbit, just within the limits of feasibility for traditional micrometer astrometry; orbit is highly elliptical, with <math>e=0.79</math> on the better of two published orbital solutions;</p>

$\rho$	Per	3 06.7 +38 56	3.39v <sup>†</sup> 1.65	M4 II	11	-1.6 310	0.167 129	+28	<p>amenability of this binary to spectroscopic orbit solution (via radial velocities: radial-velocity data are available since ~1900) and to interferometry make it a useful candidate for stellar-mass studies, and therefore a useful test for theoretical predictions of luminosity classes and spectral types from given masses (<a href="#">1999A&amp;A...348.127P</a>)</p> <p>¶ Aa is mag. ~3.6 and Ab is mag. ~3.8; our <math>m_v</math>, B-V values are for <math>\gamma</math> Per Aa,Ab combined light</p> <p>¶ Astron. Alm. (epoch 2021.5) assigns MK type G5 III</p> <p>semiregular variable: 3.3–4 in V, ~50 d, ~120 d, ~250 d rough period ~50 d, with possibly also longer (rough?) period(s)</p> <p>(AAVSO(VSX) assessment in 2022: status flag = confirmed variable; variability classification symbol = “<b>SRB</b>”)</p> <p>Aa=compos. spectrum Aa1,2 ecl.,2.09-3.30 in V, 2.9 d Algol Aa2 is K2IV?</p> <p>¶ in older terminology, <math>\beta</math> Per Aa1 = <math>\beta</math> Per A, <math>\beta</math> Per Aa2 = <math>\beta</math> Per B, <math>\beta</math> Per Ab = <math>\beta</math> Per C; but WDS, following the current terminology, uses the “B” and “C” for other purposes (since there are optical neighbours B,C,D,E,F,G,H; all are between 5” and 100” from the Aa1,Aa2,Ab triple, and all are fainter than mag. 10); system is hierarchical, with outlying Ab experiencing the close Aa1,Aa2 pair (Aa1–Aa2 distance is just 14.14 <math>R_{\odot}</math>) as essentially a point mass; angular separation between Aa1,Aa2 and Ab is ~0.1” (WDS 1973, 2010); orbital solutions have been published both for the tight binary that is <math>\beta</math> Per Aa1,Aa2 and for the wider binary that is <math>\beta</math> Per Aa,Ab</p> <p>¶ among the most visually prominent of the eclipsing binaries, and for theoreticians the most familiar of the semidetached binaries (i.e., binaries in which one of the two Roche equipotential surfaces is fully occupied, the other not)</p> <p>(AAVSO(VSX) assessment in 2022: status flag = confirmed variable; variability classification symbol = “<b>EA/SD</b>”)</p> <p>¶ Aa2 is tidally locked, in a rapid circular orbit with Aa1; the consequent rapid spin of Aa2 causes dynamo action in Aa2 convection zone, with Aa2 consequently having complex magnetosphere (mass-transfer stream possibly even deflected out of Aa1,Aa2 orbital plane by magnetics; <a href="#">2012ApJ...760...8R</a>; Aa2 has additionally a meridional coronal loop, approximately as high as the diameter of Aa2 (the size exceeds what has been anticipated from modelling) believed pointing at all times to Aa1), X-ray emission, varying radio morphology (double-lobed when radio-brightest) and CME episodes (<a href="#">2017ApJ...850..191M</a>) suggests the 1997 Aug. 30 superflare event supplies “arguably the best candidate” for a non-solar CME)</p> <p>¶ the (unsteady) Aa2-to-Aa1 mass transfer, while ongoing, and indeed responsible for an annulus around Aa1, is no longer copious (in contrast with the copious transfer still present in, e.g. <math>\beta</math> Lyr)</p> <p>¶ it is not the (now modest) unsteady mass transfer, but possibly instead the Applegate mechanism (<a href="#">1992ApJ...385..621A</a>), implicating a stellar magnetic activity cycle, which dominates the Aa2,Aa1 period variation (increase-decrease-increase cycle, not quite strictly periodic, 32 y; there are additionally period modulations of 1.9 y and 180 y; as viewed 2021 Jan. 28, AAVSO(VSX) asserts period 2.86736 d,</p>
$\beta$	PerAa1 <sup>†</sup>	3 09.7 +41 03	2.12v <sup>†</sup> -0.05	B8 V <sup>†</sup>	36	-0.1 90	0.003 119	+4 SB	

but as viewed 2022 Mar. 3 gives updated value 2.86734 d);  
 full amplitude  
 of the Aa1 Aa2 period variation is ~0.8 s;  
 such alternating period changes  
 in binaries are still not, however,  
 well understood  
 ¶ it is the (several My ago rapid and  
 copious) mass transfer  
 that resolves the “Algol paradox”  
 of a lower-mass more evolved (in this case, sub-giant)  
 star in orbit with a higher-mass  
 less evolved (indeed MS) star;  
 masses are well known in this particular case:  
[2015MNRAS.451.4150K](#), having disentangled the  
 $\beta$  Per Aa1, Aa1, Ab spectra, determines their  
 masses within  $\pm 2\%$ ,  
 corroborating [2012ApJ...752...20B](#)  
 ¶  $\beta$  Per Aa2 elemental abundances below corona and flare  
 (investigated in [2015MNRAS.451.4150K](#))  
 are of special  
 interest, since mass transfer has  
 stripped off Aa2 outer layers,  
 opening the Aa2 interior to spectroscopic inspection  
 ¶ [1983ApJ...273L..85K](#) reports discovery of  
 Chandrasekhar eclipse-induced stellar limb  
 polarization from  $\beta$  Per Aa1, in a wide optical passband  
 ¶ MK type K2 IV is assigned to Aa2  
 in at least 3 recent papers,  
 whereas the older [1993ApJ...410..808L](#) has the  
 slightly hotter MK type K0 IV; what is essential  
 here is the agreed “IV” (as opposed to “V”), indicating  
 evolution of this (secondary) star off the MS  
 (and Astron. Alm. (epoch 2021.5) assigns MK type  
 B8 V for Aa1, as we do here, and for one companion  
 (is this Aa2, or is it Ab?) the  
 uncertainty-flagged “F;,” without luminosity class)  
 ¶  $\beta$  Per Ab, spectrally  
 Am with some F1V characteristics,  
 orbits the  $\beta$  Per Aa1,Aa2 binary with  
 period ~680 d, without eclipsing  
 ¶ [2012ApJ...752...20B](#)  
 presents CHARA interferometry  
 (~0.5 mas, H (near-IR) band)  
 of the  $\beta$  Per Aa1,Aa2,Ab system (finding orbital plane  
 of Ab to be nearly perpendicular to Aa1,Aa2 orbital plane),  
 and also summarizes earlier interferometry;  
 an approx 55-frame animation from this paper  
 can be conveniently viewed at  
[en.wikipedia.org/wiki/Algol](http://en.wikipedia.org/wiki/Algol)  
 ¶  $E(B-V)=+0.03$   
 ¶ [2013ApJ...773...11](#) suggests that  $\beta$  Per system variability  
 is documented in the “Cairo  
 Calendar” papyrus (New Kingdom,  
 dated to 1271–1163 BCE); al-Sufi  
 (Persia, ca 964 CE) is, however, silent  
 on question of  $\beta$  Per variability  
 ¶ AAVSO has briefing notes, with some history,  
 at [www.aavso.org/vsots\\_betaper](http://www.aavso.org/vsots_betaper);  
[1910ApJ...32..185S](#) is the discovery paper  
 for  $\beta$  Per Aa1,Aa2 secondary minimum, from the  
 beginnings of photoelectric-cell photometry;  
[1998A&AT...15..357P](#) analyzes “Algol paradox” history;  
[arxiv.org/pdf/astro-ph/0611855.pdf](http://arxiv.org/pdf/astro-ph/0611855.pdf)  
 “Appendix B.,” is a tabular history of  
 $\beta$  Per-pertinent investigations  
 from antiquity to 1999; in this  
 same K. Wecht 2006 Lehigh Univ  
 PhD thesis, Table 2.5.1  
 summarizes 1966-through-1983 observational coverage,  
 as tabulated in the less  
 Web-accessible 1986 work of Budding  
 in open cluster Melotte 20, a.k.a. Collinder 39 **Mirfak**  
 near edge of HR diagram IS (slightly too  
 hot to be a straightforward Cepheid);  
 entered into NSV as suspected variable;  
 AAVSO(VSX) notes many AAVSO observations,  
 shows range 1.72–1.86 in V,

$\alpha$  Per A 3 26.0 +49 57 1.79<sup>†</sup> 0.48<sup>†</sup> F5 Ib ~6.4 -4.2 510 0.035 138 -2 V

δ	Eri	3 44.4	-9 41	3.53 <sup>†</sup>	0.92	K0 IV	111	3.7	29.5	0.749	353	-6	and indicates existence of NSV entry, but is not yet able either to assign a conjectural variability type or to deny variability (further photometric study advisable?) (AAVSO(VSX) assessment in 2022: status flag = suspected variable; no variability classification symbol assigned) ¶ m <sub>v</sub> , B-V values seem unavailable in GCPD; we take values from <a href="#">2002yCat.2237...0D</a> earlier suggestion of variability is now discounted (AAVSO(VSX) assessment in 2022: status flag = confirmed non-variable; classification symbol = "CST")
δ	Per Aa	3 44.6	+47 52	3.01 <sup>†</sup>	-0.12	B5 IIIIn <sup>†</sup>	6	-3.0	500	0.050	149	+4 SB	[THIS STAR ONLY IN ONLINE VERSION OF TABLE] slight var., possibly of γ Cas type, 2.99-3.04 in V (but WDS indicates possible variability of the type presented by α CVn A ("α <sup>2</sup> CVn")) (a star discussed later in this table); AAVSO(VSX) asserts variability, and assigns γ Cas variability as a possible-and-yet-not-certain variability type (further photometric study advisable?) (AAVSO(VSX) assessment in 2022: status flag = confirmed variable; classification symbol = "GCAS:") ¶ cluster affiliation is controverted ¶ Astron. Alm. (epoch 2021.5) assigns MK type B5 III ¶ E(B-V) = +0.04
γ	Hyi	3 46.9	-74 10	3.24 <sup>†</sup>	1.61	M2 III	15.2	-0.8	~214	0.126	24	+16	slight semireg. var., 3.22-3.29 in V (AAVSO(VSX) assessment in 2022: status flag = confirmed variable; classification symbol = "SRB") ¶ evolutionary status is uncertain
η	Tau Aa	3 48.9	+24 11	2.87	-0.09	B7 IIIIne <sup>†</sup>	8	-2.6	400	0.048	156+10	V? SB	brightest member of Pleiades Alcyone ¶ m <sub>v</sub> , B-V values seem unavailable in GCPD; we take values from <a href="#">2002yCat.2237...0D</a> ¶ η Tau Aa,Ab is known as a binary both from spectroscopy and from occultation (but this tight pair seems not to have been resolved interferometrically, at any rate as of 2021 Oct. 25; it is therefore on the strength of occultation alone that WDS is able to write "η Tau Aa," "η Tau Ab"), with Ab of mag. 4.6; additionally, in this crowded part of the celestial sphere, WDS catalogues as neighbours of the tight Aa,Ab binary B, C, Da, Db, E, F, G, H; of these neighbours to Aa, not only Ab, but also B, C, Da, and Db are brighter than mag. 10, and possess astrometric data from as early as the 19th century (except that the tight Da,Db pair was first resolved in the 21st century, through speckle interferometry); <a href="#">1972JBAA...82.431K</a> describes the 18.6-year 1940-through-2050 cycle of lunar occultation possibilities ¶ rapid rotator (2.29 d?), with "Be" and "shell-spectrum" phenomena (BS5: "rotationally unstable"), making this star an appropriate target for periodic low-cadence (e.g. once-yearly) amateur-spectroscopy monitoring ¶ the overall η Tau assemblage is known to present both rotational (spotted) and slowly-pulsating-B variability, at the millimagnitude level (AAVSO(VSX) assessment in 2022: status flag = confirmed variable; variability classification symbol = "ROT+SPB")
ζ	Per A <sup>†</sup>	3 55.6	+31 57	2.87 <sup>v?</sup>	0.11 <sup>†</sup>	B1 Ib	4	-4.0	800	0.011	150	+20 SB	¶ significant dimming by ISM dust; E(B-V) = +0.03 B: 9.16, B8 V, 12.8", PA 205° → 208°, 1824 → 2020 orbit ≥ 50,000 y ¶ since the SB that is ζ Per A is not as yet resolved, even in interferometry, WDS is not as yet able (at any rate as of 2021 Oct. 25) to write "ζ Per Aa," "ζ Per Ab"; ζ Per B experiences ζ Per Aa,Ab as essentially a point mass, and is too slow in its orbit to yield spectroscopic (radial-velocity)

$\gamma$	Eri A	3 59.1	-13 27	2.95 <sup>v†</sup>	1.59	M1 IIIb <sup>†</sup>	16	-1.0	200	0.129	151	+62	<p>data; further, WDS gives, as celestial-sphere neighbours of the wide and slow <math>\zeta</math> Per Aa,Ab-B pairing, <math>\zeta</math> Per C, D, and E (with the A,E angular separation, as measured in 2015, wide, at 120")</p> <p>¶ <math>m_v</math>, B-V values are for <math>\zeta</math> Per A; GCPD gives the corresponding values for <math>\zeta</math> Per AB combined light as 2.85, 0.12 (and for <math>\zeta</math> Per B, gives 9.16 and 0.23; and for <math>\zeta</math> Per D, gives 9.90 and 0.33; further, for the elusive <math>\zeta</math> Per E gives 10.35, 0.71)</p> <p>¶ as of 2022 July 29, AAVSO(VSX) notes existence of NSV entry, notes existence of more than 30 AAVSO observations, and gives possible range 2.80–2.93 in V (further photometric study advisable?) (AAVSO(VSX) assessment in 2022: status flag = suspected variable; no variability classification symbol assigned)</p> <p>¶ significant dimming by ISM dust; E(B-V) = +0.33 (pronounced reddening) slight var. (2.88–2.96 in V), possibly slow-irreg. Zaurak (AAVSO(VSX) assessment in 2022: status flag = confirmed variable; variability classification symbol = "LB:")</p> <p>¶ Ca, Cr weak</p> <p>¶ further photometric and spectroscopic studies advisable? (Kaler, at <a href="http://stars.astro.illinois.edu">stars.astro.illinois.edu</a>, writes, "must be one of the least-studied of the cooler bright stars")</p> <p>¶ Astron. Alm. (epoch 2021.5) assigns MK type M0.5 IIIb Ca-1</p>
$\varepsilon$	Per A <sup>†</sup>	3 59.4	+40 05	2.89 <sup>†</sup>	-0.18	B0.5 IV <sup>†</sup>	5	-3.6	600	0.028	149	+1 SB2	<p>B: 8.9, B9.5 V, 9.1", PA 10°→10°, 1821→2015 orbit <math>\geq</math> 16,000 y; since the SB2 has not yet been resolved, even in interferometry, WDS is not as yet able (at any rate as of 2021 Oct. 25) to write "<math>\varepsilon</math> Per Aa," "<math>\varepsilon</math> Per Ab"; <math>\varepsilon</math> Per B experiences the unresolved SB2 that is <math>\varepsilon</math> Per A as essentially a point mass, and is too slow in its orbit to yield spectroscopy (radial-velocity) orbital data; further, WDS gives, as celestial-sphere neighbours of the wide and slow <math>\varepsilon</math> Per AB pairing, <math>\varepsilon</math> Per C and D, with D a little brighter than mag. 10 (at mag 9.25), and at the wide angular separation of 163" from <math>\varepsilon</math> Per A; the unresolved SB2 that is <math>\varepsilon</math> Per A does have published orbital solutions, with period 14.069 d (indicating a tight pairing), and with <math>e=0.5</math></p> <p>¶ slight variable, of <math>\beta</math> Cep type (2.89–2.91 in V; AAVSO(VSX) assessment in 2022: status flag = confirmed variable; variability classification symbol = "BCEP"); one of the most extreme spectroscopic variables (periods 2.27 h and 8.46 h)</p> <p>¶ E(B-V) = +0.10</p>
$\lambda$	Tau	4 02.0	+12 33	3.49 <sup>v†</sup>	-0.12	B3 V	7	-2.4	480	0.017	209+18	SB2 <sup>†</sup>	<p>ecl.: 3.37–3.91 in V, 3.953 d; secondary is A4 IV system is of <math>\beta</math> Per type; AAVSO(VSX) as viewed 2022 July 08 gives period 3.9529478 d, eclipse duration 14.231 h (AAVSO(VSX) assessment in 2022: status flag = confirmed variable; variability classification symbol = "EA/DM")</p> <p>¶ shape distortion (mutual tides), reflection effect, some evidence of mass transfer</p>
$\alpha$	Ret A	4 14.7	-62 25	3.33	0.92	G8 II–III	20.2	-0.1	162	0.065	40	+36 SB?	<p>AAVSO(VSX) situation in 2022: no status flag (so not confirmed variable, not suspected variable, not confirmed non-variable), a fortiori no variability classification symbol assigned</p>
$\varepsilon$	Tau Aa +1P <sup>†</sup>	4 30.0	+19 14	3.53 <sup>†</sup>	1.01 <sup>†</sup>	K0 III <sup>†</sup>	22.2	0.3	150	0.113	110	+39 V?	<p>in Hyades; Aa,Ab 0.2" (2005), mags. ~3.6, ~6.0 as of 2021 Oct. 26, WDS records just one (interferometric) measurement of the <math>\varepsilon</math> Tau Aa,Ab binary</p> <p>¶ <math>m_v</math>, B-V values seem unavailable in GCPD; we take <math>m_v</math> value from <a href="http://2002yCat.2237...0D">2002yCat.2237...0D</a></p>

and calculate B–V from [2009ApJ...694.1085V](#) with [2002yCat.2237....0D](#)

¶ AAVSO(VSX) situation in 2022: no status flag (so not confirmed variable, not suspected variable, not confirmed non-variable), a fortiori no variability classification symbol assigned  
 ¶ metals-rich  
 ¶ first known instance of a planet-host in an open cluster; unusually massive among the currently known planet-hosts  
 ¶ Astron. Alm. (epoch 2021.5) assigns MK type G9.5 III CN0.5

[THIS STAR ONLY IN ONLINE VERSION OF TABLE]

system Aa-plus-Ab is a.k.a.  $\theta^2$  Tau; in Hyades Chamukuy companion in elongated orbit, with published orbital solutions (period is 140.728 d;  $e=0.73$ ; 0.23 au min, 1.3 au max); we here state mag. for Aa,Ab combined light (separately these are mag.  $\sim 3.7$ , mag.  $\sim 4.9$ , making Aa alone a little fainter than what this table consider a “bright star,” even though the naked-eye bright point that is  $\theta$  Tau Aa,Ab is the brightest Hyades member; one of the components in the Aa,Ab pair, typically presumed to be Aa, is a  $\delta$  Sct variable, with V range somewhat less than 0.1 mag., for which 12 periods are known, 1.64 h to 2.22 h, the ranges in some cases small (0.5 millimag., 30 millimag.); this is one of the intensely studied cases of  $\delta$  Sct variability); the SB system  $\theta$  Tau Aa,Ab forms a wide pair with the bright SB system  $\theta$  Tau Ba,Bb, a system a.k.a.  $\theta^1$  Tau (mag.  $\sim 3.9$ ; astrometry of Aa,Ab with respect to Ba,Bb is  $340'' \rightarrow 348''$ , PA  $346^\circ \rightarrow 347^\circ$ , 1800 $\rightarrow$ 2016); not only the Aa,Ab but also the Ba,Bb pair has a published orbital solution; Aa,Ab and Ba,Bb are in turn gravitationally bound, with each of the two tight pairs in this quadruple of stars experiencing the other tight pair as essentially a point mass; the entire Aa,Ab,Ba,Bb system, notably including at least three of the four individual masses, has been much studied since the 1990s, drawing on data from occultation, spectroscopy, and interferometry, even though the various challenges include some troublesome rotational broadening of spectral lines, since both Aa and Ab are rapid rotators; determination of masses, plus (helpfully, even *HIPPARCOS*-independent, via orbit model) determination of distances, confers on this Aa,Ab,Ba,Bb system, as on several other stars in the Hyades, a specially enhanced stellar-astrophysics significance, as isochrone-anchored data points for plotting the empirical (i.e. the theory-free) mass-luminosity relation, and therefore for constraining models of stellar evolution ([1997ApJ...485..167T](#) recapitulates the strategic position as follows: “The Hyades is unique in this respect. In no other case have dynamical masses been determined over a range covering much of the Main Sequence /.../ for stars of the same age and known chemical composition.”)

¶  $m_v$ , B–V values seem unavailable in GCPD; we instead use [1993A&AS..100..591O](#)

¶ AAVSO(VSX) assessment in 2022: status flag = confirmed variable; variability classification symbol = “DSCTC+E:”

¶ the  $\theta$  Tau Aa name “Chamukuy,” IAU-official since 2017, is the Yucatec Mayan name for a small bird

A: 3.8; B: 4.3, B9 IV; 0.2" (2019); orbit 12 y orbit very elongated, with A-to-B distance 1.9 au min, 17.5 au max

¶  $m_v$ , B–V values are for  $\alpha$  Dor AB combined light  
 ¶  $\alpha$  Dor system present slight variability, 3.236–3.276 in V, period 2.94247 d ([en.wikipedia.org/wiki/Alpha\\_Doradus](#) has light curve, from *TESS*), of the  $\alpha$  CVn A (“ $\alpha^2$  CVn”) type (AAVSO(VSX) assessment in 2022:

$\theta$  Tau Aa<sup>†</sup> 4 29.9+15 55 3.41<sup>†</sup> 0.19<sup>†</sup> A7 III 22 0.1 150 0.112 104 +40 SB<sup>†</sup>

$\alpha$  Dor A<sup>†</sup> 4 34.5 –55 00 3.26v<sup>†</sup>–0.10<sup>†</sup> A0p V: (Si)<sup>†</sup> 19 –0.3 169 –0.059 ~79? +26

$\alpha$	Tau A	+1P <sup>†</sup> 4 37.2 +16 33	0.87	1.54	K5 III <sup>†</sup>	49	-0.7	67	0.199	161	+54 SB	<p>status flag = confirmed variable;  variability classification symbol = “ACV”  ¶ Astron. Alm. (epoch 2021.5) assigns MK temperature type A0p Si (and does not assign an MK luminosity class)  slight irregular var., 0.86–0.89 in V Aldebaran  (AAVSO(VSX) assessment in 2022:  status flag = confirmed variable;  variability classification symbol = “LB:”);  recent literature proposes oscillations,  and also proposes rotational modulation from  modest photospheric-activity features  (with possibly an activity cycle (<a href="#">2015A&amp;A...580A.31H</a>):  the features could be (cool)  starspots, but could alternatively  be large convection cells;  the general topic of activity  in K giants is not yet well understood)  ¶ foreground star, not true Hyades member;  among the nearest of the red giants; evolution has  proceeded beyond the “FDU”  stage that accompanies helium-core contraction on RGB  ¶ 49 lunar occultations occurred over the period  2015 Jan. 29/2018 Sep. 03 (and yet there is a surprisingly  large scatter in the occultation  determinations of <math>\alpha</math> Tau angular diameter;  <a href="#">1972JBA...82..43IK</a> describes the overall 18.6-year  1940-through-2050 cycle of lunar occultation possibilities)  ¶ in contrast with its celestial-sphere neighbour  <math>\alpha</math> Ori, <math>\alpha</math> Tau is of modest mass (with recent literature  variously offering <math>\sim 1.2 M_{\odot}</math>, <math>\sim 1.3 M_{\odot}</math>, <math>\sim 1.5 M_{\odot}</math>):  Appendix C of <a href="#">2018ApJ...865L..20F</a>  tabulates values for mass,  luminosity, radius, age, and several other parameters,  on the strength of five separate  2008-through-2012 spectroscopy investigations  ¶ <a href="#">2013A&amp;A...553A...3O</a> reports “MOLsphere” (molecule-  harbouring atmosphere) inhomogeneities,  from VLTI/AMBER, thereby helping  advance the still poorly  understood topic of RGB mass loss (especially in a context  in which dust condensation might appear not to play  a significant role; in general, it is RGB mass loss that  is puzzling, AGB mass loss that is straightforward)  ¶ Astron. Alm. (epoch 2021.5) assigns MK type  K5<sup>+</sup> III  ¶ although <a href="#">2019A&amp;A...625A..22R</a> casts doubt on  <a href="#">2018ApJ...865L..20F</a>, <a href="#">2015A&amp;A...580A.31H</a>  exoplanet assertion, exoplanet is asserted in NASA  exoplanet catalogue (as viewed 2021 Aug. 07)  slightly variable, 3.15–3.21 in V Tabit  further photometric study advisable?  (AAVSO(VSX) assessment in 2022:  status flag = suspected variable;  no variability classification symbol assigned)</p>
$\pi^3$	Ori A	4 51.1 +7 00	3.19	0.45	F6 V	124	3.7	26.3	0.464	89	+24 SB2	<p>slightly variable, 3.15–3.21 in V Tabit  further photometric study advisable?  (AAVSO(VSX) assessment in 2022:  status flag = suspected variable;  no variability classification symbol assigned)</p>
$\iota$	Aur	4 58.5 +33 12	2.68 <sup>v</sup>	1.53	K3 II	7	-3.2	500	0.016	155	+18V	<p>var?.: 2.63–2.78 in V?; poss. “+2P” (brown dwarfs?) Hassaleh  as of 2022 July 29, AAVSO(VSX) notes  a record of more than 85 AAVSO observations,  and notes existence of NSV entry  (AAVSO(VSX) assessment in 2022:  status flag = suspected variable;  no variability classification symbol assigned);  further photometric study advisable?  ¶ X-ray “hybrid star” (unusual combination of  (hot) corona, cool wind)  ¶ dimming by ISM dust, <math>\sim 0.6</math> mag.  slow ecl.: 2.92–3.83 in V, <math>\sim 27.1</math> y (dim <math>\sim 700</math> d) Almaaz  more formally, period has been  asserted as <math>9896.0 \pm 1.6</math> d (although AAVSO(VSX)  has 9892 d): as again discussed  twice below, there are spectroscopic,  as distinct from photometric, phenomena  indicative of an eclipsing mass before the  onset, and continuing after the end, of the  photometric eclipse;  AAVSO(VSX) assessment in 2022:  status flag = confirmed variable;  variability classification symbol = “EA/GS”  ¶ <math>m_v</math>, B–V values are for <math>\epsilon</math> Aur AB combined light  ¶ Astron. Alm. (epoch 2021.5) assigns MK type A9 Ia</p>
$\epsilon$	Aur A	5 03.7 +43 51	2.98 <sup>†</sup>	0.54 <sup>†</sup>	F0Iab? <sup>†</sup>	$\sim 2^{\dagger}$	-8.0:~1450 <sup>†</sup>	$\sim 0.003$	n.a.	-3	SB <sup>†</sup>	<p>slow ecl.: 2.92–3.83 in V, <math>\sim 27.1</math> y (dim <math>\sim 700</math> d) Almaaz  more formally, period has been  asserted as <math>9896.0 \pm 1.6</math> d (although AAVSO(VSX)  has 9892 d): as again discussed  twice below, there are spectroscopic,  as distinct from photometric, phenomena  indicative of an eclipsing mass before the  onset, and continuing after the end, of the  photometric eclipse;  AAVSO(VSX) assessment in 2022:  status flag = confirmed variable;  variability classification symbol = “EA/GS”  ¶ <math>m_v</math>, B–V values are for <math>\epsilon</math> Aur AB combined light  ¶ Astron. Alm. (epoch 2021.5) assigns MK type A9 Ia</p>



¶  $\epsilon$  Aur B MK type is ~B5 V

¶  $\epsilon$  Aur ranks among the longest-period eclipsing binaries (exceeding even V383 Sco, with period 13.5 y; WW Vul, with period 13.9 y; and VV Cep, with period 20.3 y: the current long-period record, however, is held by TYC 2505-672-1, at ~69.1 y [with dimming ~3.45 y])

¶ determination of orbit elements has proven troublesome, with [2012A&A...544A.91M](#) urging caution even in respect of recent careful studies

¶ it is remarkable that, even though the eclipsing entity is physically very extended (because the eclipse is protracted), and even though orbital dynamics indicates that the entity is quite massive, nevertheless no visible radiation from an eclipsing body is readily observable (i.e. it is remarkable that this SB is essentially a single-lined SB)

¶ although the (notably protracted)  $\epsilon$  Aur eclipse is largely flat-bottomed, nevertheless even during eclipse the (dimmed) spectrum of the primary can be seen, with no visible-wavelength colour preference in the attenuation (except that there are absorption lines, as from a semi-transparent atmosphere around the eclipsing mass, at the start and the end of the dimming); the [1937ApJ...86..570K](#) explanation, postulating a large semitransparent totally eclipsing mass, with the non-selective opacity due to scattering off free electrons, is now universally abandoned in favour of the [1953AJ.....58..219K](#) and [1965ApJ...141..976H](#) hypothesis of an almost edge-on ([2010ApJ...714..549H](#)) cool opaque gas-dust low-mass disk or disk-like entity (spiral arm? cf. [2013PASP..125..775G](#)) (rotating while orbiting, and several au in diameter, presenting a temperature gradient ~550 K to ~1150 K (representing the portions respectively farthest from and closest to the primary star), and in terms of its vertical development not a (thick) hockey puck but a (thin) wafer, of much larger radius than the primary star; [2013PASJ...65L...1S](#) gives evidence for clumping in the disk; [2015ApJ...220...14K](#) raises the possibility that the disk is slightly tilted out of the binary-system orbital plane, with consequent precession), shrouding a B-type star (B5V?) or star pair (the more dramatic hypothesis of a shrouded black hole is not now generally favoured: [2010AJ...140..595VW](#), e.g. reports null result from X-ray search), with the disk geometry making the eclipses of the primary star, as observed from Earth, only partial; the disk may have been formed by mass transfer from the primary star, and indeed [2013PASP..125..775G](#) and [2018MNRAS.479.2161G](#) report putative spectroscopic detection of narrow mass-transfer stream; the former paper stresses that the detection of rare-earth elements within the putative stream spectrum (an indication that the primary is highly evolved?) now poses a fresh puzzle, in a system traditionally classed as puzzling

¶ [2012ApJ...748L..28H](#) and [2012MNRAS.423.2075M](#) discuss the question of gas-to-dust ratio in the disk; [2015ApJ...798...11P](#), [2012MNRAS.423.2075M](#), and [2010ApJ...714..549H](#) suggest not-very-small values in the distribution of dust-particle

diameters, with the first two of these three papers suggesting carbonaceous chemistry; additionally, [2011AJ....142..174S](#) spectroscopy finds CO absorption bands, symptomatic of sublimation, with indications that large particles dominate ¶ [2013ARep....57..991P](#) and [2013PASP..125..775G](#) document indications that the structure of the disk does not greatly change from one eclipse to the next ¶ the brightening around mid-eclipse has in the post-1970 papers repeatedly been attributed to a central opening in the postulated disk: however, (a) dissenter [2011A&A...530A.146C](#) has instead suggested intrinsic variability in the primary (that indeed has various quasi-periods or periods, with 67 d and 123 d prominent, with also variations in radial velocity, and (unblended) spectral line width, and other periodic or quasi-periodic behaviour, including possible orbitally excited non-radial pulsation; there seems as yet, however, to be no extensive asteroseismology), and (b) dissenter [2011A&A...532L..12B](#) has instead suggested forward scattering by disk dust (a line of thought now supported by the key imaging-and-modelling paper [2015ApJS..220...14K](#)) ¶ *HIPPARCOS* yields  $\pi$  possibly  $< 2$  mas, distance  $\sim 2000$  ly; we now, however, choose to relinquish the *HIPPARCOS* determination, made at the limit of *HIPPARCOS* capabilities, in favour of [2019IBVS.6258....1P](#), that deduces from *Gaia* DR2  $\pi = 2.4144 \pm 0.5119$  mas, and goes on to deduce from this, via supplementary (not straightforward, Bailer-Jones et al. [2018AJ....156...58B](#)) considerations what we express here as “ $\sim 1450$  ly” ¶ section 1 of [2012A&A...546A.123G](#) and section 1 of [2012A&A...544A..91M](#) summarize past controversies regarding mass of primary (low or high?), stemming from the difficulty in determining distance ([2012A&A...546A.123G](#) assigns a high distance,  $\sim 4900$  ly, and consequently favours a high mass value,  $\sim 20 M_{\odot}$ ; however, several post-2009 papers instead assign a modest mass to the primary, suggesting various values within the range  $\sim 2 M_{\odot} - \sim 6 M_{\odot}$ : [2014MNRAS.445.2884M](#), e.g. suggests  $2.5 M_{\odot}$  for primary,  $5.4 M_{\odot}$  for secondary [and suggests disk diameter 8.9 au]: evolutionary status of the primary has been correspondingly controverted (post-AGB star, now of modest mass, with much past shedding of mass, and consequent accumulation of the low-mass opaque disk around the secondary (a view taken by various papers, including recently [2019IBVS.6258....1P](#)) or, rather, an evolutionally earlier supergiant (cf [2012JAVSO...40..647K](#)), even perhaps of high mass? – but it is clear that the primary is at any rate sufficiently evolved to have left the MS, and there are indications that it is pulsating and a wind source; angular diameter is 2.1 mas) ¶ most recent photometric eclipse started 2009 Aug. 12, ended 2011 Aug. 23  $\pm 15$  d; next secondary (shallow, for the casual observer elusive) eclipse is possibly 2025 Dec. 20 through 2028 Mar. 29; next (deep, easy observable) primary photometric eclipse starts in 2036; monitoring even outside both the primary eclipse and the secondary eclipse is useful,

												<p>in part because of intrinsic variations in the primary star (cf <a href="#">2012JAVSO.40.647K</a>); in part because the postulated dense disk has an extended "atmosphere" yielding (e.g.) H<math>\alpha</math> absorption even outside photometric eclipse (<a href="#">2011A&amp;A...530A.146C</a>), with spectral premonitions starting ~3 y before the onset of the photometric eclipse; and in part because the opaque primary-star-eclipsing disk is potentially liable to thermal changes, visible in mid-infrared outside primary and secondary eclipse (<a href="#">2011AJ...142..174S</a>)</p> <p>¶ the Kloppenborg et al. CHARA interferometric imaging of the eclipsing disk is perhaps the single largest 21st-century advance in <math>\epsilon</math> Aur studies: <a href="#">2010Natur.464.842G</a> supplies journalistic background, including a recapitulation of <a href="#">2010ApJ...714..549H</a> modelling; <a href="#">2010Natur.464.870K</a> is the formal Kloppenborg et al. discovery paper (with the first spatially resolved image for any eclipsing binary during eclipse); and <a href="#">2015ApJS...220...14K</a> is a Kloppenborg-et-al update, with additional interferometry, now including also PTI and NPOI (and supplying also an overall history of <math>\epsilon</math> Aur studies)</p> <p>¶ news sources include <a href="#">mysite.du.edu/~rstencel/epsaur.htm</a> (Prof. R. Stencel, Univ of Denver, on the Kloppenborg-2010 team) and <a href="#">twitter.com/epsilon_Aurigae</a>; <a href="#">2012JAVSO.40.618S</a> summarizes the 2009–2011 campaign from an AAVSO perspective; an 18-paper archive, of NSF-supported ~2009-through~2011 AAVSO eclipse campaign, is at <a href="#">www.aavso.org/citizen-sky-epsilon-aurigae-papers</a></p> <p>¶ since the eclipsing companion of <math>\epsilon</math> Aur A has not yet been resolved, even in interferometry, WDS is not as yet (at any rate as of 2021 Oct. 27) able to write "<math>\epsilon</math> Aur Aa," "<math>\epsilon</math> Aur Ab"; WDS does, on the other hand, catalogue celestial-sphere neighbours B,C,D,E,F,G,H,I,J,K; all of these are fainter than mag. 10, except for <math>\epsilon</math> Aur E (at the wide angular separation of 207" from <math>\epsilon</math> Aur A, and at mag. 9.6 just barely clearing our mag.-10 threshold for comment-worthiness of a bright-star neighbour on celestial sphere)</p>	
$\epsilon$	Lep	5 06.5	-22 20	3.17 <sup>†</sup>	1.47	K4 III	15	-0.9	210	0.076	164	+1	<p>slightly var.? (3.12–3.20 in V?) as of 2022 July 29, AAVSO(VSX) notes availability of more than 30 AAVSO observations, and notes existence of NSV entry (AAVSO(VSX) assessment in 2022: status flag = suspected variable; no variability classification symbol assigned); further photometric study advisable?</p>
$\eta$	Aur	5 08.2	+41 16	3.17	-0.18	B3 V <sup>†</sup>	13	-1.2	240	0.075	155	+7 V?	<p>¶ evolutionary status is uncertain: RGB or AGB? slight rotating ellips. var?: 3.16–3.18 in V, 2.5617 d Haedus AAVSO(VSX) finds photometric status as rotating no-eclipses ellipsoid to be possible-and-yet-not-certain (AAVSO(VSX) assessment in 2022: status flag = confirmed variable; variability classification symbol = "<b>ELL:</b>"); further photometric study advisable? ¶ it has been suggested that variations are present also in spectroscopy ¶ weak magnetic field detected, ~2<math>\times</math> strength of Earth's dipole field</p>
$\beta$	Eri A	5 09.0	-5 03	2.78 <sup>†</sup>	0.13	A3 IVn	36	0.6	89	0.112	228	-9	<p>slight var.? (2.72–2.80 in V?) as of 2022 July 29, AAVSO(VSX) notes availability of 59 AAVSO observations, and notes existence of NSV entry (AAVSO(VSX) assessment in 2022: status flag = suspected variable;</p>

$\mu$	Lep	5 14.0 -16 11	3.29 <sup>†</sup> -0.11	B9p IV: (HgMn) <sup>†</sup>	18	-0.5 190	0.050 109	+28	no variability classification symbol assigned; further photometric study advisable?); unexplained brightening episode, over 2 h, by ~3 mag, in 1985 (recalling the 1972 unexplained brightening of $\epsilon$ Peg) elusive var. 0.001 in V, period 3.0113 d ( <i>TESS</i> ) variable of $\alpha$ CVn A (“ $\alpha^2$ CVn”) type; <a href="https://en.wikipedia.org/wiki/Mu_Leporis">en.wikipedia.org/wiki/Mu_Leporis</a> has a light curve, from <i>TESS</i> data (AAVSO(VSX) updated assessment, as viewed 2023 Jan. 28: status flag = confirmed variable; variability classification symbol = “ACV”) ¶ among the brightest of the Hg-Mn stars ¶ Astron. Alm. (epoch 2021.5) assigns MK temperature type B9p HgMn (and does not assign an MK luminosity class) ¶ X-ray emission noted from putative companion, at angular distance 0.93”
$\beta$	Ori A <sup>†</sup>	5 15.7 -8 11	0.14v <sup>†</sup> -0.03	B8 Ia	4	-6.9 900	0.001 69	+21 SB	B: 6.8, B5 V, 9.7” (2017); C: 7.6; BC: 0.1” Rigel A-BC orbit $\geq 25,000$ y, BC orbit $\sim 400$ y ¶ $m_v$ , B-V values are for $\beta$ Ori ABC combined light ¶ the $\beta$ Ori system presents variability in the $\alpha$ Cyg class, range 0.08-0.20 in V AAVSO(VSX) assessment in 2022: status flag = confirmed variable; variability classification symbol = “ACYG” ¶ E(B-V) = +0.00
$\alpha$	Aur <u>Aa,Ab</u> <sup>†</sup>	5 18.4 +46 01	0.07v <sup>†</sup> 0.80	G6:III + G2:III	76	-0.5 43	0.433 170	+30 SB2	composite; Aa: 0.7, Ab: 0.9; Aa,Ab < 0.1” <b>Capella</b> ¶ $m_v$ , B-V values are for $\alpha$ Aur Aa,Ab combined light; Aa,Ab are resp. mags. $\sim 0.08$ , $\sim 0.18$ in V passband ¶ under IAU rules, “Capella” designates Aa, not Ab ¶ orbit of the Aa, Ab SB is 140.0 d, with orbit exactly or very nearly circular; Aa,Ab is the first binary with orbit studied interferometrically (Anderson-Pease, Mt Wilson, 1910: the binary is informally known as “The Interferometrist’s Friend”); full system, however, appears to be $\alpha$ Aur Aa+Ab+H+L, where H and L are red dwarfs, of respective mags. 9.99 and 13.5, sharing the proper motion of Aa+Ab and perhaps possessing further gravitationally bound companions (with $\alpha$ Aur B,C,D,E,F,G,I,J,K being mere line-of-sight coincidences; of these 9 celestial-sphere neighbours to “The Interferometrist’s Friend” which is the Aa,Ab SB, only G, at wide angular separation of 522” from Aa,Ab, is brighter than mag. 10 (at mag. 8.10); additionally, WDS documents the celestial-sphere neighbours M,N,O,P, of which only the sparsely observed M and N are brighter than mag. 10 (at mag. 6.29 and mag. 9.84, respectively); the crowding, surely with abundant line-of-sight coincidences, in this region of the celestial sphere is perhaps to be expected, given the celestial-sphere adjacency of $\alpha$ Aur to the Milky Way ¶ $\alpha$ Aur Ab is in rapid evolutionary transition, currently crossing the Hertzsprung Gap ¶ $\alpha$ Aur system presents possible variability, 0.03-0.16 in V; as of 2022 July 29, AAVSO(VSX) notes the availability of 95 AAVSO observations, and notes existence of NSV entry (AAVSO(VSX) assessment in 2022: status flag = suspected variable; no variability classification symbol assigned); elsewhere, RS CVn-type variability has been suggested; further photometric study advisable? ¶ $\alpha$ Aur system is among the brightest of X-ray sources ecl.: 3.31-3.60 in V, 8.0 d; A: 3.6; B: 4.9, 1.8” (2019) $\eta$ Ori Aa is unresolved SB2, strictly 7.989268 d, eclipsing ( $\beta$ Per type), to be informally thought of as “the Aa1, Aa2 binary” (but since the SB2 is not as yet resolved, even in interferometry, WDS is not as yet able to formally catalogue the stars as $\eta$ Ori Aa1, $\eta$ Ori Aa2, and uses the informal terminology only in a note); Ab experiences “Aa1,” “Aa2” as essentially a point mass; the Ab-with- “Aa1,Aa2” period is 9.44 y,
$\eta$	Ori <u>Aa</u> <sup>†</sup>	5 25.6 -2 23	3.36v <sup>†</sup> -0.17 <sup>†</sup>	B0.5 Ve <sup>†</sup>	3	-4.0 1000	$\sim 0.004?$	n.a.+20 SB2	

$\gamma$	Ori A	5 26.4	+6 22	1.64 <sup>†</sup>	-0.22	B2 III <sup>†</sup>	13	-2.8	250	0.015	212	+18 SB?	<p>spectroscopically SB but not SB2 (with at least one orbital solution published); in a very slow orbit with the three-star “Aa1,Aa2 binary”-plus-Ab system, and so in turn experiencing that triple as essentially a point mass, is <math>\eta</math> Ori B (mag. 4.9; AB astrometry is 0.9”→1.9”, PA 87°→77°, 1848→2020, with orbit <math>\geq</math> 2000 y)</p> <p>¶ one or other of “Aa1,” “Aa2” is a pulsator, with period 8 h (possible <math>\beta</math> Cep variability? but one or other of the Aa components is an instance of the “Be phenomenon,” and additionally a shell spectrum has been observed from one or other of “Aa1,” “Aa2”); as of 2022 July 29, AAVSO(VSX) assessment is as follows: status flag = confirmed variable; variability classification symbol = “EA+BCEP:”</p> <p>¶ <math>m_v</math>, B–V values are for <math>\eta</math> Ori Aa,Ab combined light</p> <p>¶ Astron. Alm. (epoch 2021.5) assigns MK type B1 IV</p> <p>¶ BSC5: “expanding circumstellar shell”</p> <p>possible slight variability, 1.59–1.64 in V possible variability in <math>\gamma</math> Ori system noted ~1988 (the system was previously believed constant, and was used in a defining of the UBV photometric scheme); as of 2022 Jul. 29, AAVSO(VSX) notes the availability of just one AAVSO observation, and notes the existence of NSV entry (AAVSO(VSX) assessment in 2022: status flag = suspected variable; no variability classification symbol assigned); further photometric study advisable?</p> <p>¶ BSC5: “expanding circumstellar shell”</p>	<b>Bellatrix</b>
$\beta$	Tau	5 27.8	+28 38	1.65	-0.13	B7 III <sup>†</sup>	24	-1.4	130	0.175	173	+9 V SB	<p>BSC5: “expanding circumstellar shell”</p> <p>¶ lunar occultations possible as far N as southern California: we use simply the designation “<math>\beta</math> Tau,” even though WDS has “<math>\beta</math> Tau Aa”; there is a single observation of an alleged “<math>\beta</math> Tau B,” of undocumented mag., at angular separation 33.40”, from 1898, and additionally a single observation of an alleged “<math>\beta</math> Tau Ab,” of mag. 2.40, at angular separation just 0.10”, possibly an occultation observation, from 1930; this sparse observational material suggests to us that a lunar-occultation campaign (for observers at suitable latitudes, such as in southern California, although not in Canada or in northern Europe) might be worth undertaking, perhaps to yield a null result rebutting the 1898 and 1930 suggestions of multiplicity, perhaps to yield a non-null result confirming at least one of these two suggestions of multiplicity</p> <p>¶ often, but not invariably, classified as Hg–Mn star: Mn 25<math>\times</math> solar (and Ca, Mg only ~0.12<math>\times</math> solar: radiative lofting, gravitational settling)</p> <p>¶ AAVSO situation in 2022: no status flag (so not confirmed variable, not suspected variable, not confirmed non-variable), a fortiori no variability symbol assigned</p> <p>¶ E(B–V)=0.00</p> <p>B: 7.5, 2.7”, PA 268°→10°, 1875→2017</p> <p>¶ although <math>\beta</math> Lep B has been held to be possibly variable, AAVSO(VSX), as viewed 2022 July 29, has no entry (no status flag, so not confirmed variable, not suspected variable, not confirmed non-variable; a fortiori no variability symbol assigned)</p> <p>¶ <math>m_v</math>, B–V values are for <math>\beta</math> Lep AB combined light</p> <p>¶ duplicity now suspected also in <math>\beta</math> Lep A, through 2002 adaptive-optics observation at Haleakala: separation 2.58”</p>	<b>Elnath</b>
$\beta$	Lep A <sup>†</sup>	5 29.3	-20 45	2.84 <sup>†</sup>	0.82 <sup>†</sup>	G5 II	-20.3	-0.6	160	0.086	183	-14 V?	<p>¶ although <math>\beta</math> Lep B has been held to be possibly variable, AAVSO(VSX), as viewed 2022 July 29, has no entry (no status flag, so not confirmed variable, not suspected variable, not confirmed non-variable; a fortiori no variability symbol assigned)</p> <p>¶ <math>m_v</math>, B–V values are for <math>\beta</math> Lep AB combined light</p> <p>¶ duplicity now suspected also in <math>\beta</math> Lep A, through 2002 adaptive-optics observation at Haleakala: separation 2.58”</p>	<b>Nihal</b>
$\delta$	Ori <u>Aa</u> <sup>†</sup>	5 33.1	-0 17	2.23 <sup>v†</sup>	-0.22 <sup>†</sup>	O9.5 II	5	-4.4	700	0.001	137	+16 SB	<p>a good marker of celestial equator</p> <p>¶ Aa is an Algol-type eclipsing binary, not as yet resolved (so WDS is not as yet, at any rate as of 2021 Oct. 30, able to write “<math>\delta</math> Ori Aa1,” “<math>\delta</math> Ori Aa2”) with period 5.73249 d,</p>	<b>Mintaka</b>

$\alpha$	Lep A	5 33.7	-17 48	2.58 <sup>†</sup>	0.21	F0 Ib <sup>†</sup>	1.5	-6.6	2000	0.004	72	+24	<p>yielding a combined-light variation reported at AAVSO(VSX) (for the entire Aa,Ab system? or just for the unresolved Aa SB?) over the mag. range 2.20–2.32 (in the V band); (AAVSO(VSX) assessment in 2022: status flag = confirmed variable; variability classification symbol = “EA”)</p> <p>¶ WDS assigns to the unresolved Aa SB the MK type O9.5 II, and to the single Ab star (mag. 3.76) the MK type (with luminosity class tentative) “O9.7 III.”; angular separation of the unresolved Aa pair and the single star Ab has increased over recent decades (0.2”→0.3”, 1978→2019); at least one orbital solution has been published for Aa,Ab; there is also a bright celestial-sphere neighbour, <math>\delta</math> Ori C, at mag. 6.8 (AC astrometry 50”→56”, PA 0°→4°, 1777→2017)</p> <p>¶ <math>m_v</math>, B–V values are for <math>\delta</math> Ori Aa,Ab combined light</p> <p>¶ yielded first detection of ISM (Hartmann, 1904, through non-moving Ca line in the SB)</p> <p>¶ E(B–V)=+0.07</p> <p>slight variability?, (2.56–2.62 in V?) as of 2022 Jul. 29, AAVSO(VSX) notes availability of a single AAVSO observation, and notes existence of NSV entry (AAVSO(VSX) assessment in 2022: status flag = suspected variable; no variability classification symbol assigned) (further photometric study advisable?)</p> <p>¶ evolutionary status uncertain (has helium fusion already started in core?); helium-fusion past yields now abundances N 5× solar, Na 2× solar</p>	Arneb
$\beta$	Dor	5 33.8	-62 28	3.52v <sup>†</sup>	0.66	F7–G2 Ib	3.2	-3.7	1000	0.013	4	+7 V	<p>Cepheid variable: 3.41–4.08 in V, 9.84 d period not quite constant; evolutionary status uncertain;</p> <p>AAVSO(VSX) assessment in 2022: status flag = confirmed variable; variability classification symbol = “DCEP”)</p> <p>¶ observed by <i>FUSE</i>, <i>XMM-Newton</i> missions [THIS STAR ONLY IN ONLINE VERSION OF TABLE]</p>	
$\lambda$	Ori A <sup>†</sup>	5 36.4	+9 57	3.39v? <sup>†</sup>	-0.18 <sup>†</sup>	O8 III <sup>†</sup>	3	-4.2~	1100	0.004	216	+34	<p>WDS, citing <a href="#">1985A&amp;AS...60..183L</a>, remarks that B may be a mere line-of-sight coincidence; WDS also gives, in addition to two faint celestial-sphere neighbours, neighbours <math>\lambda</math> Ori D (mag. 9.6; AD angular separation was 78” in 2012) and <math>\lambda</math> Ori F (mag. 9.2; AF angular separation was 151” in 2012)</p> <p>¶ <math>m_v</math>, B–V values are for <math>\lambda</math> Ori AB combined light</p> <p>¶ the most prominent member of the “<math>\lambda</math> Ori Cluster,” a.k.a. Collinder 69</p> <p>¶ within Sharpless 264 (i.e. Sh2-264), a.k.a. the “<math>\lambda</math> Ori Ring,” a.k.a. the “Angelfish Nebula,” a gas ring 150 ly in diameter (possibly, but not certainly, remnant from a Type II supernova)</p> <p>¶ <math>\lambda</math> Ori assemblage possibly harbours variability: as of 2022 July 29, AAVSO(VSX) notes the availability of 140 AAVSO observations, and notes the existence of NSV entry, and assigns possible range 3.38–3.54 in V passband; AAVSO(VSX) assessment is status flag = suspected variable, with no variability classification symbol assigned (further photometric study advisable?)</p> <p>¶ E(B–V)=+0.12</p>	Meissa
$\iota$	Ori Aa <sup>†</sup>	5 36.6	-5 54	2.77 <sup>†</sup>	-0.24 <sup>†</sup>	O9 III	~1.4	-6.5	2000	0.001	108	+22 SB2	<p>Aa,Ab 0.1” (2016), mags. 3.0, 6.3</p> <p>B: 7.3, B7 IIIp (He wk), 12.5”, PA 134°→146°, 1779→2018, orbit <math>\geq</math> 700,000 y;</p> <p>there is additionally a celestial-sphere neighbour <math>\iota</math> Ori C, at mag. 9.8, with AC angular separation 49” in 2002;</p> <p><math>\iota</math> Ori Aa is SB, not as yet resolved (so WDS is not as yet, at any rate as of 2022 Feb. 4, able to write “<math>\iota</math> Ori Aa1,” “<math>\iota</math> Ori Aa2”), with</p>	Hatysa

$\epsilon$	Ori A	5 37.4	-1 11	1.69v <sup>†</sup> -0.18	B0 Ia	2	-7.2 2000	0.002	118	+26 SB	<p>published orbital solutions (29.134 d, <math>e=0.76</math>, 0.11 au min, 0.8 au max): the elongated orbit, and a disparity in ages, suggest duplicity through many-body interaction-with-expulsion, rather than through the coGenesis that is usual for a binary; the <math>\iota</math> Ori Aa, <math>\iota</math> Ori Ab pairing does not for its part possess published orbital solutions</p> <p>¶ colliding winds make <math>\iota</math> Ori A a strong X-ray source</p> <p>¶ <math>m_v</math>, B-V values seem unavailable in GCPD; we take values from <a href="#">2002yCat.2237....0D</a> (but we are uncertain whether these values are meant for <math>\iota</math> Ori Aa,Ab combined light or for <math>\iota</math> Ori Aa,Ab,B combined light)</p> <p>¶ <math>\iota</math> Ori Aa,Ab system presents a slight variability, 2.76–2.79 in V, of an AAVSO(VSX) (but not GCVS) ellipsoid-photospheres type; additionally, and independently of this HB phenomenon, <math>\iota</math> Ori B is variable (AAVSO(VSX) assessment in 2022: status flag = confirmed variable; variability classification symbol = “<b>HB</b>”)</p> <p>¶ brightest member of Sword asterism</p> <p>¶ E(B-V)=+0.07</p> <p>var. of <math>\alpha</math> Cyg type, 1.64–1.74 in V supergiant, nonradially pulsating (AAVSO(VSX) assessment in 2022: status flag = confirmed variable; variability classification symbol = “<b>ACYG</b>”)</p> <p>¶ luminosity (etc) controverted: Crowther (2006) 275,000 <math>L_{\odot}</math>, Searle (2008) 537,00 <math>L_{\odot}</math>, Puebla (2015) 832,000 <math>L_{\odot}</math></p> <p>¶ E(B-V)=+0.08</p>	<b>Abnilam</b>
$\zeta$	Tau	5 39.1	+21 09	2.92v <sup>†</sup> -0.20	B2 IIIpe (shell) <sup>†</sup>	7	-2.7 400	0.020	175	+20 SB <sup>†</sup>	<p>var.?, ecl.?, (and <math>\gamma</math> Cas?), 2.80–3.17 in V, 133.0 d Tianguan although eclipse-generated variability is asserted at AAVSO(VSX), with period 132.9735 d (AAVSO(VSX) assessment in 2022: status flag = confirmed variable; variability classification symbol = “<b>E/GS+GCAS</b>”), the occurrence of eclipses has been disputed (cf <a href="https://en.wikipedia.org/wiki/Zeta_Tauri">en.wikipedia.org/wiki/Zeta_Tauri</a>); <math>\gamma</math> Cas variability would be consistent with the observed Be-phenomenon-cum-shell, and is accepted by AAVSO(VSX), although not accepted throughout the literature</p> <p>¶ <math>m_v</math>, B-V values are for the combined light of the <math>\zeta</math> Tau SB</p> <p>¶ the primary in the SB pairing is one of the best-known “Be phenomenon” stars, and is possibly one of the keys to the solution of currently unsolved Be-phenomenon problems; consistently with the shell-spectrum history, the disk is just 5° away from being seen edge-on (<a href="#">2013A&amp;ARv.2L...69R</a>, p. 58n); although the disk gases move in Keplerian orbits, their orbits are not circular, and consequently the material has some nonzero radial velocity even at the midpoint of transit; a further consequence of this kinematics is that the orbiting gas is less dense at apastron than at periastron; shell spectrum underwent three full cycles of V/R variation from 1997 to 2010, with these cycles generally taken as making the precession, under gravitational influence of the elusive SB companion, of a one-armed density wave within the Be disk (for geometry and time variations of disk, cf Fig. 7 of <a href="#">2010AJ....140.1838S</a>, Fig. 8 of <a href="#">2015A&amp;A...576A.112E</a>); however, in more recent years, the V/R cycling has been absent; precession notwithstanding, the disk has been observed to be stable, and therefore must be being fed by decretion from the host-star photosphere at a nearly constant rate; as a step toward the eventual discovery of the excitation structure of some conveniently observable Be-phenomenon disk, <a href="#">2012ApJ...744...19K</a> reports spectro-interferometry from two different <math>\zeta</math> Tau primary-star radii, in hydrogen Brackett <math>\gamma</math> and in a set of hydrogen Pfund lines (while drawing also on hydrogen Balmer <math>\alpha</math> data from previous literature); the emission is found to originate</p>	

$\alpha$	Col A	5 40.5	-34 04	2.65 <sup>†</sup>	-0.12	B7 IVe <sup>†</sup>	12	-1.9	260	0.025	176	+35 V?	<p>at roughly the same disk radius for hydrogen Balmer <math>\alpha</math> and hydrogen Brackett <math>\gamma</math>, and at a smaller radius for the hydrogen Pfund lines; the <a href="#">2012ApJ...744...19K</a> <math>\zeta</math> Tau study provides some observational support for the viscous decretion-disk, Keplerian-rotation model prevalent in recent Be-phenomenon theorizing, and additionally supports the density-wave-in-disk hypothesis for the observed V/R cycles; modelling efforts are ongoing, with <a href="#">2015A&amp;A...576A.112E</a> serving as a progress report</p> <p>¶ its rapid rotation and Be-phenomenon and shell-spectrum histories notwithstanding, the <math>\zeta</math> Tau primary has already evolved some distance off the MS, to “giant” stage (in general, giants are not expected to be rapid rotators); <a href="#">2012ApJ...744...19K</a> assumes an equatorial radius of <math>7.7 R_{\odot}</math></p> <p>¶ the nature of the elusive low-flux? <math>\sim 1 M_{\odot}</math> SB companion, of period 133.0 d, is unknown (could even be a neutron star); separation (with orbit nearly circular) is <math>\sim 1.17</math> au; since interferometry seems so far to have failed to resolve the companion, WDS, at any rate as of 2022 Feb. 4, is constrained to write “<math>\zeta</math> Tau” rather than “<math>\zeta</math> Tau A” and “<math>\zeta</math> Tau B”; the elusive companion may be producing a truncation in the Be-phenomenon disk, in the sense of a radical change in the dependence of disk density on radius (<a href="#">2013A&amp;ARv.21...69R</a>)</p> <p>¶ under IAU rules, the name “Tianguan” applies only to the primary, not to the entire SB system</p> <p>slight var.? (<math>\gamma</math> Cas type?) 2.62–2.66 in V? Phact</p> <p>AAVSO(VSX) as viewed 2022 July 29 indicates possible slight variability, finds no record of AAVSO observations, notes existence of NSV entry, indicates <math>\gamma</math> Cas-type variability as possible-but-not-certain, offers a V-passband range as a possibility (AAVSO(VSX) assessment in 2022: status flag = suspected variable; variability classification symbol = “GCAS:”) (further photometric study advisable?)</p> <p>¶ rapid rotator, with mass loss to disk, and so an instance of the “Be phenomenon”; H<math>\alpha</math> is variable, and H<math>\beta</math> profile varies rapidly; nevertheless, the Be disk is stable (unlike, e.g. the Be disk of <math>\gamma</math> Cas), indicating that the process of decretion-from-photosphere is in this case proceeding at a constant rate</p> <p>¶ E(B–V) = 0.00</p>
$\zeta$	Ori <u>Aa</u>	5 41.9	-1 56	1.76 <sup>†</sup>	-0.21 <sup>†</sup>	O9.5 Ib	4	-5.0	960	0.005	58+18	SB2 <sup>†</sup>	<p>B: 3.7, B0 III, 2.4”, PA 152°→167°, 1822→2017 Alnitak orbit <math>\geq 1500</math> y; B is a very rapid rotator, and also is possibly a <math>\beta</math> Cep variable, with departures from the classic rotational-broadening spectral profile (<a href="#">2013A&amp;A...554A..52H</a> is first presentation of high-quality spectroscopy for <math>\zeta</math> Ori B; for the overall <math>\zeta</math> Ori system, AAVSO(VSX) as of 2022 Jul. 29 notes the availability of 1146 AAVSO observations, and gives the V-passband range 1.74–1.77 (AAVSO(VSX) assessment: confirmed variable; variability classification symbol = “VAR”) (further photometric study advisable?)</p> <p>¶ there is additionally a celestial-sphere neighbour <math>\zeta</math> Ori C, of mag. 9.6 (AC astrometry: 56”→58”, PA 8°→10°, 1839→2017), said by WDS to be possibly a physical companion; <math>\zeta</math> Ori Aa,Ab now possesses an orbital solution from a combination of high-resolution spectroscopy (Aa,Ab is now found to be SB2, not merely SB) with full-period NPOI interferometry campaign (<a href="#">2013A&amp;A...554A..52H</a>; period is 7.3 y); since minimum distance of the O-star <math>\zeta</math> Ori Aa from the B1 IV star <math>\zeta</math> Ori Ab is 9.5 au, this system is a good candidate for mass determinations (helpfully for the determinations, the distance,</p>



ζ	Lep	5 48.0	-14 49	3.54	0.10	A2 Vann <sup>†</sup>	~46.3	1.9	~70.5	0.015	266	+20SB?	<p>even at periastron, precludes a significant mass-transfer stream linking Aa with Ab; and in fact ζ Ori Aa is not only the brightest of the O-type stars in our visual sky, but is the first O-giant to have been assigned a mass via orbital computation); although there is no mass transfer between ζ Ori Aa and ζ Ori Ab, nevertheless Aa does eject mass vigorously, consistently with its membership in MK type O; at least one orbital solution has also been published for Aa,Ab-with-B (where ζ Ori B experiences the Aa,Ab binary as essentially a point mass)</p> <p>¶ m<sub>v</sub>, B-V values are for ζ Ori Aa,Ab,B combined light</p> <p>¶ we here take the distance D from the Solar System (in our notation, and writing here to just 2 significant figures) 960 ly suggested in <a href="#">2013A&amp;A...554A..52H</a> on the basis of orbital solution (the suggestion is made with 7% uncertainty, and additionally with the caveat that ζ Ori B photometry would indicate a larger D, in our notation ~1300 ly); from this D we deduce the corresponding value of π, as 4 mas; <i>HIPPARCOS 2007</i> stated instead a different D (in our notation, and to just one significant figure, D = 700 ly)</p> <p>¶ <a href="#">2013A&amp;A...554A..52H</a> suggests age of ~7 My (but elsewhere a still lower age, below 4 My, has been suggested)</p> <p>¶ E(B-V) = +0.09</p>
													<p>rapid rotator (period ~0.2 d or ~0.3 d)</p> <p>¶ has debris disk, has first known extrasolar asteroid belt</p> <p>¶ Astron. Alm. (epoch 2021.5) assigns MK type A2 Van</p> <p>¶ approached to within ~4 ly or ~5 ly of Sun ~1 My ago</p> <p>¶ AAVSO(VSX) situation in 2022: no status flag (so not a confirmed variable, not a suspected variable, not a confirmed non-variable; a fortiori no variability classification symbol)</p>
κ	Ori	5 48.9	-9 40	2.06 <sup>†</sup>	-0.18	B0.5 Ia <sup>†</sup>	5	-4.4	600	0.002	131	+21 V?	<p>[THIS STAR ONLY IN ONLINE VERSION OF TABLE]</p> <p>Saiph</p> <p>slight var. of α Cyg type, 2.04–2.09 in V (AAVSO(VSX) assessment in 2022: status flag = confirmed variable; variability classification symbol = “ACYG”)</p> <p>¶ evolutionary status unclear, high mass-loss rate</p> <p>¶ carbon-deficient (with metallicity otherwise unremarkable)</p> <p>¶ E(B-V) = +0.07</p>
β	Col	5 51.8	-35 46	3.11	1.16	K1.5 III <sup>†</sup>	37.4	1.0	87	0.408 <sup>†</sup>	8	+89 <sup>†</sup> V	<p>Wazn</p> <p>although high space velocity indicates that β Col is an interloper from outside galactic thin disk, nevertheless this star is richer than Sun in the elements beyond He</p> <p>¶ AAVSO(VSX) situation in 2022: no status flag (so not a confirmed variable, not a suspected variable, not a confirmed non-variable); a fortiori no variability classification symbol</p>
α	Ori Aa <sup>†</sup>	5 56.4	+7 25	0.48v <sup>†</sup>	1.86	M2 Iab <sup>†</sup>	7 <sup>†</sup>	-5.5	500 <sup>†</sup>	0.030 <sup>†</sup>	68	+21 <sup>†</sup> SB	<p>semireg., late-type supergiant var.: ~0–1.7 in V <b>Betelgeuse</b></p> <p>variability was discovered by J. Herschel in 1839; the latest minimum, early in 2020, at ~1.7 in V, sank below even the minima of 1927 and 1941 (each ~1.2); journalism on this 2020 event includes <a href="#">www.sciencenews.org/article/betelgeuse-star-dim-supernova-death-what-happened</a>; three currently offered explanations are dust cloud from mass ejection, (gigantic) starspot, and fortuitous coincidence of minima from three separate cyclical variations; recovery began 2020 Feb. 22, with a rise to ~0.3 in V by 2020 late April; AAVSO reports mag. 0.65 in V on 2021 Jan. 11, 0.60 in V on 2022 Feb. 4;</p> <p><a href="#">2018A&amp;A...615A.116M</a> suggests on basis of magnetic variations a scenario on which evolution of giant</p>

photospheric convective cells, generating magnetism through local dynamos, is responsible for the observed long secondary ~2100-day photometric period; there are additionally ~200- ~400-day photometric periodicities, plus a stochastic variation ascribed to photospheric granulation (AAVSO(VSX) assessment in 2022: status flag = confirmed variable; variability classification symbol = “**SRC**”)

¶  $m_v$ , B–V values are for  $\alpha$  Ori Aa,Ab combined light

¶ Astron. Alm. (epoch 2021.5) assigns MK type M1–M2 Ia–Iab

¶ brightest star in IR sky, also brightest in bolometric sky

¶ nearest RSG (contrast with  $\sigma$  Cet, as AGB); greatest angular diameter of almost any star other than Sun (near-IR limb-darkened disk ~42 mas; but R Dor, having approx 1/3 radius of  $\alpha$  Ori, is less distant, and so attains still greater angular diameter);

[en.wikipedia.org/wiki/List\\_of\\_largest\\_stars](https://en.wikipedia.org/wiki/List_of_largest_stars) supplies context, giving radii for many supergiants; reduction of  $\alpha$  Ori angular diameter over period 1993/2009 has been asserted

¶ [2017AJ...154...11H](#) reviews the longstanding  $\alpha$  Ori distance problem: parallaxes, including *HIPPARCOS*, labour under the difficulty of accurately determining photocentre of visually extended object, awkwardly harbouring even plumes and hotspots; we now give in our table these authors’ values for  $\pi$  (rounding from their 4.51 mas) and by implication for D (strictly 717 ly  $\pm$  20%)

¶ very slow rotator (true period difficult; 8.4 y has been suggested)

¶ [2019A&A...628A.101H](#) announces dust halo with inner radius 1.5 R\*;

[2016A&A...585A..28K](#) locates 3 R\* as the interface between hot-gas and more outlying dust envelopes

¶ CO shells inner 50 R\* to 150 R\*, outer as far as 250 R\*

¶ runaway star, exceeding local speed-of-sound in ISM: bow shock 6’–7’, from stellar wind meeting ISM, plus linear bar at 9’ (it has been suggested that the bar is a relic of collapsing wind from a previous BSG phase, and it also has been suggested that the bar is a feature intrinsic to the embedding ISM, unconnected with any  $\alpha$  Ori Aa wind)

¶ although RSG stars pose a more serious mass-loss problem for astrophysics than do the AGB stars, since it is not immediately clear what mechanism is lifting RSG stellar material above the photospheres (convection? pulsation? magnetism?), there is now a possible partial resolution in this particular case:

[2018A&A...609A..67K](#), using ALMA, finds  $\alpha$  Ori anisotropic mass loss, with plume of ejecta; the authors suggest that plume is associated with strong “rogue” convection cell, observable as photospheric hot spot (in contrast with the cool spots encountered on such MS stars as the Sun)

¶ progenitor mass possibly ~20  $M_{\odot}$  (making  $\alpha$  Ori very massive), age since arrival on ZAMS possibly 8.0–8.5 My (making  $\alpha$  Ori very young)

¶ present evolutionary status of  $\alpha$  Ori uncertain: has this RSG previously been a BSG? (and [2017MNRAS.465.2654W](#) suggests history may have been complicated by a stellar merger)

¶  $\alpha$  Ori is SN Type II-P progenitor, the core collapse being due within, (perhaps much within) 1 My: although SN will plateau for several months, yielding a star visible even in daytime, with the brilliance of a quarter Moon or full Moon,

$\beta$	Aur <u>Aa,Ab</u>	6 01.3	+44 57	1.90 <sup>†</sup>	0.03 <sup>†</sup>	A1 IV + A1 IV	~40.2	-0.1	81	0.056	269	-18 SB2	<p>the SN radiation from so distant a source will not constitute a terrestrial biohazard</p> <p>¶ <i>Sky &amp; Telescope</i>. feature article 2019-05 on <math>\alpha</math> Ori can usefully be supplemented with Fig. 13 from <a href="#">2018A&amp;A...609A..67K</a> (multi-wavelength composite, showing ejecta plume condensing to dust at a few <math>R^*</math>, and showing also two areas of local photospheric magnetic activity): AAVSO has backgrounder at <a href="http://www.aavso.org/vsots_alphaori">www.aavso.org/vsots_alphaori</a></p> <p>¶ WDS documents the putative detection, from a small amount of work in speckle interferometry, of two close companions (and is therefore compelled to write “<math>\alpha</math> Ori Aa,” “<math>\alpha</math> Ori Ab,” and “<math>\alpha</math> Ori Ac”); since we reproduce WDS designations, we are in turn obliged to refer to Betelgeuse not as <math>\alpha</math> Ori A but as <math>\alpha</math> Ori Aa); however, since the WDS-documented speckle interferometry observations are from no later than 1983, and since current interferometry detects no close companions, it is now likely that Betelgeuse is unperturbed by any other star (the very faint WDS-catalogued stars <math>\alpha</math> Ori B,C,D, E,F,G,H,I,J all lie on the celestial sphere at large angular separations from Betelgeuse, with the smallest angular separation, between A and B, measured as a quite wide 38” in 2014; of these faint celestial-sphere neighbours, even the brightest, <math>\alpha</math> Ori E, shines at a mere mag. 11)</p> <p>slight ecl.: 1.89–1.98 in V, 3.96 d (mags. equal) Menkalinan eclipsing system is of <math>\beta</math> Per type (AAVSO(VSX) assessment in 2022: status flag = confirmed variable; variability classification symbol = “<b>EA/DM</b>”)</p> <p>¶ orbit is found in the published orbital solutions to be either circular or nearly circular (possibly <math>e=2.8e-06</math>); <math>\beta</math> Aur Aa,Ab was spectroscopically identified as a binary in 1890 (and is said to be only the third binary ever to be spectroscopically identified); orbit has been studied interferometrically since 1990s</p> <p>¶ <math>m_v</math>, B–V values are for <math>\beta</math> Aur Aa,Ab combined light</p> <p>¶ under IAU naming rules, “Menkalinan” denotes Aa, not Ab</p>
$\theta$	Aur <u>A</u>	6 01.3	+37 13	2.64 <sup>†</sup>	-0.08 <sup>†</sup>	A0p II: (Si) <sup>†</sup>	~19.7	-0.9	166	-0.086	~149	+30 SB	<p>B: 7.2, G2 V, 4.2”, PA <math>7^\circ \rightarrow 304^\circ</math>, 1871→2019 Mahasim orbit <math>\geq 1200</math> y, with AB distance <math>\geq 185</math> au</p> <p>¶ at least one of the components in the SB that is <math>\theta</math> Aur A is magnetic, and an oblique rotator; there are abundant anomalies in photosphere patches, with Si and Cr 10<math>\times</math> and 100<math>\times</math> solar, respectively; consistently with rotation, the <math>\theta</math> Aur A SB pairing presents weak variability, with AAVSO(VSX) assigning range 2.63–2.66 in V, period 3.618664 d (and yet a period of 1.37 d has been asserted elsewhere) (AAVSO(VSX) assessment in 2022: status flag = confirmed variable; variability classification symbol = “<b>ACV</b>”); although AAVSO(VSX) asserts “<math>\alpha^2</math> CVn-” (<math>\alpha</math> CVn A-) type variability, <a href="#">2007A&amp;A...464.1089S</a> finds that observed variations in H<math>\alpha</math>, H<math>\beta</math>, and H<math>\gamma</math> profiles cannot be modelled with photosphere inhomogeneities (this is the <math>\alpha</math> CVn A (“<math>\alpha^2</math> CVn”) variability scenario), and instead proposes changes in atmospheric pressure structure, as ions moving in the star’s magnetic field undergo Lorenz-force deflections</p> <p>¶ <math>m_v</math>, B–V values are for <math>\theta</math> Aur AB combined light var.: 3.1–3.7 in V, 2979 d; B: 6.2, 1.8” (2020) Propus (AB astrometry in detail: 1.1”→1.8”, PA <math>300^\circ \rightarrow 258^\circ</math>, 1881→2020); a potentially confusing blend of two variabilities: the unresolved single-line SB which is <math>\eta</math> Gem A is an Algol-type eclipsing system, with each eclipse lasting 17 weeks, system range possibly 3.1–3.8 in V, period 2969 d</p>
$\eta$	Gem <u>A</u> <sup>†</sup>	6 16.3	+22 30	3.28v <sup>†</sup>	1.59 <sup>†</sup>	M3 III	8	-2.0	400	-0.064	~259	+19 SB	

											(most recent minimum around 2020 Oct. 22, with mag 3.766 in V band reported at AAVSO; next eclipse may therefore be expected to begin late in 2028); additionally, however, one or the other component of this SB presents semiregular instability of a type more or less analogous to o Cet, with one or more periods, average period 234 d (AAVSO(VSX) assessment in 2022: status flag = confirmed variable; variability classification symbol = "EA/GS+SRA") ¶ m <sub>v</sub> , B-V values are for η Gem AB combined light ¶ liable to lunar, and also to very rare planetary, occultations (making η Gem A not only an SB, but also an occultation binary)	
ζ	CMa Aa,Ab	6 21.2 -30 04	3.02 <sup>†</sup> -0.20 <sup>†</sup>	B2.5 V	9.0	-2.2 360	0.008	61	+32	SB <sup>†</sup>	Furud	
											SB is recently resolved, as ζ CMa Aa, ζ CMa Ab, with just 4 observations (from 2019 and 2020) documented in WDS as of 2021 Nov. 4 (Aa,Ab angular separation surely much less than 1"); SB period is 675 d; WDS asserts mags. 3.6, 3.8; the pairing with ζ CMa B (mag. 7.8) is wide (167"→170", PA 338°→340°, 1833→2016) ¶ m <sub>v</sub> , B-V values are for ζ CMa Aa,Ab combined light ¶ variability has been claimed somewhere in what is now resolved as the ζ CMa Aa,Ab pair (possibly of the β Cep type: as of 2022 July 29, AAVSO(VSX) notes the existence of an NSV entry, but finds no record of AAVSO observations, and is able to state V-passband range only as "3.02-?" (AAVSO(VSX) assessment in 2022: status flag = suspected variable; variability classification symbol = "BCEP:") (further photometric study advisable?)	
β	CMa A	6 23.7 -17 58	1.98 <sup>†</sup> -0.24	B1 II-III	7	-3.9 ~490	0.003	256	+34	SB	Mirzam	
											slight var., β Cep type, 1.97-2.00 in V, 0.25130 d (AAVSO(VSX) assessment in 2022: status flag = confirmed variable; variability classification symbol = "BCEP") (we give here the AAVSO(VSX) period and V-passband range, as viewed 2022 Jul. 08); the brightest of the β Cep pulsators; has multiple modes, with beat period 50 d; it is not known why ε CMa, while physically similar, is not a pulsator ¶ near the boundary of the "Local Bubble" ISM cavity ¶ E(B-V) = +0.01	
μ	Gem A	6 24.4 +22 30	2.87 <sup>v†</sup> 1.64	M3 IIIab	14	-1.4 230	0.124	153	+55	V?	Tejat	
											semiregular variable: 2.75-3.02 in V (AAVSO(VSX) assessment in 2022: status flag = confirmed variable; variability classification symbol = "SR") ¶ on AGB ¶ subject to lunar occultations	
α	Car	6 24.4 -52 43	-0.72 0.16 <sup>†</sup>	A9 Ib <sup>†</sup>	11	-5.5 ~310	0.031	41	+21		Canopus	
											visible both in X-ray (magnetically heated corona; also rapid rotator, strongly convective) and in radio ¶ evolutionary status not fully clear, and colour unusual in its luminosity class ¶ AAVSO(VSX) situation in 2022: no status flag (so not a confirmed variable, not a suspected variable, not a confirmed non-variable), and a fortiori no variability classification symbol ¶ Astron. Alm. (epoch 2021.5) assigns MK type A9 II	
v	Pup	6 38.5 -43 13	3.17 <sup>†</sup> -0.11	B8 IIIIn <sup>†</sup>	9	-2.1 370	0.004	186	+28	SB		
											slight var., 3.16-3.20 in V; instance of "Be phenom.?" AAVSO(VSX) asserts λ Eri-type variability; shell spectrum has been suggested, with "central quasi-emission peak" (cf <a href="#">1999A&amp;A...348.831R</a> ) (AAVSO(VSX) assessment in 2022: status flag = confirmed variable; variability classification symbol = "LERI") ¶ rapid rotator, with period < 1.7 d ¶ distance was ~27 ly 3.6 My ago	
γ	Gem <u>Aa</u>	6 39.1 +16 23	1.93 <sup>†</sup> 0.00 <sup>†</sup>	A1 IVs	30	-0.7 110	0.057	166	-13	SB <sup>†</sup>	Alhena	
											γ Gem Aa,Ab is SB system in highly elongated orbit, known historically as an occultation binary (the brightest ever to be observed in an asteroid occultation: 381 Myrrha, in 1991) and as SB, but also reported in 2014 as resolved with adaptive	

$\varepsilon$	Gem A	6 45.4 +25 06	2.99 <sup>†</sup>	1.40 <sup>†</sup>	G8 Ib	4	-4.0	800	0.014	204	+10 SB	optics at the USA military facility “Starfire Optical Range,” thereby facilitating study of component masses (2014A.....147..65D, Fig. 6: that author finds period 12.634 y, $e=0.89$ , in good agreement with period and eccentricity from other published orbital solutions for $\gamma$ Gem Aa,Ab); average Aa,Ab distance is $\sim 8.5$ au; Ab is mag. $\sim 7.5$ ¶ AAVSO(VSX) situation in 2022: no status flag (so not a confirmed variable, not a suspected variable, not a confirmed non-variable), and a fortiori no variability classification symbol ¶ $E(B-V) = +0.03$ slight irreg. var., 2.97–3.05 in V Mebsuta (AAVSO(VSX) assessment in 2022: status flag = confirmed variable; variability classification symbol = “LC”) ¶ unusually yellow in the general population of supergiants; $m_v$ , B–V values are for $\varepsilon$ Gem A alone (not for $\varepsilon$ Gem AB combined light) ¶ among the few supergiants liable to lunar and planetary occultations ¶ celestial-sphere neighbour $\varepsilon$ Gem B (112" $\rightarrow$ 110", PA 93° $\rightarrow$ 94°, 1825 $\rightarrow$ 2013) is mag. $\sim 9.6$
$\alpha$	CMa <u>A</u> <sup>†</sup>	6 46.2 -16 45	-1.45 <sup>†</sup>	0.00 <sup>†</sup>	A0mA1 Va <sup>†</sup>	$\sim 379$	1.5	8.6	$\sim 1.339$	$\sim 204$	-8 SB	Sirius B: 8.5, WDA; 11.2" (2020); orbit 50.1 y separation 8.2 au min (3"), 31.5 au max (11", in 2019); although $\alpha$ Cma AB, now among the more celebrated visual binaries, was first resolved in 1862 (Alvan G. Clark, using 18.5-inch refractor), binary was conjectured as early as 1834 (by Bessel, on the grounds of variability in proper motion); <a href="https://cdsarc.u-strasbg.fr/viz-bin/getCaFile_Redirect/?-plus=-%2b&amp;B/wds/./notes.dat">cdsarc.u-strasbg.fr/viz-bin/getCaFile_Redirect/?-plus=-%2b&amp;B/wds/./notes.dat</a> , as notes for “WDS 06451-1643,” has vivid and extended discussion of 19 <sup>th</sup> -century struggles with obtaining accurate angular-separation and PA measurements for the $\alpha$ Cma AB system; as of 2021 Nov. 6, WDS reports the existence of 2061 measurements, for the timespan 1862 $\rightarrow$ 2020 (in contrast with the more difficult, and less well observed, $\alpha$ CMi AB system, where the secondary is again a WD, and the primary is again an intrinsically luminous ( $\sim 7$ Suns; Sirius radiates with the power of $\sim 25$ Suns) and nearby and notably hot star, but where smaller angular separation and greater magnitude difference pose a greater challenge: as of 2021 Nov. 6, WDS reports the availability of just 99 $\alpha$ CMi AB measurements); $\alpha$ Cma B is the brightest WD in the visual sky (its nearest competitors being $\alpha^2$ Eri B, at mag. 9.5, and $\alpha$ CMi B, at mag. 10.8); <a href="http://www.atnf.csiro.au/outreach/education/senior/astrophysics/binary_types.html">www.atnf.csiro.au/outreach/education/senior/astrophysics/binary_types.html</a> compares and contrasts a visual-wavelengths image of the $\alpha$ Cma AB pair (with A much brighter than B) and a CHANDRA X-ray image of this same pair (with B much brighter than A; A is said to appear in the image largely through reflecting UV emitted by the much hotter B) ¶ $m_v$ , B–V values are for $\alpha$ Cma A (which, however, can differ only minimally from $\alpha$ Cma AB combined light, since the $\alpha$ Cma B contribution is minuscule) ¶ AAVSO(VSX) assessment in 2022: status flag = confirmed non-variable; variability classification symbol = “CST” ¶ IRAS detected IR excess, a signature of dust (rather unexpected in a binary) ¶ Fe abundance of $\alpha$ Cma is $\sim 2\times$ or $\sim 3\times$ solar ¶ $\alpha$ Cma B is unusually massive for a WD (1.02 $M_\odot$ ; Chandrasekhar Limit is, however, 1.4 $M_\odot$ ; spectral type of $\alpha$ Cma B is DA [= hydrogen-only]) ¶ $E(B-V) = -0.03$
$\xi$	Gem	6 46.6 +12 52	3.35 <sup>†</sup>	0.44	F5 IV	56	2.1	58.7	0.223	211	+25 V? <sup>†</sup>	previous assertion of variability is now discounted Alzirr AAVSO(VSX) asserts constant light ¶ possibly SB, with components of $\sim$ equal mass ¶ rapid rotator (but just barely over the internal-structure

$\alpha$	Pic	6 48.4	-61 58	3.26	0.22	A6 Vn kA6 <sup>†</sup>	-34	0.9	100	0.252	345	+21	<p>transition, or “F5 rotation break,” that causes some stars to rotate rapidly, others to experience braking through magnetics and winds</p> <p>¶ AAVSO(VSX) assessment in 2022: status flag = confirmed non-variable; variability classification symbol = “CST”</p> <p>¶ X-ray source (suggesting significant corona)</p> <p>rapid rotator; shell, with time-varying spectral absorption features</p> <p>¶ X-ray emission suggests a companion, otherwise undetected</p> <p>¶ AAVSO(VSX) assessment in 2022: no status flag (so not a confirmed variable, not a suspected variable, not a confirmed non-variable), and a fortiori no variability classification symbol</p>
$\tau$	Pup	6 50.5	-50 39	2.93	1.20	K1 III	18	-0.8	180	0.077	154	+36 SB <sup>†</sup>	<p>SB period 1066.0 d, separation ~3 au, orbit of low eccentricity; since the SB is not as yet resolved, even in interferometry, WDS is not as yet able to write “<math>\tau</math> Pup A” and “<math>\tau</math> Pup B”</p> <p>¶ AAVSO(VSX) assessment in 2022: no status flag (so not a confirmed variable, not a suspected variable, not a confirmed non-variable), and a fortiori no variability classification symbol</p>
$\kappa$	CMa	6 50.7	-32 32	3.88v <sup>†</sup>	-0.21	B1.5 IVne <sup>†</sup>	4.9	-3.0	700	0.010	293	+14	<p>var., <math>\gamma</math> Cas type, 3.40–3.97 in V (was at faint end of its range before 1963; AAVSO reports visually ~3.3 in 2021 January, ~3.6 in Mar. 2022); an instance of the “Be phenomenon” (AAVSO(VSX) assessment in 2022: status flag = confirmed variable; variability classification symbol = “GCAS”)</p>
$\varepsilon$	CMa A <sup>†</sup>	6 59.6	-29 00	1.50 <sup>†</sup>	-0.21 <sup>†</sup>	B2 II	8.0	-4.0	410	0.004	68	+27	<p>B: mag.7.5 (7.9”, PA 161°→162°, 1850→2008) <b>Adhara</b></p> <p>AB distance 900 au, period at least 7500 y</p> <p>¶ brightest known source of extreme UV (~75 nm) in Earth’s night sky; hydrogen Lyman <math>\alpha</math> (121.6 nm) observed by NASA <i>OAO-3</i></p> <p>¶ <math>m_v</math>, B–V values are for <math>\varepsilon</math> CMa AB combined light</p> <p>¶ AAVSO(VSX) situation in 2022: no status flag (so not confirmed variable, not suspected variable, not confirmed non-variable), and a fortiori no variability classification symbol</p>
$\sigma$	CMa A	7 02.6	-27 58	3.47v <sup>†</sup>	1.73 <sup>†</sup>	K7 Ib <sup>†</sup>	3	-4.2	1100	0.008	308	+22	<p>¶ E(B–V) =+0.02</p> <p>irregular var.: 3.41–3.51 in V Unurgunite</p> <p>(AAVSO(VSX) assessment in 2022: status flag = confirmed variable; variability classification symbol = “LC”)</p> <p>¶ authorities are in some disagreement on MK type (possibly M, rather than K)</p> <p>¶ <math>m_v</math>, B–V values seem unavailable in GCPD; we take these values from <a href="#">2002yCat.2237....0D</a></p>
$\alpha^2$	CMa	7 04.0	-23 52	3.02 <sup>†</sup>	-0.08 <sup>†</sup>	B3 Ia	1	-6.6	3000	0.004	329	+48 SB	<p>[THIS STAR ONLY IN ONLINE VERSION OF TABLE]</p> <p>slight var., <math>\alpha</math> Cyg type, 2.98–3.04 in Hp band, 24.44 d (AAVSO(VSX) assessment in 2022: status flag = confirmed variable; variability classification symbol = “ACYG”)</p> <p>¶ <math>m_v</math>, B–V values seem unavailable in GCPD; we take these values from <a href="#">2002yCat.2237....0D</a></p>
$\delta$	CMa	7 09.3	-26 26	1.84	0.68	F8 Ia <sup>†</sup>	2	-6.6	2000	0.005	317	+34 SB	<p>¶ E(B–V) =+0.03</p> <p>Wezen</p>
$L_2$	Pup A+1P? <sup>†</sup>	7 14.3	-44 41	4.73v <sup>†</sup>	1.54	M5 IIIe	16	0.4	210	0.342	18	+53 V?	<p>¶ as of 2022 July 29, AAVSO(VSX) notes slight variability suspected, of unassigned type, range 0.004 mag. in V passband, with just one AAVSO observation available (AAVSO(VSX) assessment in 2022: status flag = confirmed variable; variability classification symbol = “VAR”) (further photometric studies advisable?)</p> <p>¶ <math>m_v</math>, B–V values seem unavailable in GCPD; we take these values from <a href="#">2002yCat.2237....0D</a></p> <p>¶ slow rotator (possibly ~1 y); N 2<math>\times</math> solar, Na 6<math>\times</math> solar semireg. late-type var., 2.6–8.0 in V, 140.6 d HR2748 fainter since 1995, with typical 2-mag. 140-day V-band variation now between 6 and 8 (AAVSO(VSX) assessment in 2022: status flag = confirmed variable;</p>

$\pi$	Pup Aa	7 18.0 -37 08	2.71 <sup>†</sup>	1.62	K3 Ib	4	-4.3	800	0.012	303	+16	<p>variability classification symbol = “<b>SRB</b>”);  for May and June 2022, the AAVSO  principal database lacks  V-band photometry, but has visual estimates  in the range [7.3, 8.1]: further photometric  studies now advisable, from observers equipped  with electronic detector and Johnson V passband?  ¶ on AGB, shedding mass  ¶ exoplanet possible but not certain (could be  mere cloud of gas and dust)  B: 7.9, 66”, PA 214°→213°, 1826→2009  Aa,Ab system has received only 2 observations,  both in 1991 (0.7” and again 0.7”, PA 148°  and 152°), with Ab at mag. ~6.5  ¶ as of 2022 Jul. 29, AAVSO(VSX)  states range as “2.8–2.87”  in Hp passband, and finds no record of  AAVSO observations  (AAVSO(VSX) assessment in 2022:  status flag = confirmed variable;  variability classification symbol = “<b>SRD</b>:”);  further photometric study advisable?  B: 8.2, K3 V, 5.5”, PA 198°→229°, 1822→2018 Wasat  orbit 1200 y  ¶ m<sub>v</sub>, B–V values are for <math>\delta</math> Gem AB combined light  ¶ AAVSO situation in 2022:  no status flag (so not confirmed variable, not suspected  variable, not confirmed non-variable); and a fortiori  no variability classification symbol)  ¶ lunar occultations possible; planetary occultations  possible-yet-rare: since the SB which is <math>\delta</math> Gem A is not  as yet resolved (even interferometrically) with  astrometry, WDS is not as yet (at any rate as of  2022 Feb. 5) able to write “<math>\delta</math> Gem Aa,” “<math>\delta</math> Gem Ab”; SB  period is 6.129y; a companion (the secondary  component in the SB?) has, however, been  noted through occultation  ¶ in evolutionary transition, having completed  stable core-hydrogen fusion  ¶ Astron. Alm. (epoch 2021.5) assigns MK type F0 V<sup>+</sup>  <b>[THIS STAR ONLY IN ONLINE VERSION OF TABLE]</b>  B: 6.8, 178” (2010) is mere optical companion Aludra  ¶ variable in <math>\alpha</math> Cyg class of non-radial pulsators;  AAVSO(VSX), as viewed 2021 Jan. 28  and again 2022 July 09, gives  range 2.36–2.50 in V, period 4.70433 d  (AAVSO(VSX) assessment in 2022:  status flag = confirmed variable;  variability classification symbol = “<b>ACYG</b>”)  ¶ m<sub>v</sub>, B–V values seem unavailable in GCPD;  we take these values from <a href="#">2002yCat.2237...0D</a>  ¶ strong wind; ejected circumstellar mass  inferred from IR excess  ¶ E(B–V) = +0.02</p>
$\delta$	Gem A	7 21.5 +21 56	3.53	0.34	F0 IV <sup>†</sup>	54	2.2	60	0.018	237	+4 SB <sup>†</sup>	<p>¶ m<sub>v</sub>, B–V values are for <math>\delta</math> Gem AB combined light  ¶ AAVSO situation in 2022:  no status flag (so not confirmed variable, not suspected  variable, not confirmed non-variable); and a fortiori  no variability classification symbol)  ¶ lunar occultations possible; planetary occultations  possible-yet-rare: since the SB which is <math>\delta</math> Gem A is not  as yet resolved (even interferometrically) with  astrometry, WDS is not as yet (at any rate as of  2022 Feb. 5) able to write “<math>\delta</math> Gem Aa,” “<math>\delta</math> Gem Ab”; SB  period is 6.129y; a companion (the secondary  component in the SB?) has, however, been  noted through occultation  ¶ in evolutionary transition, having completed  stable core-hydrogen fusion  ¶ Astron. Alm. (epoch 2021.5) assigns MK type F0 V<sup>+</sup>  <b>[THIS STAR ONLY IN ONLINE VERSION OF TABLE]</b>  B: 6.8, 178” (2010) is mere optical companion Aludra  ¶ variable in <math>\alpha</math> Cyg class of non-radial pulsators;  AAVSO(VSX), as viewed 2021 Jan. 28  and again 2022 July 09, gives  range 2.36–2.50 in V, period 4.70433 d  (AAVSO(VSX) assessment in 2022:  status flag = confirmed variable;  variability classification symbol = “<b>ACYG</b>”)  ¶ m<sub>v</sub>, B–V values seem unavailable in GCPD;  we take these values from <a href="#">2002yCat.2237...0D</a>  ¶ strong wind; ejected circumstellar mass  inferred from IR excess  ¶ E(B–V) = +0.02</p>
$\eta$	CMa A	7 25.0 -29 21	2.45v <sup>†</sup> –0.08 <sup>†</sup>		B5 Ia	2	-6.5	2000	0.007	325	+41 V	<p>rapid rotator, possibly ~1 d, with modest variability  in the hydrogen Balmer emission; disk of ejected matter  has diameter ~4× diameter of <math>\beta</math> CMi itself  (BSC5: “rotationally unstable”);  an instance of the “Be phenomenon”;  although GCVS assertion of  <math>\gamma</math> Cas-type variability has not been corroborated,  <a href="#">2007ApJ...654..544S</a> reports, using <i>MOST</i>,  millimagnitude “slowly pulsating B-type” variability;  AAVSO(VSX) as viewed 2021 Jan. 16  gives V-mag. range 2.84–2.92, but as viewed  2022 March 03 and 2022 July 09  the narrower range 2.89–2.90;  (AAVSO(VSX) assessment in 2022:  status flag = confirmed variable;  variability classification symbol = “<b>SPBe</b>”);  in contrast with e.g. the Be-phenomenon star <math>\gamma</math> Cas A,  the Be disk is in this case considered very stable  (<a href="#">2013A&amp;ARv.2L...69R</a>), indicating constancy in the  process of decrection from the host-star photosphere;  <a href="#">2012ApJ...744...19K</a> reports confirmation of Keplerian  rotation in the Be disk (an important follow-on  to the discovery of Keplerian rotation in</p>
$\beta$	CMi A	7 28.4 +8 14	2.89 <sup>†</sup>	-0.10	B8 Ve <sup>†</sup>	~20.2	-0.6	~162	0.064	234	+22	<p>Gomeisa</p>





(past mass transfer may be a complicating factor)  
 ¶  $\alpha$  CMi A radiates with the power of  $\sim 7$  Suns  
 (and so is not dramatically unlike  $\alpha$  CMa A,  
 which radiates as  $\sim 25$  Suns)  
 ¶  $m_v$ , B–V values are for  $\alpha$  CMi A (which, however,  
 can differ only minimally from  
 $\alpha$  CMi AB combined light, since the  
 $\alpha$  CMi B contribution is minuscule)  
 ¶ asteroseismology of  $\alpha$  CMi A is somewhat uncertain  
 (*MOST* mission 2004 did not find pulsations, and yet  
*WIRE* mission 1999 and 2000 did); at the level of  
 tens of millimag., or coarser, the previous claim  
 of variability is now discounted  
 (AAVSO(VSX) assessment in 2022:  
 status flag = confirmed non-variable;  
 variability classification symbol = “**CST**”)  
 ¶ the WD  $\alpha$  CMi B is physically unlike the WD  
 $\alpha$  CMa B, attaining only  $\sim 0.2$  of the density of  
 $\alpha$  CMa B, and being of a rare spectral type DQZ6.5  
 (elusive in ground-based spectroscopy: but at long  
 last (a) helium-not-hydrogen character of spectrum  
 was noted at HST through filter photometry in a  
 set of bands running from 1600 Å to 7828 Å,  
 and (b) detailed spectroscopy was performed at  
*HST* through STIS camera, over the range  
 1800 Å to 10000 Å; in this spectral type,  
 H features are absent and C, Mg, and Fe features  
 are present)  
 ¶ from an astrometrist’s perspective, the  $\alpha$  CMi AB  
 system contrasts with the less difficult, and  
 consequently better measured,  
 $\alpha$  CMa AB system, in which the secondary is  
 again a WD, and the primary is again  
 an intrinsically luminous and  
 nearby and notably hot star, with the magnitude  
 difference less severe and the typical angular  
 separation greater: as of 2021 Nov. 6, WDS  
 reports the existence of just 99  $\alpha$  CMi AB  
 astrometry measurements, as against the much  
 larger tally of 2061  $\alpha$  CMa AB measurements;  
 Bond et al. write, in [2015IAUDS.187....1M](#),  
 “Charles Worley, double-star observer at the USNO,  
 asserted to two of us, more than two decades ago,  
 that he was the only living astronomer who had seen  
 Procyon B with his own eye”  
 ¶ WDS documents, for the  $\alpha$  CMi AB system, celestial-  
 sphere neighbours  $\alpha$  CMi C,D,E,F,G,H; of these,  
 only E and G are brighter than mag. 10, at mags.  
 9.2 and 8.8 respectively (at the large respective angular  
 separations, from  $\alpha$  CMi A, of 467” [2009] and 356”  
 [2012])

$\beta$  Gem A+1P<sup>†</sup> 7 46.8 +27 58 1.14<sup>†</sup> 1.00<sup>†</sup> K0 IIIb 97 1.1 33.8 0.628 266 +3 V

Pollux

the nearest of the giants; unusual in being a giant  
 known to harbour an exoplanet (and the brightest  
 known exoplanet host in Earth’s sky); as of 2015,  
 exoplanet is IAU-named “Thestias”  
 ¶ possible slight variability: as of 2022 Jul. 29,  
 AAVSO(VSX) notes the availability of 34  
 AAVSO observations, notes the existence of  
 an NSV entry, and suggests  
 the possible V-passband band range 1.14–1.15  
 (AAVSO(VSX) assessment in 2022:  
 status flag = suspected variable;  
 variability classification symbol = “**VAR:**”),  
 further photometric study advisable?  
 ¶  $m_v$ , B–V values are for  $\beta$  Gem AB combined light  
 ¶ subject to rare lunar occultations, for observers  
 S of Earth’s equator  
 ¶ Castor-Pollux comparison is a helpful test of  
 naked-eye night colour response

$\xi$  Pup A<sup>†</sup> 7 50.3 –24 55 3.33<sup>†</sup> 1.24<sup>†</sup> G6 Iab–Ib<sup>†</sup> 3 –4.5 1200 0.005 260 +3 SB<sup>†</sup>

Azmid

full system comprises an unresolved tight binary (an SB)  
 and widely separated  $\xi$  Pup B, with B experiencing the  
 unresolved SB which is  $\xi$  Pup A as essentially a  
 point mass (B is mag. 13; 4.6” $\rightarrow$ 5.1”,  
 PA 189° $\rightarrow$ 191°, 1899 $\rightarrow$ 1964; orbit  $\geq$  26,000 y?)  
 ¶ SB primary has high metallicity, with exact  
 evolutionary status uncertain

												¶ SB primary is near, but is a little too cool to lie within, the HR diagram IS ¶ $m_v$ , B–V values are for $\xi$ Pup AB combined light ¶ possible slight variability in the $\xi$ Pup assemblage: as of 2022 Jul. 29, AAVSO(VSX) has no record of AAVSO observations, but does indicate range 3.31–3.45 in V passband and period 31.15265 d (AAVSO(VSX) assessment: status flag = suspected variable; variability classification symbol = “SRD:”)
$\chi$	Car	7 57.4 –53 03	3.46 <sup>†</sup> –0.18	B3 IV(p?) <sup>†</sup>	7	–2.3	500	0.035	304	+19 V		Si II anomalous strength now discounted ¶ the assertion, in <a href="https://en.wikipedia.org/wiki/Chi_Carinae">en.wikipedia.org/wiki/Chi_Carinae</a> as viewed 2022 Jul. 29, that $\chi$ Car is free of variability seems to us at the Handbook to be based on an erroneous reading of <a href="https://doi.org/10.1093/mnras/410.190T">2011MNRAS.410.190T</a> ; we instead favour AAVSO(VSX), which as viewed 2022 Jul. 29 asserts slight $\beta$ Cep-type variability (although, admittedly, without available AAVSO observations), asserting range 0.015 in V, period 2.4 h (AAVSO(VSX) assessment has status flag = confirmed variable; variability classification symbol = “BCEP”) ¶ the MK luminosity class “IV” (phenomenologically “giant”) notwithstanding, $\chi$ Car is in astrophysical terms in the last part of its stable core-hydrogen-fusion phase; Astron. Alm. (epoch 2021.5) assigns MK temperature type B3p Si without assigning an MK luminosity class blue supergiant Naos ¶ rapid rotator (1.78 d), despite ~2300 km/s stellar wind (in which spiral structure was announced in 2017 by <i>BRITE</i> mission team), with mass loss rate > 1e-6 $M_{\odot}$ /y ¶ high space velocity (impelled by past nearby supernova? or, rather, impelled by multibody gravitational interactions in its stellar birth family?); possibly ejected from Trumpler 10 OB association ¶ distance has been controverted ¶ He, N overabundant ¶ has been suspected of being a variable of the $\alpha$ Cyg type; however, as of 2022 Jul. 30, AAVSO(VSX), while confident of variability, and while giving V-passband range 2.24–2.26, period 1.78 d, classifies this star as possibly (not certainly) spotted-and-rotating, and notes that no AAVSO observations are available (AAVSO(VSX) assessment in 2022: status flag = confirmed variable; variability classification symbol = “ROT:”); further photometric study advisable? ¶ E(B–V) = +0.04 var.: 2.68–2.87 in V, 3.38114 h prototype of the “ $\rho$ Pup stars” (but in the AAVSO taxonomy, a $\delta$ Sct-type variable: 2022 AAVSO(VSX) assessment has status flag = confirmed variable, variability classification symbol = “DSCT”); main period is ~3.3 h (0.15 mag.); photosphere temperature is notably low in the overall population of stars presenting $\delta$ Sct variability ¶ Astron. Alm. (epoch 2021.5) assigns MK type F5 (Ib–II)p ¶ IR excess (circumstellar ring, at separation 50 au?) eruptive WR var.: 1.81–1.87 in V; Aa is a.k.a. $\gamma^2$ Vel AAVSO(VSX) as of 2022 Jul. 30 reports availability of 64 AAVSO observations, gives period 78.53 d (AAVSO(VSX) assessment: status flag = confirmed variable;
$\zeta$	Pup	8 04.4 –40 04	2.25 <sup>†</sup> –0.27 <sup>†</sup>	O5 Iafn <sup>†</sup>	3.0 <sup>†</sup>	–5.4	1080 <sup>†</sup>	0.034 <sup>†</sup>	299	–24 <sup>†</sup> V?		
$\rho$	Pup A	8 08.5 –24 22	2.80v <sup>†</sup> 0.42	F2mF5 II: (var) <sup>†</sup>	51.3	1.4	64	0.095	299	+46 SB		Tureis
$\gamma$	Vel Aa <sup>†</sup>	8 10.3 –47 24	1.79 –0.24	O7.5 III-1 <sup>†</sup>	3 <sup>†</sup>	–5.9~1100 <sup>†</sup>		0.012	330+35	SB2 <sup>†</sup>		

variability classification symbol = “WR”)

¶  $\gamma$  Vel Aa is a double-lined SB not as yet resolved, even in interferometry (so WDS is not as yet able to write “ $\gamma$  Vel Aa1,” “ $\gamma$  Vel Aa2”), but with published orbital solutions (period 78.5 d,  $e=0.4$  or  $0.5$ ); faint Ab (mag. 13.4) is poorly known, with just one astrometry result, from 1997 (Aa,Ab angular separation  $4.7''$ ); distance between components of the unresolved  $\gamma$  Vel Aa SB is  $0.8(?)$  au min,  $1.6$  au max);  $\gamma$  Vel B, a.k.a.  $\gamma^1$  Vel, is itself a resolved SB pair (so in WDS  $\gamma$  Vel Ba,  $\gamma$  Vel Bb: period is 1.48 d); AB astrometry is  $43'' \rightarrow 41''$ , PA  $222^\circ \rightarrow 221^\circ$ ,  $1826 \rightarrow 2017$

¶ the (carbon-rich) WR component in the unresolved SB which is  $\gamma$  Vel Aa is of spectral type WC8, and is the nearest and visually brightest of all WR stars (presenting “a unique opportunity to spatially resolve a WR wind by means of interferometry” [[2007A&A...464..107M](#)]), and is an exceptionally massive WR ( $9.0 M_\odot$ ; but at birth,  $> 30 M_\odot$ ); this SB is the best studied of all O-WR binaries: in the SB pair it is the WR component, rather than the O component, that dominates spectrally (although we assign an O type, and Astron. Alm. (epoch 2021.5) rather similarly assigns the uncertainty-flagged MK type “O9 I;” since the V-passband light is overwhelmingly from the more massive ( $18.5 M_\odot$ ) O-type component), making the  $\gamma$  Vel Aa SB the “Spectral Gem of the Southern Skies,” and a notable sight within broader “Vela complex” (dominated by the Gum Nebula, within which lie the Vela SNR, the *IRAS* Vela shell, and the Vela pulsar; some literature, including [2011A&A...525A.154S](#), indeed proposes intersection between the Vela SNR and a  $\gamma$  Vel SWB, taking the *IRAS* Vela shell as marking the meeting of SNR and SWB)

¶  $m_v$ , B–V values are for  $\gamma$  Vel Aa,Ab combined light  
¶ like  $\eta$  Car (bright to mag.  $\sim 0$  for several years after 1837, but now too faint, and now too lacking in firm future-outburst prognoses, to qualify for the RASC Handbook “Brightest Stars” list), the unresolved  $\gamma$  Vel Aa SB is a colliding-wind pair ([2017MNRAS.468.2655L](#) Fig. 1 sketches the collision geometry), and in consequence is a UV and X-ray source (and in consequence may also possibly resemble  $\eta$  Car in being a  $\gamma$ -ray source

[cf [2017ApJ...847..40R](#); as of at any rate 2017, it seems that no other colliding-winds-binary stellar  $\gamma$ -ray sources are known]); it is the wind from the (WR) secondary that dominates, with mass-loss rate at least  $100\times$  greater than for the (O-type) primary; the WR wind may feature some clumping, but is to a good approximation spherically symmetric until it encounters the O-star wind; orbital motion of the two SB components around their centre of mass yields a spiral structure in the wind-collision area, particularly salient at periastron

¶ [2017ApJ...847..40R](#) summarizes recent observations of the  $\gamma$  Vel Aa SB, in radio and IR and optical, including interferometry, noting inter alia discrepancies in the available determinations of mass-loss rates from the WR star (a copious  $3e-6 M_\odot /y?$  or a still more copious  $8e-5 M_\odot /y?$ )

¶ notable among recent observational studies are [2017MNRAS.468.2655L](#) (VLTI/AMBER near-IR spectro-interferometry, with also 3-D hydrodynamic modelling) and [2012MNRAS.427..581R](#)

¶ likely destiny of  $\gamma$  Vel Aa (WR) secondary is as (exotic) stripped-core SN (same prognosis as for  $\eta$  Car; this

															<p>contrasts with <math>\alpha</math> Ori, which will for its part instead explode as a (not exotic) hydrogen-spectrum SN)</p> <p>¶ dust emission is absent (even though formation of circumstellar dust is common in stars that, like the <math>\gamma</math> Vel Aa (WR) secondary, undergo copious mass outflow)</p> <p>¶ distance ~1200 ly, in contrast with our ~1100 ly, has also been recently asserted, on basis of VLTI/AMBER</p> <p>¶ we take MK type for <math>\gamma</math> Vel Aa from <a href="#">1999A&amp;A...345..163D</a> (as what must be considered an emendation of our (slightly cooler) Garrison-approved MK type from earlier editions of this table; admittedly, MK determination of <math>\gamma</math> Vel Aa is still difficult, because the raw spectrum is a composite comprising not only the O and WR stars, but also emission from the wind-collision zone)</p> <p>¶ Ba,Bb appear in combined light as mag. 4.1; there are additionally wide celestial-sphere neighbours C (mag. 7.3; AC separation was 62" in 2009) and D (mag. 9.2; AD separation was 94" in 2000)</p> <p>¶ neither the traditional Suhail al-Muhlif nor the modern Regor (devised within NASA, to commemorate 1967 fire victim Roger Chaffee) is presently IAU-approved name for any of the five stars <math>\gamma</math> Vel Aa primary, <math>\gamma</math> Vel Aa secondary, Ab, Ba, Bb</p>
$\beta$	Cnc A+1P	8 17.8	+9 07	3.52 <sup>†</sup>	1.48	K4 III <sup>†</sup>	11	-1.3	300	0.068	224	+22 V?			<p>Tarf</p> <p>"barium star," with Ba abundance ~6<math>\times</math> solar, presumably as contamination from defunct companion (but no companion remnant has been found)</p> <p>¶ GCPD offers not only the m, value of 3.52, but also, as an alternative possibility supported by a smaller number of authorities, 3.54</p> <p>¶ AAVSO(VSX) assessment in 2022: status flag = confirmed variable; variability classification symbol = "VAR")</p> <p>¶ Astron. Alm. (epoch 2021.5) assigns MK type K4 III Ba 0.5</p>
$\varepsilon$	Car A	8 23.0	-59 35	1.86v <sup>†</sup>	1.27	K3:III	5	-4.5	600	0.034	311	+2			<p><b>[THIS STAR ONLY IN ONLINE VERSION OF TABLE]</b></p> <p>B is possibly ecl., with AB mag. 1.82–1.94 in V <b>Avior</b> (AAVSO(VSX) as of 2022 Jul. 08 notes existence of NSV entry, has no record of AAVSO observations (AAVSO(VSX) assessment: status flag = suspected variable; variability classification symbol = "E:"); further photometric study advisable? – cf also <a href="#">2004AJ...127.2915P</a>, Table 5, which treats B as an unresolved binary (or is it that the eclipses (if real at all), involve not an unresolved "<math>\varepsilon</math> Car B primary," "<math>\varepsilon</math> Car B secondary" pair, but more straightforwardly the <math>\varepsilon</math> Car A, <math>\varepsilon</math> Car B pair?); full WDS-catalogued system is <math>\varepsilon</math> Car A (mag. 2.2) and <math>\varepsilon</math> Car B (mag. 3.9), with <math>\varepsilon</math> Car B not further resolved, and with the <math>\varepsilon</math> Car AB pairing reported by WDS as very tight, the angular separation being just 0.4" in 2019; the AB system is sparsely observed (WDS documents just 6 satisfactory astrometry measurements, 1991→2019); AAVSO general database (as distinct from AAVSO(VSX)) seems to have no record of observations, the <a href="#">1959AJ...64..127G</a> report of variability notwithstanding; Kaler comments at <a href="#">stars.astro.illinois.edu/sow/avior.html</a>, with reference to the overall paucity of observations, that "if there is a stellar category of 'bright stars getting no respect', [<math>\varepsilon</math> Car A] probably holds the record"</p> <p>¶ B is of MK type (uncertainty-flagged) "B2: V"</p> <p>¶ the IAU-official name "Avior" for <math>\varepsilon</math> Car A is of uncertain etymology, and yet its origins are known: here, as also with <math>\alpha</math> Pav A (IAU-officially "Peacock"), the name stems from the 1930s RAF Air Almanac project, which directed HM Nautical</p>

o	UMa A+IP 8 32.2 +60 38	3.37 <sup>†</sup>	0.85	G5 III	~18.2	-0.3	~179	0.172 <sup>†</sup>	231	+20 <sup>†</sup>	Almanac Office that no air-navigation star was to be left nameless slight var.?: 3.30?–3.36 in V? AAVSO(VSX) as of 2022 Jul. 30 notes existence of NSV entry and availability of 176 AAVSO observations (AAVSO(VSX) assessment: status flag = confirmed variable; variability classification symbol = “VAR”); further photometric study advisable? ¶ m <sub>v</sub> , B–V values are for o UMa AB combined light (where, however, the contribution of o UMa B, shining at mag. ~15, is minuscule) ¶ o UMa A is currently in rapid evolutionary transition, crossing the Hertzsprung Gap ¶ despite high space velocity, a member of the galaxy thin disk	Muscida		
δ	Vel <u>Aa</u> <sup>†</sup> 8 45.4	-54 48	1.96v <sup>†</sup>	0.04 <sup>†</sup>	A1 Va	40	0.0	81	~0.107	~164	+ 2 V?	Aa,Ab brightest known ecl. binary (1.95–2.43, V) orbital period 45.15023 d, primary (resp. secondary) eclipse duration 0.587 d (resp. 0.91 d), with eclipse indicated by photometry to be total; system is β Per type, with average Aa,Ab distance 90.61 au, resolved both interferometrically and with VLT adaptive optics; <a href="#">2007A&amp;A...469..633K</a> offers orbital solution (0.23≤e≤0.37, with orbit inclination to plane of sky near one or other of the two extremes 87.5°, 92.5°), discusses masses, finds unexpectedly large stellar diameters (so both stars are evolved?); B experiences Aa,Ab as essentially a point mass, the Aa,Ab-with-B orbital period being 142 y (at least one orbital solution has been published); B is mag. 5.6, 2.2″→0.8″ (min. angular separation was in 2000), PA 177°→195°, 1894→2019; AAVSO(VSX) assessment in 2022 has status flag = confirmed variable; variability classification symbol = “EA” ¶ WDS documents faint celestial-sphere neighbours C,D,E, and also a sparsely observed close brighter celestial-sphere neighbour F (mag. 5.8, with just 3 observations 1991→1999) ¶ m <sub>v</sub> , B–V values are for δ Vel Aa,Ab,B combined light	Alsephina	
ε	Hya <u>A</u> <sup>†</sup>	8 48.0	+6 20	3.38 <sup>†</sup>	0.68 <sup>†</sup>	G5:III	25	0.4	130	~0.232	259	+36 SB <sup>†</sup>	composite A: 3.8; B: 4.7, 0.2″ (2018); C: 7.8, 2.9″ (2020); B is of poorly known MK type “A:”; orbital solutions have been published both for AB (15.09 y) and for the much wider AB+C system (C experiences AB as essentially a point mass; period is 590 y) ¶ C is unresolved SB, with orbital period 9.9 d ¶ m <sub>v</sub> , B–V values are for ε Hya ABC combined light ¶ our π, D are from 2018 <i>Gaia</i> parallax, which is known to ±2% (since π is stated as 20.7182 ±0.3925 mas), and which we take to supersede 2007 <i>HIPPARCOS</i> ¶ the ε Hya system presents slight variability: AAVSO(VSX), as of 2022 Jul. 30, notes the availability of 52 AAVSO observations, states V-passband range as 3.35–3.39 (AAVSO(VSX) assessment: status flag = confirmed variable; variability classification symbol = “BY:”); further photometric study advisable?	Ashlesha
ζ	Hya	8 56.6	+5 51	3.10	1.00	G9 II–III	~21 <sup>†</sup>	-0.4	~157	0.101	279	+23	¶ Astron. Alm. (epoch 2021.5) assigns MK type G9 IIIa ¶ AAVSO situation in 2022: no status flag (so not confirmed variable, not suspected variable, not confirmed non-variable), and a fortiori no variability classification symbol	
ι	UMa <u>A</u> <sup>†</sup>	9 00.8	+47 57	3.14 <sup>†</sup>	0.20	A7 IVn	~68.9	2.3	47.3	~0.491	~244	+9 SB <sup>†</sup>	A+BC 2.4″, PA 349°→90°, 1831→2017 A+BC orbit 818 y; BC 0.9″, period ~39 y; both the AB binary system and the AB,C binary system (in which C experiences the ι UMa A, ι Uma B binary as essentially a point mass) possess published orbital solutions; A is itself SB, orbit 4028 d, making this a quadruple system; the system is not, as in many	Talitha

$\lambda$	Vel A	9 08.9 -43 32	2.20v <sup>†</sup> 1.66	K4 Ib-IIa <sup>†</sup>	6.0 -3.9 540	0.028 299 +18	cases of multiplicity, hierarchical and stable, but kinematically unstable (disruption in ~0.1 My?); B is mag. 9.9 M1 V and C is mag. 10.1 M1 V; since the A SB has not yet been resolved, even interferometrically, WDS is not yet able to write “ι UMa Aa,” “ι UMa Ab” ¶ AAVSO(VSX) as of 2022 Jul. 30 notes the availability of 42 AAVSO observations and the existence of an NSV entry, suggests a V-passband range of 3.12–3.18, and treats the system as a possible, underexamined, case of slight and rapid variability (AAVSO(VSX) assessment: status flag = suspected variable; variability classification symbol = “S”; is this symbol perhaps a misprint for “S:”?); further photometric study advisable? – WDS indicates a possible line of research by by indicating, in a note, that the ι UMa A SB pairing presents δ Sct variability irregular variable, 2.12–2.32 in V (AAVSO(VSX) assessment in 2022: status flag = confirmed variable; variability classification symbol = “LC”) ¶ probably on or approaching AGB, but could still be on RGB ¶ Astron. Alm. (epoch 2021.5) assigns MK type K4.5 Ib ¶ has slow wind, whose origins are said to be poorly understood	<b>Suhail</b>
a <sup>†</sup>	Car	9 11.6 -59 04	3.43 <sup>†</sup> -0.20	B2 IV–V	7 -2.3 500	0.022 312+23 SB2 <sup>†</sup>	slight variability, 3.41–3.44 in V AAVSO(VSX), as of 2022 Jul. 30, notes availability of a single AAVSO observation, and classifies this tentatively (not with certainty) as Be-star variability without the γ Cas variability common in cases of the “Be phenomenon” (noting that in many such cases the variability is found to be of the λ Eri type; AAVSO(VSX) assessment has status flag = confirmed variable; variability classification symbol = “BE:”); further photometric and further spectroscopic study advisable? ¶ orbit 6.74 d, with light curve indicating tidal distortion; since the SB is as yet unresolved, even in interferometry, WDS is not yet able to write “a Car A,” “a Car B” ¶ there is some uncertainty whether observable light is solely from primary, or whether primary and secondary make approximately equal contributions ¶ not to be confused with α Car	HR 3659
β	Car	9 13.4 -69 49	1.67 0.00	A1 III	28.8 <sup>†</sup> -1.0 113	0.191 305 -5 V?	rapid rotator (< 2.1 d), despite having finished stable core hydrogen fusion ¶ quasi-periodic variation, ~0.5 h, in hydrogen Balmer lines ¶ AAVSO(VSX) situation in 2022: no status flag (so not confirmed variable, not suspected variable, not confirmed non-variable), and a fortiori no variability classification symbol possible slight variability (2.23–2.28 in V?)	<b>Miaplacidus</b>
ι <sup>†</sup>	Car	9 17.7 -59 22	2.25 <sup>†</sup> 0.18	A7 Ib	4.3 -4.6 800	0.022 302 +13	AAVSO(VSX) as of 2022 Jul. 30 has no record of AAVSO observations, notes existence of NSV entry (AAVSO(VSX) assessment: status flag = suspected variable; no variability classification symbol); further photometric study advisable? ¶ despite being slow rotator, has magnetic activity (as inferred from X-rays) ¶ not to be confused with I (letter el) Car B: 8.8, 223”, PA 33°→43°, 1823→2016 suspected var., mag. 3.12–3.17 (beginning to evolve into a Mira?); AAVSO(VSX) as of 2022 Jul. 30 notes the unavailability of any AAVSO observations, notes the existence of NSV entry, cites	Aspidiske
α	Lyn A	9 22.5 +34 18	3.14 <sup>†</sup> 1.55	K7 IIIab	16 -0.8 -203	0.224 274 +38		

													Pannekoek as a discoverer (AAVSO(VSX) assessment: status flag = suspected variable; no variability classification symbol); further photometric study advisable?	
κ	Vel	9 22.8	-55 07	2.49	-0.19	B2 IV-V	6	-3.8	600	0.016	315	+22 SB <sup>†</sup>		Markeb <sup>†</sup>
													IAU name "Markeb" is not to be confused with "Markab" (the IAU name for α Per A) ¶ orbit 116.65 d, average separation possibly ~1.1 au; since the SB has not yet been resolved (even interferometrically), WDS is not yet able to write "κ Vel A" and "κ Vel B" ¶ mass loss rate ~1e-9 M <sub>o</sub> /y ¶ system is X-ray source ¶ AAVSO(VSX) situation in 2022: no status flag (so not confirmed variable, not suspected variable, not confirmed non-variable), and a fortiori no variability classification symbol ¶ ISM absorption has varied over the years (ISM cloud in transit?)	
α	Hya A <sup>†</sup>	9 28.7	-8 46	1.98 <sup>†</sup>	1.44	K3 II-III <sup>†</sup>	18	-1.7	180	0.038	336	-4 V?	possible slight variability (1.93-2.01 in V?) AAVSO(VSX), as of 2022 Jul. 30, notes existence of NSV entry and the availability of a single AAVSO observation (AAVSO(VSX) assessment: status flag = suspected variable; no variability classification symbol); further photometric study advisable? ¶ slow rotator (possibly 2.4 y), with Ba mildly overabundant ¶ asteroseismology has been studied ¶ α Hya B (mag. 9.7; 284", PA 55°→155°, 1833→2015) might be a true binary component (with orbit ≥ 870,000 y, separation ≥ 15,700 au)	Alphard
N	Vel	9 31.9	-57 08	3.14 <sup>†</sup>	1.55	K5 III	13.6	-1.2	240	0.033	280	-14	slight semiregular var., 3.12-3.18 in V, 82.0 d (AAVSO(VSX) assessment in 2022: status flag = confirmed variable; variability classification symbol = "SR") ¶ evolutionary status uncertain (helium core fusion impending, or already ended?)	HR 3803
θ	UMa A	9 34.4	+51 34	3.18 <sup>†</sup>	0.46	F6 IV <sup>†</sup>	74.2	2.5	44.0	1.088	241	+15 SB <sup>†</sup>	possibly slight variability (3.16-3.20 in V) AAVSO(VSX), as of 2022 Jul. 30, notes existence of NSV entry, notes the unavailability of AAVSO observations (AAVSO(VSX) assessment: status flag = suspected variable; no variability classification symbol); further photometric study advisable? ¶ luminosity class, and also SB status, have been controverted, with postulated SB companion remaining undetected in speckle interferometry ¶ m <sub>v</sub> , B-V values are for θ UMa AB combined light (where, however, the contribution of θ UMa B, at mag. -14, is minuscule)	
o	Leo <u>Aa</u> <sup>†</sup>	9 42.4	+9 47	3.52	0.49	F9 III + A5m	24.2	0.5	135	0.148	255+27	SB2 <sup>†</sup>		Subra
													IAU name "Subra" applies only to o Leo Aa; Aa,Ab is a tight binary, with angular separation ~4 mas, but nevertheless now interferometrically resolved ( <a href="#">2001AJ....121.1623H</a> ; from this paper we take MK type, and also our π-cum-D (as a distance derived from comparing seen angular size of orbit with orbit physical size, yielding a result in good agreement with the 2007 <i>HIPPARCOS</i> trigonometric parallax, and yet with smaller uncertainty); period is 14.498 d, with orbit nearly circular, distance between the two stars 0.165 au; our assertion of eclipsing in earlier editions of the Handbook was erroneous, although it is true that the orbit is seen rather close to edge-on; lunar occultation as reported in <a href="#">1978AJ....83.1100A</a> failed to split Aa,Ab ¶ m <sub>v</sub> , B-V values are for o Leo Aa,Ab combined light ¶ AAVSO(VSX) situation in 2022: no status flag (so not confirmed variable, not suspected variable, not confirmed non-variable), and a fortiori no variability type symbol ¶ o Leo Aa is a rare instance of a star that has ended	

												core hydrogen fusion, and yet in which the convection typical of an evolved star has not removed the chemical peculiarities possible in a core-hydrogen fuser (where the still-quiet atmosphere facilitates radiative lofting and gravitational settling)
I <sup>†</sup>	Car	9 45.9 -62 37	3.36v <sup>†</sup> 1.02	F9–G5 Ib	2	-4.7 2000	0.015 302	+3 V				<p> <b>[THIS STAR ONLY IN ONLINE VERSION OF TABLE]</b>            Cepheid variable: 3.28–4.18 in V, 36 d            (AAVSO(VSX) assessment in 2022: status flag = confirmed variable; variability classification symbol = “<b>DCEP</b>”); AAVSO(VSX) as viewed both 2021 Jan. 18 2022 Jul. 13 gives period 35.551609 d; an exceptionally luminous, and consequently exceptionally slow, Cepheid (compare both the visual brightness and the intrinsic luminosity with less dramatic δ Cep A (in this table), η Aql A (Okab; in this table), and ζ Gem Aa (Mekbuda; almost, but not quite, bright enough for inclusion in this table): Kaler remarks that “if Carina had been in the northern hemisphere, the collection of these variables might well have been called the ‘Carinids’”); radius, in its pulsation cycle, has been measured as 160 R<sub>⊙</sub> min, 194 R<sub>⊙</sub> max            ¶ circumstellar envelope of ejected matter, radius 10 au–100 au            ¶ lower-case ell Car; not to be confused with i (lower-case i) Car (HR 3663), I Car (HR 3699), L Car (HR 4089), I (upper-case i) Car (HR 4102) (and note additionally that Bayer nomenclature does not use the label “λ Car”)  <b>[THIS STAR ONLY IN ONLINE VERSION OF TABLE]</b>            poss. slight variability (2.95–3.04 in V?)            AAVSO(VSX), as at 2022 Jul. 30, notes the availability of 3 AAVSO observations, and notes existence of NSV entry (AAVSO(VSX) assessment: status flag = suspected variable; no variability classification symbol; further photometric study advisable?); some photometric work is possibly reported in <a href="#">1998AN...319.239A</a> (pulsation as in Cepheids?)            ¶ slow rotator, period possibly as long as 200 d            ¶ currently residing in the Hertzsprung Gap?            ¶ the Arabic or quasi-Arabic name Algenubi (more classically, al Ras al Asad al Janubiyah et al.) is not presently IAU-official            A: 3.01; B: 5.99, B7 III, 5.1”, PA126°→128°, 1836→2015 orbit ≥ 19,500 y, separation ~2000 au            ¶ m<sub>v</sub>, B–V values are for v Car AB combined light            ¶ AAVSO(VSX) situation in 2022: no status flag (so not confirmed variable, not suspected variable, not confirmed non-variable), and a fortiori no variability type symbol            ¶ the duplicity causes parallax to be poorly known            ¶ m<sub>v</sub>, B–V values are for φ Vel A alone (however, the contribution of B, at mag. ~12, to the φ Vel AB combined light is minuscule)            ¶ AAVSO(VSX) situation in 2022: no status flag (so not confirmed variable, not suspected variable, not confirmed non-variable), and a fortiori no variability type symbol  <b>[THIS STAR ONLY IN ONLINE VERSION OF TABLE]</b>            B: 8.4, 0.4”, PA 84°→239°, 1937→2015            ¶ possibly variable (3.46–3.60 in V?); AAVSO(VSX), as of 2022 Jul. 30, notes existence of NSV entry, notes that no AAVSO observations are available for the η Leo system            (AAVSO(VSX) assessment: status flag = suspected variable; no variability classification symbol); further photometric study advisable?            ¶ m<sub>v</sub>, B–V values are for η Leo AB combined light            ¶ mass-loss rate ~5e-8 M<sub>⊙</sub>/y (&gt; 10,000× solar mass-loss rate); BSC5: “chromospheric shell”         </p>
ε	Leo	9 47.2 +23 40	2.98 <sup>†</sup> 0.81	G1 II	13.2	-1.4 250	0.047 259	+4 V?				
v	Car A	9 47.7 -65 11	2.96 0.27	A6 II	2.3 <sup>†</sup>	-5.3~1400 <sup>†</sup>	0.028 307	+14				
φ	Vel A	9 57.7 -54 41	3.53 -0.09	B5 Ib	2.0	-4.9 1600	0.014 285	+14				
η	Leo A	10 08.6 +16 39	3.49v? <sup>†</sup> -0.03 <sup>†</sup>	A0 Ib <sup>†</sup>	3	-4.5 1300	~0.003	n.a.				



$\alpha$  Leo A<sup>†</sup> 10 09.6 +11 51 1.36<sup>†</sup> -0.11<sup>†</sup> B8 IVn<sup>†</sup> 41 -0.6 79 0.249 271 +6 SB<sup>†</sup>

¶ we follow WDS in asserting duplicity (with WDS indicating that the A,B pair has been split both through lunar occultation and in speckle interferometry, and documenting 7 observations 1937→2015; the assertion is, admittedly, questioned by Kaler at [stars.astro.illinois.edu/sow/etaleo.html](http://stars.astro.illinois.edu/sow/etaleo.html))

**Regulus**

$\alpha$  Leo A is SB orbit 40.11 d, with the secondary in the pair that is  $\alpha$  Leo A now detected ([2011IBVS.5987....1R](https://arxiv.org/abs/2011IBVS.5987....1R) reported null photometry result from *MOST*, at the high precision of ~0.5 millimag., but a spectroscopic detection is reported in [2020ApJ...902...25G](https://arxiv.org/abs/2020ApJ...902...25G); since the secondary is not yet resolved, even interferometrically, WDS is not yet able, at any rate as of 2022 Mar. 3, to write “ $\alpha$  Leo Aa” and “ $\alpha$  Leo Ab”); at the level of coarse photometry, slight variability has been asserted, with AAVSO(VSX) showing possible range 1.33–1.40 in V; as of 2022 Jul. 30, AAVSO(VSX) notes existence of NSV entry, notes the unavailability of AAVSO observations (AAVSO(VSX) assessment: status flag = suspected variable; no variability classification symbol); further photometric study advisable?  
 ¶  $m_v$ , B–V values are for the combined light of the  $\alpha$  Leo A SB two-star system  
 ¶ the primary in  $\alpha$  Leo A is an exceptionally rapid rotator (15.9 h), making the star an oblate spheroid ( $R_{\text{pol}} \sim 3.14 R_{\odot}$ , but  $R_{\text{eq}} \sim 4.16 R_{\odot}$ ) and rendering the photosphere equator ~3000 K cooler than the photosphere poles (and possibly inducing meridional flow in the envelope); this is the first rotating star not in an eclipsing binary system to have its gravitational low-latitudes darkening detected, and the first to have its inclination angle and low-latitudes darkening measured through a direct application of spectroscopy-constrained interferometry (inaugural science run of CHARA, [2005ApJ...628.439M](https://arxiv.org/abs/2005ApJ...628.439M)); in contrast with pole-on rapid rotators such as  $\alpha$  Lyr A, the  $\alpha$  Leo A primary is seen nearly equator-on; in [2011ApJ...732...68C](https://arxiv.org/abs/2011ApJ...732...68C), Fig. 5 presents an image as fitted to CHARA interferometry (the luminosity contours display the perturbing effect of limb darkening upon the rotation-induced gravity darkening; since the poles are near the limb, the brightest regions, as viewed from Earth, do not quite coincide with the poles); the aperture-synthesis imaging of [2017NatAs...1..690C](https://arxiv.org/abs/2017NatAs...1..690C) Fig. 5 displays the photosphere temperature variation (a joint consequence of limb darkening and oblateness), along with oblateness and axis orientation; according to [2017NatAs...1..690C](https://arxiv.org/abs/2017NatAs...1..690C), the  $\alpha$  Leo A primary (i) has attained 96.5% of its breakup speed (earlier literature had suggested 86%), and (ii) is the first rapid rotator found to exhibit Chandrasekhar rotation-induced stellar limb polarization (the related phenomenon of eclipse-induced stellar limb polarization was admittedly detected earlier, with  $\beta$  Per, as reported in [1983ApJ...273L..85K](https://arxiv.org/abs/1983ApJ...273L..85K))  
 ¶ despite the large rotation-induced latitude variation in photospheric effective temperature, at all latitudes the envelope is radiative (since the photospheric effective temperature, even at the equator, never falls so low as to approach the ~8300 K radiation-to-convection transition value); rotation-induced meridional circulation, on the other hand, disturbs the usual radiative-equilibrium picture of a radiative envelope ([2011ApJ...732...68C](https://arxiv.org/abs/2011ApJ...732...68C), p. 11a); since meridional circulation transfers angular momentum, the envelope cannot be presumed to be in solid-body rotation  
 ¶ the rapid rotation, the membership in MK type B, and the near-MS evolutionary status notwithstanding, the question of Be-phenomenon behaviour

is answered in the negative by [2005ApJ...628.439M](#); the authors do, however, remark on p. 446 that the historical record contains a lone report of marginal hydrogen Balmer- $\alpha$  emission, from February 1981 (might amateur-spectroscopist monitoring now be advisable?)

¶ [2011ApJ...732...68C](#) revises the mass of the primary upward, offering 4.15  $M_{\odot}$  in place of the [2005ApJ...628.439M](#) determination of  $\sim 3.5 M_{\odot}$  ¶ A+BC almost unchanged since 1779 (179"; PA 307°→304°, 1779→2019); AB is nevertheless known to be a true binary pairing, rather than a mere line-of-sight coincidence; AB distance  $\geq 4200$  au, orbit  $\geq 125,000$  y; BC combined-light  $m_v$ , B-V are 8.13, 0.88; BC is no longer underobserved (PA: 89°→94°, 4.0"→2.20", 1867→2019, with orbit  $\geq 880$  y)

¶ a puzzling discrepancy between the ages of the  $\alpha$  Leo SB primary and  $\alpha$  Leo B (surely condensed from the same ISM cloud, at the same time) is perhaps to be explained by the peculiarities in the evolution of rapid rotators

¶ we adopt here the MK classification of [2003AJ...126.2048G](#), while recalling that earlier editions of our RASC brightest-stars table used instead B7 Vn, essentially in accordance with [1953ApJ...117..313J](#); Astron. Alm. (epoch 2021.5) likewise assigns MK type B7 Vn

¶ the  $\alpha$  Leo system is occasionally occulted by Mercury, Venus (e.g. 1959 Jul. 07, 2044 Oct. 01), Moon (e.g. 2017 Sep. 18; [1972JBA...82.431K](#) describes the 18.6-year 1940-through-2050 cycle of possibilities), and asteroids (e.g. 166 Rhodope 2005 Oct. 19 ([2008mgm.conf.2594S](#) reports GTR effect of light bending, not only from general solar gravitational field but also from Rhodope field), 163 Erigone 2014 Mar. 20 (cloud-defeated 2014 Erigone campaign is documented at [occultations.org/regulus2014](#))

¶ E(B-V) = +0.01 or "IIIne"; shell star ¶ rapid rotator (< 1.2 d,  $\sim 85\%$  of breakup speed); instance of "Be phenomenon"; AAVSO(VSX), following GCVS, on the one hand asserts slight variability (3.30–3.32 in V), as would be consistent with  $\lambda$  Ori-type variability, and on the other hand implicitly denies the  $\gamma$  Cas-type outburst variability that often accompanies the "Be phenomenon" (AAVSO(VSX) assessment in 2022: status flag = confirmed variable; variability classification symbol = "BE"); on the side of spectroscopy, as distinct from photometry, BSC5 reports variable hydrogen Balmer- $\alpha$  irregular variable, 3.34–3.44 in V HR 4050 (AAVSO(VSX) assessment in 2022: status flag = confirmed variable; variability classification symbol = "LC")

¶ metallicity is uncertain ¶ evolutionary state is uncertain (has core already started He fusion?) ¶ Astron. Alm. (epoch 2021.5) assigns MK type K2.5 II B (6.0, 331", PA 337°) is mere optical companion Adhafera (since A,B parallax discrepancy is large, 12 mas for  $\zeta$  Leo A, but 33 mas for the decidedly less distant  $\zeta$  Leo B)

¶ AAVSO(VSX), as of 2022 Jul. 30, has no record of available AAVSO observations, offers possible range 3.42–3.46 in V passband (AAVSO(VSX) assessment: status flag = suspected variable;

$\omega$  Car 10 14.3–70 09 3.31<sup>†</sup> -0.09 B8 IIIIn<sup>†</sup> 9.5 -1.8 340 0.037 281 +7 V

q Car A 10 17.9–61 27 3.36v<sup>†</sup> 1.54 K3 IIa<sup>†</sup> 5.0 -3.1 660 0.026 286 +8

$\zeta$  Leo A 10 18.0+23 18 3.44<sup>†</sup> 0.31 F0 IIIa<sup>†</sup> 12 -1.2 270 0.020 110 -16 SB

												no variability classification symbol); further photometric study advisable? ¶ Astron. Alm. (epoch 2021.5) assigns MK type F0 III ¶ in rapid evolutionary transition, currently residing in Hertzsprung Gap
λ	UMa	10 18.5+42 48	3.45	0.03	A1 IV <sup>†</sup>	24	0.3	140	0.186	256	+18 V	Tania Borealis despite MK luminosity class “IV,” has not yet finished core hydrogen fusion ¶ mildly metallic, but insufficiently metallic to warrant MK “Am” ¶ AAVSO(VSX) situation in 2022: no status flag (so not confirmed variable, not suspected variable, not confirmed non-variable), and a fortiori no variability classification flag ¶ seems mild IR excess (indicating circumstellar debris) 4.7” (2020), PA 99°→127°, 1820→2020 (510.3 y); Algieba max = ~5”, around 2100 separation ≥ 170 au, orbital parameters not yet well known ¶ A, B are of mildly unequal masses, and therefore are of mildly disparate evolutionary stage (Kaler <a href="http://stars.astro.illinois.edu/sow/algieba.html">stars.astro.illinois.edu/sow/algieba.html</a> : “best understood as being in different stages of gianthood”; cf this same source for further discussion of the uncertainties in various γ Leo parameters, including the respective mags of A and B) ¶ we take m <sub>v</sub> , B–V values (following normal procedure in this Handbook table) from GCPD, but correcting here what we believe to be a clerical error in GCPD, leading to “AB” being stated as of 2022 Aug. 05 where in our view “B” was meant: additionally, GCPD gives, as directly measured m <sub>v</sub> , B–V values for γ Leo AB combined light, 1.98, 1.14 ¶ γ Leo A, or γ Leo B, or both, may harbour variability: AAVSO(VSX), as of 2022 Jul. 30, notes existence of NSV entry, notes availability of 180 AAVSO observations, offers V-passband range 1.84–2.03, presumably for AB combined light (AAVSO(VSX) assessment: status flag = suspected variable; no variability classification symbol); further photometric study advisable? ¶ γ Leo A “+ 1P” is an exception to the tendency for exoplanets to be found around the more metallic stars (but the “+1P” could be modelled as a brown dwarf); and indeed even “+2P” is now considered possible ¶ high space velocity of the γ Leo AB pair, plus their low metallicity, suggests system is interloper from more remote galactic region ¶ WDS documents celestial-sphere neighbours γ Leo Ca, Cb, D,E,F, of which all but Ca and Cb are fainter than mag. 10; Ca and Cb, a tight pair first split in 1981, shine with a combined light of mag. 9.64, the contribution coming almost entirely from Ca (the faint Cb has been detected only at 750 nm), and are widely separated from A (341”, at PA 288°, in 2016); the AC pairing is found through analysis of the respective proper motions to be a mere line-of-sight coincidence (211”→341”, PA 294°→288°, 1851→2016) ¶ γ Leo AB, and indeed also the next “Sickle” star ζ Leo, serve to mark the radiant of the Leonids meteor shower slight var., 3.03–3.10 in V Tania Australis AAVSO(VSX) considers the system to present slow irregular evolved-star variability, and additionally to possibly present β Lyr-type eclipses (AAVSO(VSX) assessment in 2022: status flag = confirmed variable; variability classification symbol = “ <b>EB</b> :+ <b>LB</b> ”) ¶ SB period 230 d; since the SB is not yet resolved, even interferometrically, WDS cannot yet write “μ UMa A” and “μ UMa B”; ¶ Ca II emission ¶ Astron. Alm. (epoch 2021.5) assigns MK type M0 III
γ	Leo A +1P <sup>†</sup>	10 21.3+19 43	2.29v <sup>†</sup>	1.20 <sup>†</sup>	K1 IIIb Fe–0.5 <sup>†</sup>	26	–0.3	130	–0.333 <sup>†</sup>	~118	–37 <sup>†</sup> SB	
γ	Leo B	10 21.3+19 43	3.46v <sup>†</sup>	0.92 <sup>†</sup>	G7 III Fe–1 <sup>†</sup>	26	0.2	130	–0.346 <sup>†</sup>	~118	–36 <sup>†</sup> V	
μ	UMa	10 23.7+41 23	3.05 <sup>†</sup>	1.58	M0 IIIp <sup>†</sup>	14	–1.2	230	0.089	293	–21 SB <sup>†</sup>	

p	Car	10 32.9–61 48	3.31 <sup>v</sup> –0.10	B4 Vne <sup>†</sup>	7	–2.6	500	0.021	304	+26	<p>¶ Kaler (<a href="https://stars.astro.illinois.edu/sow-taniaas.html">stars.astro.illinois.edu/sow-taniaas.html</a>) terms this “a rare ‘hybrid star’” (in the sense of blowing both a fast-and-thin wind and a slower-and-dense wind), and additionally notes the puzzle posed by X-ray emission in the presence of cool photosphere variable of <math>\gamma</math> Cas type, 3.22–3.55 in V and instance of the “Be phenomenon” (AAVSO(VSX) assessment in 2022: status flag = confirmed variable; variability classification symbol = “GCAS”</p> <p>¶ fast rotator; BSC5: shell; variable hydrogen Balmer-line profiles</p>
$\theta$	Car	10 43.8–64 31	2.76 <sup>†</sup> –0.22 <sup>†</sup>	B0.5 Vp	7	–3.0	460	0.022	303	+24 SB <sup>†</sup>	<p>chemically anomalous; of three published orbital solutions for this unresolved SB, the two that seem most reliable (from 1995, 1988) assert periods of 2.20 d, 2.13 d respectively (with <math>e</math> values 0.0, 0.24 respectively); the short SB period, even given the low <math>e</math>-values (i.e. the lack of severely close periastron passages) suggests mass transfer could be the culprit in the anomalies</p> <p>¶ since the SB is not as yet resolved, even interferometrically, WDS is not yet able to write “<math>\theta</math> Car A,” “<math>\theta</math> Car B”</p> <p>¶ <math>m_v</math>, B–V values seem unavailable in GCPD; we take these values from <a href="#">2002yCar.2237...0D</a></p> <p>¶ slight variability detected by <i>TESS</i> mission, range 0.003 mag. in V</p> <p>(AAVSO(VSX) assessment in 2022: status flag = confirmed variable; variability classification symbol = “HB”</p> <p>¶ the primary is the brightest of the “blue stragglers”; at <a href="https://stars.astro.illinois.edu/sow/thetacar.html">stars.astro.illinois.edu/sow/thetacar.html</a>. Kaler discusses difficulties in determination of the primary’s temperature and of its (short) rotation period</p> <p>¶ E(B–V) = +0.06</p> <p>A: 2.82; B: 5.65, 2.3”, PA 55°→58°, 1880→2019 period variously given as 116.24 y (Hoffleit) and 138 y (Heintz); A-to-B distance possibly 8 au min, 93 au max, 51 au average</p> <p>¶ A is itself an SB, not as yet resolved, even in interferometry (so WDS is not as yet able to write “<math>\mu</math> Vel Aa,” “<math>\mu</math> Vel Ab”)</p> <p>¶ <math>m_v</math>, B–V values are for <math>\mu</math> Vel AB combined light</p> <p>¶ AAVSO(VSX) situation in 2022: no status flag (so not confirmed variable, not suspected variable, not confirmed non-variable), and a fortiori no variability classification symbol</p> <p>¶ <math>\mu</math> Vel B is of uncertainty-flagged MK type “F8: V”</p> <p>¶ one or the other component of the SB that is <math>\mu</math> Vel A is in rapid evolutionary transition, having recently finished core hydrogen fusion</p> <p>¶ one or other component of the SB that is <math>\mu</math> Vel A is magnetic, and an X-ray emitter, with hot corona, and with violent 2-day X-ray flare detected in 1998 by IUE</p>
$\mu$	Vel $\underline{A}$ <sup>†</sup>	10 47.8–49 33	2.68 <sup>†</sup> 0.90 <sup>†</sup>	G5 III <sup>†</sup>	28	–0.1	~117	0.083	131	+6 SB	<p>¶ A is itself an SB, not as yet resolved, even in interferometry (so WDS is not as yet able to write “<math>\mu</math> Vel Aa,” “<math>\mu</math> Vel Ab”)</p> <p>¶ <math>m_v</math>, B–V values are for <math>\mu</math> Vel AB combined light</p> <p>¶ AAVSO(VSX) situation in 2022: no status flag (so not confirmed variable, not suspected variable, not confirmed non-variable), and a fortiori no variability classification symbol</p> <p>¶ <math>\mu</math> Vel B is of uncertainty-flagged MK type “F8: V”</p> <p>¶ one or the other component of the SB that is <math>\mu</math> Vel A is in rapid evolutionary transition, having recently finished core hydrogen fusion</p> <p>¶ one or other component of the SB that is <math>\mu</math> Vel A is magnetic, and an X-ray emitter, with hot corona, and with violent 2-day X-ray flare detected in 1998 by IUE</p>
v	Hya	10 50.8–16 19	3.12 1.24	K2 III <sup>†</sup>	23	–0.1	144	0.220 <sup>†</sup>	25	–1 <sup>†</sup>	<p>slow rotator (but <math>\leq 619</math> d)</p> <p>¶ low metallicity and high space velocity suggest interloper, born outside Sun’s neighbourhood</p> <p>¶ Astron. Alm. (epoch 2021.5) assigns MK type K1.5 IIIb H<math>\delta</math>–0.5</p> <p>¶ AAVSO(VSX) situation in 2022: no status flag (so not confirmed variable, not suspected variable, not confirmed non-variable), and a fortiori no variability classification symbol</p>
$\beta$	UMa	11 03.2+56 15	2.37 <sup>†</sup> –0.02	A0mA1 IV–V	~40.9	0.4	80	0.088	68	–12 SB	<p>possible slight variability (2.35–2.40 in V?) Merak</p> <p>AAVSO(VSX), as of 2022 Jul. 30, notes existence of NSV entry and availability of 77 AAVSO observations</p> <p>(AAVSO(VSX) assessment in 2022: status flag = suspected variable; no variability classification symbol assigned); further photometric study advisable?</p> <p>¶ debris disk first detected via IR excess, now marginally resolved by <i>Herschel Space Observatory</i></p>



												at La Silla (ESO) in 2002 demonstrated for the first time the feasibility of asteroseismology for a highly evolved star: $\xi$ Hya Aa has left the theoretical MS, being now near the theoretical SGB-RGB transition, and yet CORALIE was able to find solar-like oscillations (with periods, however, of 2.0 h to 5.5 h, in contrast with the “five-minute oscillations” in the Sun; $\xi$ Hya Aa, being evolved, is larger than the Sun, and so its starquakes face larger propagation distances; on the other hand, where the solar “five-minute oscillations” involve speeds of $\sim 15$ cm/s or $\sim 20$ cm/s, in the case of $\xi$ Hya Aa the observed speeds attain values only a little below 2 m/s (such a refined stellar radial-velocity measurement is now possible as a kind of spinoff from advances in exoplanet-search spectrograph engineering); both radial and non-radial oscillations are found in $\xi$ Hya Aa; the pertinent ESO press release is at <a href="http://www.eso.org/public/news/eso0215/">www.eso.org/public/news/eso0215/</a> , and the modern $\xi$ Hya Aa asteroseismology literature in the journals starts with the “Letter” which is <a href="#">2002A&amp;A...394L...5F</a> [THIS STAR ONLY IN ONLINE VERSION OF TABLE]
$\lambda$	Cen Aa <sup>†</sup>	11 36.9–63 09	3.12 <sup>†</sup>	–0.04 <sup>†</sup>	B9.5 IIn <sup>†</sup>	8	–2.4	400	0.034	258	–1 V	despite possible fast rotation ( $< 2.7$ d?), Fe is overabundant, with Si and C mildly underabundant ¶ at <a href="http://stars.astro.illinois.edu/sow/lambda_cen.html">stars.astro.illinois.edu/sow/lambda_cen.html</a> . Kaler discusses questions of visual binarity ( $\lambda$ Cen Aa, Ab, B: but WDS documents only a single observation of $\lambda$ Cen Ab, from the year 2000, and just 3 observations of the faint $\lambda$ Cen B, over the period 1937→2015) ¶ $m_v$ , B–V values are for $\lambda$ Cen Aa, Ab combined light ¶ AAVSO(VSX) situation in 2022: no status flag (so not confirmed variable, not suspected variable, not confirmed non-variable), and a fortiori no variability classification symbol
$\beta$	Leo A	11 50.3+14 26	2.14 <sup>†</sup>	0.09	A3 Va	91	1.9	36	0.511	257	0 V	<b>Denebola</b> very slight var., range 0.025 in V passband (AAVSO(VSX) assessment in 2022: status flag = confirmed variable; variability classification symbol = “DSCTC”) ¶ rapid rotator ( $< 0.65$ d) ¶ debris disk resolved by <i>Herschel Space Observatory</i> (2010A&A...518L.135M), disk structures differentiated with ground-based interferometry (2010ApJ...724.1238S) ¶ WDS documents very faint celestial-sphere neighbours $\beta$ Leo B,C, and additionally the not-so-faint D (mag. 8.5; AD astrometry is 298″→240″, PA 204°→190°, 1833→2019) and the tight (0.5″) pair Ea,Eb (mags. 6.5 and 6.6; this pair is separated by 2″ from A; but for the “AE” pairing WDS has just a single measurement, from 2009, and likewise for the Ea,Eb pairing WDS has just a single measurement, again from 2009)
$\gamma$	UMa A	11 55.1+53 34	2.44 <sup>†</sup>	0.00	A0 Van <sup>†</sup>	39	0.4	83	0.108	84	–13	Phecda rapid rotator: although in MK temperature class A, nevertheless an instance of the “Be phenomenon” (the term “Ae star” is sometimes used for this rare category); following GCVS, AAVSO(VSX) asserts slight variability for the $\gamma$ UMa system, offering range 2.41–2.45 in V passband, while refraining from asserting the eruptive $\gamma$ Cas-type variability often found with the “Be phenomenon” (could this also be an instance of $\lambda$ Eri-type variability?); additionally, AAVSO(VSX) asserts eruptive UV Ceti-type variability in the $\gamma$ UMa system (AAVSO(VSX) assessment in 2022: status flag = confirmed variable; variability classification symbol = “BE+UV”) ¶ $\gamma$ UMa A with an invisible companion (or is this, rather, the sparsely observed pairing of $\gamma$ UMa A with (visible, mag. 8.2 $\gamma$ UMa B?) is said in <a href="http://en.wikipedia.org/wiki/Gamma_Ursae_Majoris">en.wikipedia.org/wiki/Gamma_Ursae_Majoris</a> to be an astrometric binary, of period 20.5 y (and we are also not fully certain on the SB situation:

											although we have in the past applied the flag “SB” (is $\gamma$ UMa A a binary with companion unresolved?), we now withdraw the flag, on the strength of <a href="#">2010NewA...15..324G</a> (which proposes an astrometric binary-system orbit for $\gamma$ UMa AB, but additionally writes, “Spectroscopic duplicity of this star mentioned in some catalogues seems to be a mistake: it [sc the alleged duplicity] could not have been detected because of a large rotational velocity [with rotational broadening, therefore, of the spectrum lines]”) ¶ E(B–V) = 0.00
$\delta$	Cen Aa <sup>†</sup>	12 09.6–50 51	2.58v <sup>†</sup> –0.13	B2 IVne <sup>†</sup>	8	–2.9	400	0.050	262	+11 V	var. of $\gamma$ Cas type, 2.51–2.65 in V (AAVSO(VSX) assessment in 2022: status flag = confirmed variable; variability classification symbol = “GCAS”) ¶ rapid rotator (< 1.3 d), with shell spectrum; <a href="#">2008A&amp;A...488L..67M</a> summarizes recent research, and as part of a wider VLT investigation into the “Be phenomenon” not only discusses the circumstellar ejecta, but also reports discovery of binarity (Ab at angular distance 68.7 mas); WDS also documents celestial-sphere neighbours $\delta$ Cen B, $\delta$ Cen C; $\delta$ Cen B (a Be star) is mag. 4.4, with astrometry 221”→269”, PA 325° →325°, 1847→1999; $\delta$ Cen C is mag. 6.4, with astrometry 218”→217”, PA 227°→227°, 1847→1999
$\varepsilon$	Crv	12 11.3–22 45	3.00 <sup>†</sup>	1.33 K2 III <sup>†</sup>	–10.3	–1.9	320	0.072	278	+5	possible slight var (2.98–3.06 in V passband?) AAVSO(VSX), as of 2022 Jul. 30, notes existence of NSV entry and availability of 22 AAVSO observations (AAVSO(VSX) assessment in 2022: status flag = suspected variable; no variability classification symbol ); further photometric study advisable? ¶ slow rotator (but $\leq$ 3.9 y) ¶ metals somewhat overabundant ¶ evolutionary status uncertain (core-helium fusion starting, in progress, or finished?) ¶ Astron. Alm. (epoch 2021.5) assigns MK type K2.5 IIIa ¶ the etymologically Arab name “Minkar” is of merely modern origin, and is not currently IAU-official
$\delta$	Cru	12 16.4–58 53	2.79 <sup>†</sup> –0.24	B2 IV <sup>†</sup>	9.4	–2.3	350	0.037	254	+22 V?	slight var., $\beta$ Cep type, 2.78–2.80 in V, 3.62 h Imai (AAVSO(VSX) assessment in 2022: status flag = confirmed variable; variability classification symbol = “BCEP”) ¶ rapid rotator (<1.3 d; BSC5: “expanding circumstellar shell”)
$\delta$	UMa A	12 16.6+56 54	3.31	0.08 A2 Van	40.5	1.4	81	0.104	86	–13 V	Megrez AAVSO(VSX), as of 2022 Jul. 30, indicates, as a preliminary survey result, slight variability, range 3.30–3.31 in V passband, with period 11.435 h, with 2493 AAVSO observations available (AAVSO(VSX) assessment: status flag = confirmed variable; variability classification symbol = “VAR”); further photometric study advisable? ¶ possesses debris disk, of unusually low radius (Wyatt et al 2007; Pointing–Robertson drag?)
$\gamma$	Crv	12 17.0–17 40	2.58 <sup>†</sup> –0.11	B8 III <sup>†</sup>	21	–0.8	154	0.160	278	–4 SB	slight var.? (2.56–2.60 in V?) AAVSO(VSX), as of 2022 Jul. 30, notes existence of NSV entry and availability of 22 AAVSO observations, credits discovery to Ferrero (AAVSO(VSX) assessment: status flag = suspected variable; no variability classification symbol); further photometric study advisable? ¶ rather rapid rotation notwithstanding (BSC5: “expanding circumstellar shell”), Hg and Mn are overabundant, with some other elements underabundant (but rotational line broadening makes abundance determinations difficult); Astron. Alm. (epoch 2021.5) assigns MK temperature type B8p Hg Mn, and does not assign an MK luminosity class
$\alpha$	Cru A <sup>†</sup>	12 27.9–63 14	1.28 <sup>†</sup> –0.18 <sup>†</sup>	B0.5 IV	10	–3.7	~320	0.037	251	–11 SB	5.4” (1826); 35” (2020) <b>Acrux</b>
$\alpha$	Cru B <sup>†</sup>	12 27.9–63 14	1.58 <sup>†</sup> –0.17 <sup>†</sup>	B1 Vn	10	–3.3	~320	0.037?	251?	–1	PA 114°→111°, 1826→2020

orbit  $\geq 1300$  y, AB distance  $\sim 430$  au (the WDS “U” flag for AB, indicating that the pairing is a mere line-of-sight coincidence, does not seem supported in the most obvious part of the secondary literature [Kaler, *Wikipedia*]); A is SB pair (not as yet resolved, even in interferometry, so WDS cannot as yet write “ $\alpha$  Cru Aa,” “ $\alpha$  Cru Ab”), with period 75.78 d, distance between components  $\sim 0.5$  au min,  $\sim 1.5$  au max; C (itself an SB pair, unresolved), at mag 4.8, has been said to be imperfectly sharing AB proper motion (AC astrometry  $92'' \rightarrow 89''$ , PA  $216^\circ \rightarrow 203^\circ$ , 1750  $\rightarrow$  2020), and so is possibly (not assuredly) gravitationally bound with the putatively gravitationally bound three-star AB system (if C is bound, then period  $> 130,000$  y, with distance from AB  $\geq 9,000$  au); the other celestial-sphere neighbours of the unresolved  $\alpha$  Cru A SB pair are fainter than mag. 10; the IAU-official name “Acrux” applies to the “primary” (the more luminous, and more massive) component of the unresolved  $\alpha$  Cru A pair; that unresolved pair has also been called  $\alpha^1$  Cru, with  $\alpha$  Cru B correspondingly called  $\alpha^2$  Cru; it is a little puzzling that the WDS note for this  $\alpha$  Cru ABC putatively 5-star system is silent on SB status of  $\alpha$  Cru C, while suggesting instead (on strength of 1967 Batten work at DAO) possible SB status for  $\alpha$  Cru B

¶ the AB duplicity makes the two individual mag. and colour determinations “Aa-with-Ab-combined light,” “B” somewhat controverted: we use [2002A&A...384..180F](#) for these two individual determinations; our normal authority, GCPD, for its part does not give separate determinations, but does give for  $\alpha$  Cru Aa,Ab,B combined light 0.76,  $-0.25$

¶ AAVSO situation in 2022: no status flag (so not confirmed variable, not suspected variable, not confirmed non-variable), and a fortiori no variability symbol B:8.4, K2 V, 24", PA  $216^\circ \rightarrow 216^\circ$ , 1782  $\rightarrow$  2020 Algorab although A,B have common proper motion, disparity in age estimates has caused binarity to be questioned; Kaler, accepting binarity (he proposes period  $\geq 9400$  y) suggests that the  $\delta$  Crv AB system is young, and that B (radiating less powerfully than A) is a post-T-Tauri star (i.e. a star that, although already stably burning core hydrogen, has nevertheless not yet succeeded in clearing away its surrounding dust)

¶  $m_v$ , B-V values are for  $\delta$  Crv A; for  $\delta$  Crv B, GCPD gives 8.40, 0.87

¶ AAVSO(VSX) assessment in 2022: status flag = suspected variable;

no variability classification symbol slight semiregular var., 1.60–1.67 in V although classified by AAVSO(VSX) as semiregular var.,

at least 6 pulsation periods have been documented (AAVSO(VSX) assessment in 2022: status flag = confirmed variable;

variability classification symbol = “SR”)

¶ the nearest of the M giants, radius  $> 0.5$  au; evolutionary status uncertain (is core He fusion now finished?)

¶ cause of the observed Ba overabundance is unknown (undetected evolved companion?)

¶  $\gamma$  Cru B (celestial-sphere neighbour through mere line-of-sight coincidence) is mag. 6.4, with AB astrometry  $93'' \rightarrow 133''$ , PA  $41^\circ \rightarrow 24^\circ$ , 1826  $\rightarrow$  2018;  $\gamma$  Cru C is mag. 9.7, with AC astrometry  $155'' \rightarrow 168''$ , PA  $86^\circ \rightarrow 68^\circ$ , 1879  $\rightarrow$  2018 slight var.? (2.60–2.66 in V passband?)

AAVSO(VSX), as of 2022 Jul. 30, notes existence of NSV entry and availability of 22 AAVSO observations (AASVO status flag = suspected variable; no variability classification symbol); further photometric study advisable? Kraz

$\delta$	Crv A <sup>†</sup>	12 31.1–16 39	2.95 <sup>†</sup> –0.05 <sup>†</sup>	B9.5 IVn	$\sim 37.6$	0.8	87	0.252	237	+9 V
$\gamma$	Cru A	12 32.5–57 15	1.62 <sup>†</sup> 1.60	M3.5 III <sup>†</sup>	37	–0.6	89	0.267	174	+21
$\beta$	Crv	12 35.6–23 32	2.65 <sup>†</sup> 0.90	G5 II <sup>†</sup>	22	–0.6	146	0.057	179	–8 V



$\alpha$	Mus Aa	12 38.6–69 16	2.69 <sup>†</sup> –0.22	B2 IV–V	10.3	–2.2	320	0.042	252	+13 V	<p>¶ slow rotator (but <math>\leq 180</math> d)</p> <p>¶ possibly in evolutionary transition (He core about to ignite?)</p> <p>¶ assertion of weak Ba-star status is perhaps erroneous</p> <p>¶ Astron. Alm. (epoch 2021.5) assigns MK type G5 IIb</p> <p>slight var., <math>\beta</math> Cep type, 2.68–2.73 in V, 2.17 h classification of the low-amplitude variability as <math>\beta</math> Cep, accepted by AAVSO(VSX), has been questioned elsewhere (AAVSO(VSX) assessment in 2022: status flag = confirmed variable; variability classification symbol = “BCEP”)</p> <p>¶ rapid rotator (<math>&lt; 2</math> d)</p> <p>¶ we give mag. for combined light of <math>\alpha</math> Mus Aa and its very close, less luminous, companion <math>\alpha</math> Mus Ab: the pair is sparsely observed, perhaps with nothing beyond the <a href="https://doi.org/10.1093/mnras/stt1694">2013MNRAS.436.1694R</a> report that binarity has been discovered (Sydney interferometer: two 2020 observations, with angular separations <math>\sim 10</math> mas, <math>\sim 16</math> mas); individual mags. are 2.8, 5.5</p>
$\gamma$	Cen $\underline{A}$ <sup>†</sup>	12 42.8–49 05	2.82 <sup>†</sup> $\sim$ –0.01 <sup>†</sup>	A1 IV	25	–0.1	130	$\sim 0.194$	$\sim 267$	–6	orbit 84 y; 0.4" (2010),
$\gamma$	Cen $\underline{B}$ <sup>†</sup>	12 42.8–49 05	2.88 <sup>†</sup> $\sim$ –0.01 <sup>†</sup>	A0 IV	25	–0.2	130	$\sim 0.194$	$\sim 267$	–6	0.3" (2019); max = 1.7"; AB distance 8 au min, 67 au max, 37 au average; $\gamma$ Cen D (mag. 3.85), despite its large physical distance from, and large angular separation from, the $\gamma$ Cen AB pair (1.72 ly, $\sim 1^\circ$ ) is likely gravitationally bound to $\gamma$ Cen AB, experiencing that binary as essentially a point mass; $\gamma$ Cen D is a.k.a. $\tau$ Cen
$\gamma$	Vir $\underline{AB}$ <sup>†</sup>	12 42.9 –1 35	2.74 <sup>†</sup> 0.36 <sup>†</sup>	F1 V + F0mF2 V	85	2.4	39	$\sim 0.619$	$\sim 276$	–20	<p>¶ GCPD gives <math>m_v</math>, B–V values only for <math>\gamma</math> Cen AB combined light (as 2.16, –0.01); we take individual <math>\gamma</math> Cen A, <math>\gamma</math> Cen B <math>m_v</math> values from WDS, as viewed 2022 Aug. 05</p> <p>¶ <math>\gamma</math> Cen system presents slight variability, V-passband range 0.012 (<i>TESS</i> mission), period 2.401 d, considered by AAVSO(VSX) to be possibly of type “<math>\alpha^2</math> CVn” (<math>\alpha</math> CVn A) (AAVSO(VSX) assessment in 2022: status flag = confirmed variable, variability classification symbol = “ACV:”)</p> <p>¶ Arabic name Muhlifain, for <math>\gamma</math> Cen A, is not currently IAU-official</p> <p>A: 3.48; B: 3.53; 0.8" (2007); 3.0" (2020) Porrima orbit 169 y; separation 5 au min (most recently 1836 and 2005), 81 au max, 43 au average, with plane of orbit inclined <math>31^\circ</math> to plane of sky; for discussion of orbit, with observations plot showing error bars (binary astrometry being now old enough to archive data for one full orbit), cf Kaler at <a href="https://stars.astro.illinois.edu/sow/porrima.html">stars.astro.illinois.edu/sow/porrima.html</a>;</p> <p>the <math>\gamma</math> Vir AB pair has, in addition to some faint celestial-sphere neighbours, the widely separated, but not very faint, neighbours <math>\gamma</math> Vir E (mag. 8.9) and <math>\gamma</math> Vir F (mag. 9.5), neither of which is gravitationally bound to the AB pair; AE astrometry is 255" <math>\rightarrow</math> 260", PA 186<math>^\circ</math> <math>\rightarrow</math> 167<math>^\circ</math>, 1851 <math>\rightarrow</math> 2016, and AF astrometry is 482" <math>\rightarrow</math> 422", PA 269<math>^\circ</math> <math>\rightarrow</math> 267<math>^\circ</math>, 1909 <math>\rightarrow</math> 2015</p> <p>¶ <math>m_v</math>, B–V values are for <math>\gamma</math> Vir AB combined light; although GCPD does not give values for <math>\gamma</math> Vir A, GCPD does additionally give 2.80, 0.32 for <math>\gamma</math> Vir B</p> <p>¶ AAVSO(VSX), while noting existence of NSV entry, asserts constant light</p> <p>(AAVSO(VSX) assessment in 2022: status flag = confirmed non-variable; variability classification symbol = “CST”)</p> <p>¶ lunar occultations possible, planetary occultations possible-yet-rare</p>
$\beta$	Mus $\underline{Aa}$	12 47.7–68 14	3.04 <sup>†</sup> –0.19 <sup>†</sup>	B2 V <sup>†</sup>	$\sim 9.6$	–2.1	340	$\sim 0.043$ <sup>†</sup>	$\sim 258$	+42 <sup>†</sup> V	<p>A: 3.52; B: 3.98, 1.0", PA 317<math>^\circ</math> <math>\rightarrow</math> 58<math>^\circ</math>, 1880 <math>\rightarrow</math> 2019 orbit 194 y; average AB distance uncertain (101 au, or only <math>\sim 80</math> au?); orbit map, showing error bars, given by Kaler at <a href="https://stars.astro.illinois.edu/sow/betamus.html">stars.astro.illinois.edu/sow/betamus.html</a>, with Kaler’s accompanying discussion of orbit-modelling problems, underscores limitations in current <math>\beta</math> Mus AB knowledge;</p>

$\beta$	Cru A	12 49.1–59 49	1.25 <sup>†</sup> –0.24	B0.5 III <sup>†</sup>	12	–3.4 300	0.046 249 +16 SB <sup>†</sup>	<p><a href="#">2013MNRAS.436.1694R</a> reports splitting <math>\beta</math> Mus A as tight binary system <math>\beta</math> Mus Aa, <math>\beta</math> Mus Ab, in two 2010 observations with Sydney interferometer: Ab is mag. 6.6, and Aa,Ab angular separation is &lt; 50 mas</p> <p>¶ <math>m_v</math>, B–V values are for <math>\beta</math> Mus Aa,Ab,B combined light</p> <p>¶ AAVSO(VSX) situation in 2022: no status flag (so not confirmed variable, not suspected variable, not confirmed non-variable), and a fortiori no variability classification symbol</p> <p>¶ <math>\beta</math> Mus A is rapid rotator (&lt; 1 d)</p> <p>¶ <math>\beta</math> Mus B is of MK type B2.5 V</p> <p>¶ a runaway system, in the sense of presenting a high velocity relative to the general galactic rotation</p> <p>A is slight var., <math>\beta</math> Cep type, 1.23–1.31 in V Mimosa although the photometry of <math>\beta</math> Cru A is dominated by a period of 5.68 h, this star is multiperiodic (AAVSO(VSX) assessment in 2022: status flag = confirmed variable; variability classification symbol = “BCEP”)</p> <p>¶ SB period 1828.0 d, <math>e=0.38</math> (with distance between the SB components 5.4 au min, 12.0 au max); Kaler at <a href="#">stars.astro.illinois.edu/sow/mimosa.html</a> discusses other possible companions, including an X-ray visible, and yet optically invisible, object interpreted as a pre-MS star; in gross optical-astronomy terms, the unresolved <math>\beta</math> Cru A SB has celestial-sphere neighbours B,C,D,E, of which only C is brighter than mag. 10; AC astrometry is 384″→373″, PA 23°→23°, 1826→2000</p> <p>¶ <math>\beta</math> Cru A is believed to be a rapid rotator (possibly ~3.6 d)</p> <p>¶ <math>\beta</math> Cru A, its MK luminosity class “III” notwithstanding, is only about halfway through its career of stable-core hydrogen fusing</p>
$\epsilon$	UMa A	12 55.1+55 50	1.76 <sup>†</sup> –0.02	A0p IV: (CrEu)	~39.5	–0.3 83	0.112 94 –9 SB <sup>†</sup>	<p>slight var., <math>\alpha^2</math> CVn type, 1.75–1.78 in V, 5.09 d <b>Alioth</b> (AAVSO(VSX) assessment in 2022: status flag = confirmed variable; variability classification symbol = “ACV”) the brightest of the Ap stars (in the specific case of <math>\epsilon</math> UMa A, the magnetic-dipole axis is believed to be nearly perpendicular to rotation axis, yielding Cr bands nearly perpendicular to equator; dipole strength is unusually low) (but it has also been suggested that a substellar companion of mass ~14.7× Jupiter, at average inter-component distance 0.055 au, orbit 5.1 d, rather than a 5.1-d stellar rotation, is the source of the observed variability period); WDS, as viewed 2021 Nov. 17, documents just 2 astrometry measurements for the <math>\epsilon</math> UMa B companion of <math>\epsilon</math> UMa A; the discovery of <math>\epsilon</math> UMa B, via speckle interferometry, is announced in <a href="#">1978MNRAS.183.701M</a>; we at the Handbook do not know whether this discovery of <math>\epsilon</math> UMa B constitutes a resolving of the putative SB</p>
$\delta$	Vir A	12 56.8 +3 16	3.38 <sup>†</sup> 1.57	M3 III <sup>†</sup>	16	–0.5–198	0.473 <sup>†</sup> 264 –18 <sup>†</sup> V?	<p>slight var., semiregular, 3.32–3.40 in V Minelauva (AAVSO(VSX) assessment in 2022: status flag = confirmed variable; variability classification symbol = “SR”); multiperiod pulsator</p> <p>¶ high space velocity relative to galactic neighbours</p> <p>¶ evolutionary status uncertain (helium fusion recently started, or already finished?)</p>
$\alpha$	CVn A <sup>†</sup>	12 57.1+38 12	2.89 <sup>†</sup> –0.12	A0 Vp (SiEu)	28	0.1 110	0.241 283 –3 V SB	<p>¶ Astron. Alm. (epoch 2021.5) assigns MK type M3+ III B:5.5, F0 V, 19.5″, PA 234°→230°, 1777→2018 Cor Caroli orbit <math>\geq 8300</math> y (common proper motion indicates true binarity); separation <math>\geq 675</math> au; prototype for the <math>\alpha^2</math> CVn var. type; rotation period is 5.46939 d, with consequent spot-driven slight V-mag. range 2.86–2.93 (AAVSO(VSX) assessment in 2022: status flag = confirmed variable; variability classification symbol = “ACV”); the <math>\alpha</math> CVn A SB is not as yet resolved, even in interferometry (so WDS cannot as yet write “<math>\alpha</math> CVn Aa,” “<math>\alpha</math> CVn Ab”), and also <math>\alpha</math> CVn B is an as-yet-unresolved SB (so WDS cannot as yet write “<math>\alpha</math> CVn Ba,” “<math>\alpha</math> CVn Bb”)</p>

$\epsilon$	Vir A	13 03.3+10 50	2.83 <sup>†</sup>	0.93	G9 IIIab <sup>†</sup>	29.8	0.2	110	0.275	274	-14 V?	<p>¶ two correct, potentially confusing, designations are <math>\alpha</math> CVn A (signalling that this is the brighter of the binary pair) and <math>\alpha^2</math> CVn (signalling that <math>\alpha^1</math> crosses the local meridian before <math>\alpha^2</math>, lying further W); the Latin “heart-of-Charles” designation, official at IAU as of 2016, honours the “martyr king” Charles I (although Charles II is sometimes cited in error)</p> <p>possible var.? (2.65–2.84 in V passband?) Vindemiatrix AAVSO(VSX), as of 2022 Jul. 30, notes existence of NSV entry, notes absence of record of AAVSO observations, cites Smyth as discoverer (AAVSO(VSX) assessment: status flag = suspected variable; no variability classification symbol); further photometric study advisable?</p> <p>¶ one of the most notable X-ray sources in our table (X-ray luminosity, although far below <math>\alpha</math> Aur, is nevertheless almost 300<math>\times</math> solar)</p> <p>¶ Astron. Alm. (epoch 2021.5) assigns MK type G8 IIIab possible slight variability? (2.94–3.02 in V passband?) AAVSO(VSX), as of 2022 Jul. 30, notes existence of NSV entry and availability of one AAVSO observation (AAVSO(VSX) assessment: status flag = suspected variable; no variability classification symbol); further photometric study advisable?</p> <p>¶ slow rotator (but <math>\leq</math> 240 d)</p> <p>¶ evolutionary state uncertain (core-helium fusion impending, or already in progress?)</p>
$\gamma$	Hya A	13 20.2–23 18	2.99 <sup>†</sup>	0.92	G8 IIIa	-24.4	-0.1	134	0.081	121	-5 V?	<p>rapid rotator (&lt; 2d)</p> <p>¶ low metallicity</p> <p>¶ debris disk (unusually luminous, given evolutionary state of <math>\gamma</math> Cen)</p> <p>¶ AAVSO(VSX) situation in 2022: no status symbol (so not confirmed variable, not suspected variable, not confirmed non-variable), and a fortiori no variability classification symbol</p>
$\iota$	Cen	13 21.9–36 50	2.75	0.04	A2 Va <sup>†</sup>	55	1.5	59	0.352	256	0	<p>B:3.88, A1mA7 IV–V, 15”; period &gt;5000 y? Mizar (more precisely 13.9”<math>\rightarrow</math>14.6”, PA 143<math>^\circ</math><math>\rightarrow</math>153<math>^\circ</math>, 1755<math>\rightarrow</math>2020); not only is <math>\zeta</math> UMa AB a true binary; it is now additionally argued (controversy possibly continues) that the pair <math>\zeta</math> UMa Ca (Alcor), Cb is gravitationally bound to the pair AB (Bob King, <i>Sky and Telescope</i> 2015 March 25); <math>\zeta</math> UMa Aa, <math>\zeta</math> UMa Ab are an interferometrically resolved SB2, seen nearly edge-on, with at least three orbital solutions published (20.5385 d or 20.5386 d, <math>e=5.54</math> or 5.53, angle subtended by semimajor axis as projected onto plane of sky <math>\approx</math>10 mas (<a href="http://apod.nasa.gov/apod/ap970219.html">apod.nasa.gov/apod/ap970219.html</a>, as NASA “Astronomy Picture of the Day” for 1997 Feb. 19, depicts an NPOI-derived <math>\zeta</math> UMa Aa,Ab orbit), with <a href="https://doi.org/10.1002/newa.15324">2010NewA...15..324G</a> suggesting, on the basis of astrometry perturbations, a possible further unseen body); <math>\zeta</math> UMa B is an unresolved SB, period 175.6 d, with highly elliptical orbit; although the old, widely repeated claim (cf Heard <i>ApJ</i> 1949) that <math>\zeta</math> UMa C is binary is shown in <a href="http://www.leosondra.cz/en/mizar/">www.leosondra.cz/en/mizar/</a> to be unfounded, binarity is now established (with WDS accordingly writing “<math>\zeta</math> UMa Ca,” “<math>\zeta</math> UMa Cb,” at mags. 4.0 and 8.0 respectively, typical angular separation 1”, with 5 satisfactory astrometry measurements over the period 2007<math>\rightarrow</math>2009); Cb is a mid-M red dwarf, very notable as one of the few cases of a red dwarf detected as gravitationally bound to an A star (Ca is of MK type A5 Vn); the IAU-official name “Alcor” applies to <math>\zeta</math> UMa Ca; <a href="http://www.leosondra.cz/en/mizar/">www.leosondra.cz/en/mizar/</a> should be consulted also (a) for details on <math>\zeta</math> UMa multiplicity-studies history, including Galileo and Michelson (Leos Ondra, citing inter alia Fedele 1949, seems to establish that it was Galileo’s pupil Castelli, rather than (as widely asserted) Riccioli, who discovered Mizar’s visual duplicity) and (b) for a 15’ map documenting around 20 of the stars in the field, including mag. 7.6 “Stella Ludoviciana” (“Sidus Ludovicianum,” in WDS <math>\zeta</math> UMa D), a mere line-of-sight coincidence on the celestial sphere, too distant from <math>\zeta</math> UMa ABC to</p>
$\zeta$	UMa <u>Aa</u> <sup>†</sup>	13 24.9+54 48	2.05 <sup>†</sup>	0.03 <sup>†</sup>	A1 Va <sup>†</sup>	40	0.1	90	0.123	100	-6 SB2 <sup>†</sup>	

be gravitationally bound to this 6-star system (Aa,Ab; B binary-as-yet-unresolved, Ca,Cb): WDS additionally documents, as gravitationally bound to the ζ UMa ABC 6-star system, E (mag. 6.9), F (mag. 9.9), G (mag. 8.2), and H (mag. 8.6)

¶  $m_v$ , B-V values are for ζ UMa Aa, Ab,B combined light

¶ Astron. Alm. (epoch 2021.5) assigns MK type “A1 Va” (Si)”

¶ possible variability now discounted (AAVSO(VSX) situation for the ζ UMa system in 2022: status symbol = confirmed non-variable; no variability symbol assigned (but should this not now be assigned the symbol “CST”?)

¶ in the mythology of the Mi’kmaq and the St Lawrence Seaway Iroquois, as presented at [www.aavso.org/myths-uma](http://www.aavso.org/myths-uma), α, β, γ, and δ UMa are a bear at various seasons of the year passant, rampant, and expired (its four paws upward in death), pursued in the warm months by seven hunters, but once the nights are cold by a remaining above-horizon three, these persistent three being ε UMa, Mizar, and ζ UMa, with Alcor the middle hunter’s cooking pot, awaiting bear meat; [www.aavso.org/myths-uma](http://www.aavso.org/myths-uma) gives further detail, offering also a speculation about a possible bear mythology shared by Siberian and North American Paleolithic peoples, in the epoch of the Bering Strait land bridge

slight ellipsoid var., 0.96–1.00 in V, 4.0145 d **Spica** (AAVSO(VSX) assessment in 2022: status flag = confirmed variable; variability classification symbol = “ELL+BCEP”); very tight (< 1”) system resolved interferometrically (and in occultation?) into α Vir Aa (mag. 1.3), α Vir Ab (mag. 4.5), α Vir Ac (mag. 7.5); as SB2 (primary-to-secondary distance 0.12 au; the geometry is close to achieving a grazing eclipse), the brightest of the rotating-ellipsoid variable systems; the Aa,Ab orbit is highly eccentric; Aa was measured in 1975 to lie 0.50” from Ac ; although the Aa,Ab pair is at all times very close, an angular distance of 0. 1” is reported from 1975; Aa (a rapid rotator, at ~0.3 breakup speed) is itself a pulsating variable of the β Cep type (0.1738 d; shortly after the ~1970 discovery of the β Cep variability, photometric and spectroscopic variations were present; the photospheric variations soon ceased, but the spectroscopic (radial-velocity, i.e. pulsational) variations continued; [2016MNRAS.458.1964T](https://doi.org/10.1093/mnras/271.4.1964), incorporating precision *MOST* photometry, reports for Aa one radial and two non-radial pulsation modes, with one of the non-radial modes tidally induced) ¶ in an early application of intensity interferometry, [1971MNRAS.151..161H](https://doi.org/10.1093/mnras/151.1.161H) argues with the example of α Vir Aa,Ab that given supporting spectroscopy and photometry, orbit and distance of a double-lined SB can be deduced (the SB distance notably without recourse to trigonometric parallax, since distance can be deduced by comparing the angular and the physical dimensions of the ascertained orbit) ¶ the tidal-interactions studies [2016A&A...590A..54H](https://doi.org/10.1093/mnras/271.4.1964) and [2013A&A...556A..49P](https://doi.org/10.1093/mnras/271.4.1964) stress the importance of the α Vir Aa,Ab double-lined SB for critically testing the (astrophysically foundational) assumption that the individual components *x*, *y* of a binary,

α Vir Aa<sup>†</sup> 13 26.4–11 17 0.98<sup>†</sup> –0.24 B1 V<sup>†</sup> 13 –3.4 250 0.052 234 +1 SB2<sup>†</sup>

of determined masses, rotation periods, and chemical compositions, resemble in their photospheres, and even in their interiors, solitary stars  $x'$ ,  $y'$  possessing the same masses, rotation periods, and chemical compositions (could tidal effects, e.g. change internal temperature structure?); additionally, the tidal effects in the  $\alpha$  Vir Aa,Ab SB are judged in [2009ApJ...704..813H](#) to be responsible for large-scale shearing horizontal photospheric motions, spectroscopically observable as modifiers of line profiles (but [2016MNRAS.458.1964T](#) questions the judgement)

¶ assignment of individual MK types to Aa, Ab is challenging: the rather-unevolved-B MK types ([1971MNRAS.151..161H](#) B1.5 IV-V + B3V, [2007AAS...211.6301A](#) B0.5 III-IV + B2.5-B3V) are in any case consistent with rather high masses (10.9  $M_{\odot}$  + 6.8  $M_{\odot}$ , 10.25  $M_{\odot}$  + 6.97  $M_{\odot}$ , for these two respective papers)

¶ as is to be expected from the failure of Aa,Ab to be tidally locked, the system is young (with [2016MNRAS.458.1964T](#) assigning as age  $12.5 \pm 1$  My)

¶ the Aa,Ab binary is a polarimetric variable (ISM material entrained?), and a strong X-ray source (colliding winds?)

¶  $\alpha$  Vir Ab is one of the few stars known to exhibit Struve-Sahade variation ([en.wikipedia.org/wiki/Struve%E2%80%93Sahade\\_effect](http://en.wikipedia.org/wiki/Struve%E2%80%93Sahade_effect)) in its spectral line strengths

¶ [1972BAA...82..431K](#) describes the 18.6-year 1940-through-2050 cycle of lunar occultation possibilities

¶  $E(B-V)=+0.03$

¶ a good marker of celestial equator Heze (precession placed  $\zeta$  Vir exactly onto equator in Feb. 1883)

¶ rapid rotator ( $< 0.5$  d; this renders puzzling the possible evidence for chemical anomalies, which would presuppose a quiet atmosphere)

¶ slight variability, of  $\delta$  Set type, range 0.009 in V (TESS mission; AAVSO(VSX) assessment in 2022: status flag = confirmed variable; variability classification symbol = “DSCT”)

¶ Astron. Alm. (epoch 2021.5) assigns MK type A2 IV<sup>-</sup>

¶ elusive red-dwarf companion  $\zeta$  Vir B is mag. 10.0, with MK classification suggested as M4 V – M7 V; WDS documents just 9  $\zeta$  Vir AB measurements, 1.8" → 1.8", PA 145° → 154°, 2004 → 2010; the discovery paper is [2010ApJ...712..421H](#), establishing inter alia shared proper motion, and proceeding from stellar coronagraphy on adaptive-optics platforms, both at Palomar and at Hawai'i-Haleakala (rather than, as is more usual with a difficult binary, from interferometry [with  $\zeta$  Vir B directly imaged in Figs. 1 and 4]); [2010ApJ...712..421H](#), while remarking that orbital coverage is as yet too brief for an orbital solution to be attempted, nevertheless (assuming 2.04  $M_{\odot}$  for  $\zeta$  Vir A, 0.168  $M_{\odot}$  for  $\zeta$  Vir B) computes approximate lower bounds for semimajor axis, for  $e$ , and for period  $a_p$ , respectively, 24.9 au, 0.16, and 124 y; [2010ApJ...712..421H](#) is a contribution to the important, and until recently unstudied, topic of low-mass companions for A stars (another contribution to this topic is, however, the circa-2010 discovery of an elusive M-type companion for the A5 Vn star that is Alcor);

$\zeta$  Vir A 13 35.9 -0 43 3.37<sup>†</sup> 0.11 A2 IV<sup>†</sup> 44 1.6 74 0.285 280 -13

$\epsilon$	Cen Aa	13 41.4–53 35	2.30 <sup>†</sup> –0.23	B1 III <sup>†</sup>	8	–3.3	400	0.019	233	+3	<p>the topic in its turn is a building block in the overall theory of star and exoplanet genesis (could massive (e.g. A-type) stars acquire low-mass companion stars via condensation not directly from the parent molecular cloud, but rather from condensation in an unstable circumstellar disk?), the <a href="#">2010ApJ...712..421H</a> detection of red-dwarf <math>\zeta</math> Vir B additionally affords an explanation for the puzzling <math>\zeta</math> Vir X-ray emission observed by <i>ROSAT</i> (as a star of a spectral type lacking strong winds and lacking convection up to the photosphere, <math>\zeta</math> Vir A would not itself be expected to emit X-rays: similar puzzles of X-ray emission from the putatively X-ray-dark A stars arise elsewhere also, and perhaps are similarly to be solved in terms of X-ray emission from (elusive) red-dwarf companions)</p> <p>slight var., <math>\beta</math> Cep type, 2.29–2.31 in V multiperiodic, with primary period 0.169608 d (AAVSO(VSX) assessment in 2022: status flag = confirmed variable; variability classification symbol = “<b>BCEP</b>”)</p> <p>¶ rapid rotator (&lt; 2.7 d)</p> <p>¶ metals underabundant</p> <p>¶ although we here assign MK luminosity class “III,” Kaler at <a href="#">stars.astro.illinois.edu/sow/epsceen.html</a> discusses uncertainty</p> <p>¶ we state mag. for combined light; WDS documents just one single measurement, from interferometry, for the elusive <math>\epsilon</math> Cen Ab (mag 4.90, 0.2” from Aa)</p>
$\eta$	UMa	13 48.5+49 12	1.86 <sup>†</sup> –0.19 <sup>†</sup>	B3 V <sup>†</sup>	31	–0.7	104	0.122	263	–11 SB?	<p><b>Alkaid</b></p> <p>resembles <math>\alpha</math> UMa, at the other extreme of the Big Dipper, in not belonging to UMa Moving Group; <a href="#">1921LicOB...10..110T</a> asserts membership in what was at that time called the “Pleiades Group”</p> <p>¶ rapid rotator (&lt; 21 h), with some line variability (circumstellar ejecta disk?)</p> <p>¶ slight variability, 0.01 mag. in V (TESS mission), of the slowly pulsating B-star type, also known as the “53 Per type” (AAVSO(VSX) assessment in 2022: status flag = confirmed variable; variability classification symbol = “<b>SPB</b>”)</p> <p>¶ X-ray source</p> <p>¶ colour and mean temperature are anomalous for the MK type</p> <p>¶ unusually young in our Sample S (&lt; 15 My)</p> <p>¶ E(B–V)=+0.02</p>
$\nu$	Cen	13 50.9–41 48	3.41 <sup>†</sup> –0.23	B2 IV <sup>†</sup>	~7.5	–2.2	440	0.034	233	+9 SB <sup>†</sup>	<p>SB period is 2.622 d;</p> <p>slight variable, not eclipsing, but varying photometrically through a so-called “reflection” (irradiation, re-radiation) effect (AAVSO(VSX) assessment in 2022: status flag = confirmed variable; variability classification symbol = “<b>R</b>”); additionally, the primary has been said, but not by AAVSO(VSX), to be a pulsator in the <math>\beta</math> Cep class; observed mag. range is 3.40–3.42 in V</p> <p>¶ MK luminosity class “IV” notwithstanding, primary is still a stable fuser of core hydrogen</p> <p>¶ possible weak instance of “Be phenomenon” (with the outbursts possibly temporary)</p>
$\mu$	Cen Aa	13 51.0–42 35	3.34 <sup>v†</sup> –0.17 <sup>†</sup>	B2 IV–V pne <sup>†</sup>	~6.4	–2.5	510	0.031	232	+9 SB	<p>variable of <math>\gamma</math> Cas type.; 2.92–3.47 in V (AAVSO(VSX) assessment in 2022: status flag = confirmed variable; variability classification symbol = “<b>GCAS</b>”); rapid rotator, and (consistently with the <math>\gamma</math> Cas behaviour) an instance of the “Be phenomenon”; additionally said to be a multiperiodic non-radial pulsator; BSC5: “line profiles of MgII 4481 change in period 0.505 d, about five times the period of weaker absorption”; variable H<math>\alpha</math>; “variable line profiles”; short-term photometric and polarimetric variability has also been reported (cf p. 46 of <a href="#">2013A&amp;ARv..21...69R</a>, which notes a rapid rise, over just a few days, in photometric brightness or line-emission intensity, with a subsequent slower decline)</p> <p>¶ WDS documents just one single measurement for <math>\mu</math> Cen Aa,Ab, via 2010 Sydney interferometry (as Aa mag. 3.50,</p>

$\eta$	Boo A	13 55.8+18 17	2.68 <sup>†</sup>	0.58	G0 IV <sup>†</sup>	88	2.4	37	0.361	190	0 SB	<p>Ab mag. 6.70, with angular separation 0.1")</p> <p>¶ <math>m_v</math>, B–V values are for <math>\mu</math> Cen Aa,Ab combined light</p> <p>¶ Astron. Alm. (epoch 2021.5) assigns MK type "B2 IV–Vpne (shell)"</p> <p>possible variability (0.1 in V passband?) Muphrid</p> <p>AAVSO(VSX), as of 2022 Jul. 30, notes existence of NSV entry, fails to find record of AAVSO observations (AAVSO(VSX) assessment: status flag = suspected variable; no variability classification symbol); further photometric study advisable?</p> <p>¶ unusually metal-rich</p> <p>¶ an X-ray source (hot corona)</p> <p>¶ <a href="#">2007ApJ...657.1058V</a> discusses recent work (MOST helioseismology, PTI interferometry)</p> <p>¶ <math>\eta</math> Boo B is mag. 9.99; discrepant proper motions for the AB pair (126"→115", PA 119°→85°, 1822→2020) establish that their pairing is a mere line-of-sight coincidence</p>
$\zeta$	Cen	13 57.0–47 24	2.54	–0.23	B2.5 IV <sup>†</sup>	8.5	–2.8	380	0.073	232	+7 SB2	<p>slight var. (2.52–2.55 in V passband, 2.29 d) AAVSO(VSX), as of 2022 Jul. 30, notes that no AAVSO observations are available, gives period 2.2903 d (AAVSO(VSX) assessment: status flag = confirmed variable; variability classification symbol = "VAR"); further photometric study advisable?</p> <p>¶ primary is a rapid rotator (&lt;1.5 d) (BSC5: "expanding circumstellar disk," and yet not (as viewed 2022 March 3) catalogued as an instance of the "Be phenomenon" in Paris-Meudon BeSS database)</p> <p>¶ MK luminosity class "IV" notwithstanding, primary is possibly only halfway through its core hydrogen fusing</p> <p>¶ SB period 8.02 d; SB as yet unresolved, even by interferometry (so WDS not yet able to write "Cen A," "Cen B")</p>
$\beta$	Cen <u>Aa,Ab</u>	14 05.5–60 29	0.61 <sup>†</sup>	–0.24 <sup>†</sup>	B1 III + B1 III	9.0 <sup>†</sup>	–4.8	360 <sup>†</sup>	0.041	235	+6 SB2 <sup>†</sup>	<p>¶ E(B–V) = –0.02</p> <p>B:3.94, A1mA7 IV–V, 0.3" (2019) <b>Hadar</b></p> <p>(more fully: 1.1"→0.3", PA 257°→168°, 1935→2019); AB orbit is already constrained by the existing observations, with period 125–220 y, and <a href="#">2016A&amp;A...588A..55P</a> indicates that it should be possible to compute the orbit by ~2025 or ~2030 or so); the <math>\beta</math> Cen A system (reported in <a href="#">1999MNRAS.302..245R</a> as resolved at AAT through spectrally dispersed aperture-masking interferometry) is <math>\beta</math> Cen Aa, <math>\beta</math> Cen Ab, comprising a pair of fast rotators of nearly equal mass, and with oddly disparate (high) rotation speeds (the slower rotator is known to be magnetic, so magnetic braking is possible) and with orbit so eccentric (how can the molecular-cloud ISM condensation have allowed this to happen?) as to make the periastron tight (at &lt; 10 R*; so could there be tidal interaction between Aa, Ab at periastron, perturbing the variability that we discuss below?); Aa,Ab is additionally reported in <a href="#">1999MNRAS.302..245R</a>, on the strength of ESO spectra, to be not just SB, but SB2; since the masses are nearly equal, it becomes a delicate question which to take as the primary, i.e. to which to apply the label "Aa" and with it the IAU-official name "Hadar"; this question is answered by WDS in its usual terms, with "Aa" deemed to be the (very slightly) more luminous, more massive star (Aa mag. 1.29, Ab mag. 1.44), and yet the contrary decision has also been taken in the literature, since it is the less massive star that has the clearer, because the less severely (rotationally) broadened, spectrum; the observational challenges notwithstanding, the observational record for Aa,Ab is favourable, with WDS now documenting 53 measurements for 1995→2018 (not only aperture-masking interferometry, but also speckle interferometry has been done); the Aa,Ab orbit is 357 d, with <math>e=0.8</math>; <a href="#">2002A&amp;A...384..209A</a> finds both <math>\beta</math> Cen Aa and <math>\beta</math> Cen Ab to be <math>\beta</math> Cen variables (AAVSO(VSX), as of 2022 Jul. 14, gives range 0.045 in V for the system, salient period 3.8 h</p>

(AAVSO(VSX) assessment:  
status flag = confirmed variable;  
variability classification  
symbol = “**BCEP**”; the Aa,Ab  
system is of astrophysical significance, as one  
of the rare cases of  $\beta$  Cep variability amenable  
to good-precision mass studies (and indeed  
[2016A&A...588A..55P](#), reporting precision two-filter  
photometry with the *BRITTE* constellation, discusses  
prospects for future asteroseismology, noting also  
that in addition to  $\beta$  Cep (“pressure-wave”) pulsation,  
there is SPB-type (“gravity-wave,” i.e. buoyancy-driven)  
pulsation (the question that of Aa,Ab is presenting which  
of the 17 detected pulsation modes is, however, difficult,  
and the magnetic field of Ab is a further complication [with  
the overall topic of asteroseismology for magnetic  $\beta$  Cep  
variables only sparsely explored, at any rate as of ~2016])  
¶  $m_v$ , B–V values are for  $\beta$  Cen Aa,Ab,B combined light  
¶ we take  $\pi$  and D not from *HIPPARCOS* but from  
[2016A&A...588A..55P](#)  
¶ E(B–V) = +0.02  
¶ the traditional (Latin-derived?) name “Agena” is not IAU-official

$\pi$  Hya 14 07.7–26 48 3.27 1.12 K2 IIIb<sup>†</sup> ~32.3<sup>†</sup> 0.8 ~101<sup>†</sup> 0.148<sup>†</sup> 163 +27<sup>†</sup> V

negative cyanide ion lines are anomalously weak relative to metal lines, consistent with this star’s anomalously high velocity relative to Sun (suggesting interloper in our own galactic region; however,  $\pi$  Hya is more metal-rich than the celebrated interloper  $\alpha$  Boo [Arcturus])  
¶ Astron. Alm. (epoch 2021.5) assigns MK type K2– III Fe–0.5  
¶ in evolutionary terms, in “Red Clump” of core-He fusers (but uncertain whether recent arrival in clump or longtime denizen)  
¶ AAVSO(VSX) situation in 2022: no status flag (so not confirmed variable, not suspected variable, not confirmed non-variable), and a fortiori no variability classification symbol

$\theta$  Cen A 14 08.1–36 29 2.06 1.01 K0 IIIb<sup>†</sup> 55 0.8 59 0.734<sup>†</sup> 225 +1<sup>†</sup>

**Menkent**  
high velocity with respect to Sun suggests interloper status (and yet metallicity is approximately solar)  
¶ Astron. Alm. (epoch 2021.5) assigns MK type K0– IIIb  
¶ AAVSO(VSX) situation in 2022: no status flag (so not confirmed variable, not suspected variable, not confirmed non-variable), and a fortiori no variability classification symbol

$\alpha$  Boo A 14 16.7+19 04 –0.05 1.23 K1.5 III Fe–0.5<sup>†</sup> 89 –0.3 37 2.279<sup>†</sup> 209 –5<sup>†</sup> V

**Arcturus**  
high space velocity  
a metal-poor interloper (from galactic thick disk? but galaxy-merger scenario has also been suggested), and member of Arcturus Moving Group ([2009IAUS..254..139VV](#))  
¶ a magnetic cycle (< 14 y?) has been detected  
¶ still ascending RGB, with He flash impending? (but a later evolutionary stage has also been suggested)  
¶ publication of  $\alpha$  Boo A line atlas [1968pmas.book....G](#) (R. Griffin) was a major event in postwar spectroscopy  
¶  $\alpha$  Boo A has been studied in recent asteroseismology  
¶ slight variability, of a slow-and-irregular type, range –0.07 to –0.02 in V  
(AAVSO(VSX) assessment in 2022:  
status flag = confirmed variable;  
variability classification symbol = “**LB**”)  
¶ a single 1991 observation of a putative companion has constrained WDS to write “ $\alpha$  Boo A,” “ $\alpha$  Boo B”; however, 1990s assertion of multiplicity was retracted in 1998; independently of this pair of developments, there have been suggestions of a sub-stellar-mass companion at the margin of *HIPPARCOS* detectability  
slight variability, 3.54–3.55 in V passband ( $\beta$  Cep type?) AAVO(VSX), as of 2022 Jul. 30, asserts variability, but finds no record of AAVSO observations  
(AAVSO(VSX) assessment:

$\iota$  Lup 14 20.9–46 10 3.54 –0.18 B2.5 IVn<sup>†</sup> ~9.6 –1.5 340 0.013 249 +22



$\gamma$	Boo Aa <sup>†</sup>	14 33.0+38 12	3.04 <sup>†</sup>	0.19 <sup>†</sup>	A7 IV <sup>+</sup>	37.6	0.9	87	0.190	323	-37 V	<p>status flag = confirmed variable;  variability classification symbol = “<b>BCEP:</b>”);  further photometric study advisable?  ¶ rapid rotator (possibly ~0.9 d), and yet no evidence of circumstellar disk, and in particular no Be-phenomenon spectral features  ¶ the MK luminosity class “IV” notwithstanding, still performing stable core-hydrogen fusion  <b>[THIS STAR ONLY IN ONLINE VERSION OF TABLE]</b>  slight var. (<math>\delta</math> Sct type?), 3.02–3.07 in V, 6.97 h Seginus (AAVSO(VSX) assessment in 2022:  status flag = confirmed variable;  variability classification symbol = “<b>DSCTC:</b>”);  further photometric study advisable?  ¶ <math>m_v</math>, B–V values are for <math>\gamma</math> Boo Aa,Ab combined light  ¶ IR excess (from circumstellar debris, so far unexplained)  ¶ Aa,Ab resolved in speckle interferometry, angular separation 70 mas  var. of <math>\gamma</math> Cas and <math>\lambda</math> Eri types, possibly shell, 2.29–2.47 in V multiperiodic, with a salient period 0.6425 d (AAVSO(VSX) assessment in 2022:  status flag = confirmed variable;  variability classification symbol = “<b>GCAS+LERI:</b>”);  BSC5 says H<math>\alpha</math> variable, H<math>\beta</math> “sometimes bright, sometimes dark and double or multiple”;  consistently with <math>\gamma</math> Cas variability,  a rapid rotator (&lt; 1 d) and an instance of “Be phenomenon”; again consistently with <math>\gamma</math> Cas variability, Astron. Alm. (epoch 2021.5) assigns MK type “B1.5 IVpne (shell)”  AB 5.2” (2019) orbit 79.9y Toliman  min = 2” (1955); max = 22”; PA (2017) 325°;  separation 11.2 au min, 35.6 au max; Kaler at <a href="http://stars.astro.illinois.edu/sow/rigel-kent.html">stars.astro.illinois.edu/sow/rigel-kent.html</a>  has map of apparent AB orbit (note further here that Kaler’s green, violet, and blue denote micrometry, photography, and interferometry, respectively:  as the error bars suggest, the <math>\alpha</math> Cen AB orbit is one of the most precisely known wide binary-system orbits in visual-binary astrometry); since plane of orbit is inclined at 79° to plane of sky, the apparent orbit is more severely elliptical than the true orbit (for which <math>e=0.5</math>); Kaler’s map can accordingly be usefully supplemented with the apparent-versus-true-orbit diagram in <a href="http://en.wikipedia.org/wiki/Alpha_Centauri">en.wikipedia.org/wiki/Alpha_Centauri</a>  ¶ whereas magnetic activity of <math>\alpha</math> Cen A is in steep decline since 2005 (analogue of Maunder Minimum? or, rather, mere regular cycle?), <math>\alpha</math> Cen B shows more magnetic activity than <math>\alpha</math> Cen A does, and its cycle is brief (8.2 y in spot numbers, 16.4 y in magnetics; this is not unlike the Sun, for which the corresponding pair of periods is ~11y, ~22 y); and <a href="https://doi.org/10.1093/mnras/stz315">2005A&amp;A...442...315R</a> reports a flare on <math>\alpha</math> Cen B  ¶ GCPD gives not only <math>m_v</math>, B–V values directly measured for <math>\alpha</math> Cen B, but additionally <math>m_v</math>, B–V values directly measured for <math>\alpha</math> Cen AB combined light: -0.29, 0.72  ¶ AAVSO(VSX) situation in 2022:  no status flag (so not confirmed variable, not suspected variable, not confirmed non-variable), and a fortiori no variability classification symbol  ¶ although 2012 <math>\alpha</math> Cen B exoplanet claim is now discounted, an exoplanet is possible (2019 transit has been suggested; cf also <a href="https://doi.org/10.1093/mnras/stz315">2015MNRAS.450.2043D</a>)  ¶ Einstein-ring event expected with 45% probability in 2028, early in May</p>
$\eta$	Cen	14 37.0–42 16	2.36v <sup>†</sup>	-0.22	B1.5 IV pne <sup>†</sup>	11	-2.5	310	0.048	227	0 SB	<p>var. of <math>\gamma</math> Cas and <math>\lambda</math> Eri types, possibly shell, 2.29–2.47 in V multiperiodic, with a salient period 0.6425 d (AAVSO(VSX) assessment in 2022:  status flag = confirmed variable;  variability classification symbol = “<b>GCAS+LERI:</b>”);  BSC5 says H<math>\alpha</math> variable, H<math>\beta</math> “sometimes bright, sometimes dark and double or multiple”;  consistently with <math>\gamma</math> Cas variability,  a rapid rotator (&lt; 1 d) and an instance of “Be phenomenon”; again consistently with <math>\gamma</math> Cas variability, Astron. Alm. (epoch 2021.5) assigns MK type “B1.5 IVpne (shell)”  AB 5.2” (2019) orbit 79.9y Toliman  min = 2” (1955); max = 22”; PA (2017) 325°;  separation 11.2 au min, 35.6 au max; Kaler at <a href="http://stars.astro.illinois.edu/sow/rigel-kent.html">stars.astro.illinois.edu/sow/rigel-kent.html</a>  has map of apparent AB orbit (note further here that Kaler’s green, violet, and blue denote micrometry, photography, and interferometry, respectively:  as the error bars suggest, the <math>\alpha</math> Cen AB orbit is one of the most precisely known wide binary-system orbits in visual-binary astrometry); since plane of orbit is inclined at 79° to plane of sky, the apparent orbit is more severely elliptical than the true orbit (for which <math>e=0.5</math>); Kaler’s map can accordingly be usefully supplemented with the apparent-versus-true-orbit diagram in <a href="http://en.wikipedia.org/wiki/Alpha_Centauri">en.wikipedia.org/wiki/Alpha_Centauri</a>  ¶ whereas magnetic activity of <math>\alpha</math> Cen A is in steep decline since 2005 (analogue of Maunder Minimum? or, rather, mere regular cycle?), <math>\alpha</math> Cen B shows more magnetic activity than <math>\alpha</math> Cen A does, and its cycle is brief (8.2 y in spot numbers, 16.4 y in magnetics; this is not unlike the Sun, for which the corresponding pair of periods is ~11y, ~22 y); and <a href="https://doi.org/10.1093/mnras/stz315">2005A&amp;A...442...315R</a> reports a flare on <math>\alpha</math> Cen B  ¶ GCPD gives not only <math>m_v</math>, B–V values directly measured for <math>\alpha</math> Cen B, but additionally <math>m_v</math>, B–V values directly measured for <math>\alpha</math> Cen AB combined light: -0.29, 0.72  ¶ AAVSO(VSX) situation in 2022:  no status flag (so not confirmed variable, not suspected variable, not confirmed non-variable), and a fortiori no variability classification symbol  ¶ although 2012 <math>\alpha</math> Cen B exoplanet claim is now discounted, an exoplanet is possible (2019 transit has been suggested; cf also <a href="https://doi.org/10.1093/mnras/stz315">2015MNRAS.450.2043D</a>)  ¶ Einstein-ring event expected with 45% probability in 2028, early in May</p>
$\alpha$	Cen B <sup>†</sup>	14 41.2–60 56	1.36 <sup>†</sup>	0.87 <sup>†</sup>	K1 V <sup>†</sup>	750	5.7	4.3	-3.703	-283	-21 V?	<p>AB 5.2” (2019) orbit 79.9y Toliman  min = 2” (1955); max = 22”; PA (2017) 325°;  separation 11.2 au min, 35.6 au max; Kaler at <a href="http://stars.astro.illinois.edu/sow/rigel-kent.html">stars.astro.illinois.edu/sow/rigel-kent.html</a>  has map of apparent AB orbit (note further here that Kaler’s green, violet, and blue denote micrometry, photography, and interferometry, respectively:  as the error bars suggest, the <math>\alpha</math> Cen AB orbit is one of the most precisely known wide binary-system orbits in visual-binary astrometry); since plane of orbit is inclined at 79° to plane of sky, the apparent orbit is more severely elliptical than the true orbit (for which <math>e=0.5</math>); Kaler’s map can accordingly be usefully supplemented with the apparent-versus-true-orbit diagram in <a href="http://en.wikipedia.org/wiki/Alpha_Centauri">en.wikipedia.org/wiki/Alpha_Centauri</a>  ¶ whereas magnetic activity of <math>\alpha</math> Cen A is in steep decline since 2005 (analogue of Maunder Minimum? or, rather, mere regular cycle?), <math>\alpha</math> Cen B shows more magnetic activity than <math>\alpha</math> Cen A does, and its cycle is brief (8.2 y in spot numbers, 16.4 y in magnetics; this is not unlike the Sun, for which the corresponding pair of periods is ~11y, ~22 y); and <a href="https://doi.org/10.1093/mnras/stz315">2005A&amp;A...442...315R</a> reports a flare on <math>\alpha</math> Cen B  ¶ GCPD gives not only <math>m_v</math>, B–V values directly measured for <math>\alpha</math> Cen B, but additionally <math>m_v</math>, B–V values directly measured for <math>\alpha</math> Cen AB combined light: -0.29, 0.72  ¶ AAVSO(VSX) situation in 2022:  no status flag (so not confirmed variable, not suspected variable, not confirmed non-variable), and a fortiori no variability classification symbol  ¶ although 2012 <math>\alpha</math> Cen B exoplanet claim is now discounted, an exoplanet is possible (2019 transit has been suggested; cf also <a href="https://doi.org/10.1093/mnras/stz315">2015MNRAS.450.2043D</a>)  ¶ Einstein-ring event expected with 45% probability in 2028, early in May</p>
$\alpha$	Cen A+1P <sup>†</sup>	14 41.2–60 56	0.00 <sup>†</sup>	0.65 <sup>†</sup>	G2 V <sup>†</sup>	750	4.4	4.3	-3.710	-277	-22 SB	<p>Ca (Proxima), 12.4, M5e, 2.2° SW of A<b>Rigel Kentaurus</b>  still the closest known object in the population of stars and brown dwarfs, despite intense surveying of entire population over the past 20 or 30 years  ¶ gravitational binding of AB+C was finally established with high probability in <a href="https://doi.org/10.1093/mnras/stz315">2017A&amp;A...598L...7K</a>,  and an orbit is considered to be known</p>

												(~550,000 y; min > 4300 au, max 13,000 au) ¶ Ca is elusive, as one faint object in a sea of faint objects (detection was not achieved until the 1915 work of Innes, with blink comparator); nevertheless, violent flaring has been known to take Ca, briefly, up to the threshold of naked-eye visibility ( <a href="#">2018ApJ...860L..30H</a> reports peak V mag. 6.8 for a superflare of 2016 Mar. 18, of duration ~1 h; the situation in UV and millimetre waves is also extreme, as reported in <a href="#">2021ApJ...911L..25M</a> ) ¶ <a href="#">2016Natur.536..437A</a> announces an approx. Earth-mass exoplanet, $\alpha$ Cen Cb, in the habitable zone of its host $\alpha$ Cen Ca; <a href="#">2018ApJ...860L..30H</a> analyzes the germicidal implications of flaring, finding that in a habitable-zone Earth-like exoplanet atmosphere the ozone UV shield would be destroyed; this paper, however, like some others, leaves open the possibility of life on Cb; <a href="#">breakthroughinitiatives.org/initiative/3</a> advocates nanocraft exploration, and <a href="#">en.wikipedia.org/wiki/2069_Alpha_Centauri_mission</a> has footnote links to reports of small-scale discussions at NASA and in the USA Congress, envisaging the launch of some (nanocraft?) mission to celebrate the Apollo 11 centenary ¶ in 2020, a more distant exoplanet, either a super-Earth or a mini-Neptune, was suggested, as $\alpha$ Cen Ac (orbit 1930 d, whereas Ab has orbit 11 d; an unexpectedly bright detection with the VLT SPHERE instrument has been interpreted as the possible signature of rings around the putative Ac); an exoplanet Cd was announced in 2022 Feb. ¶ GCPD gives not only $m_v$ , B–V values directly measured for $\alpha$ Cen A, but additionally $m_v$ , B–V values directly measured for $\alpha$ Cen AB combined light: –0.29, 0.72 ¶ AAVSO(VSX), as of 2022 Jul. 30, notes existence of NSV entry for $\alpha$ Cen A, fails to find record of AAVSO observations, and provisionally offers the “BY Dra” variability type, with variability range –0.3 to +0.01 in V passband (AAVSO(VSX) assessment: status flag = suspected variable, variability classification symbol = “BY:”); further photometric study advisable?
$\alpha$	Lup A	14 43.5–47 29	2.30 <sup>†</sup> –0.21	B1.5 III	7	–3.5 460	0.032 221	+5 SB	slight var. of $\beta$ Cep type, 2.29–2.34 in V, 6.24 h actually multiperiodic, with primary period (unusually long) 6.23632 h given by AAVSO(VSX) as viewed 2021 Jan. 16, and again 2022 Jul. 14 (AAVSO(VSX) assessment in 2022: status flag = confirmed variable; variability classification symbol = “BCEP”)			
$\alpha$	Cir A <sup>†</sup>	14 44.4–65 05	3.18 <sup>†</sup> 0.24 <sup>†</sup>	A7 Vp (Sr)	60.4	2.1 54.1	0.303 220	+7 SB?	B: 8.5, K5 V, 15.7”, PA 263°→224°, 1826→2016 AB probably true binary, with orbit $\geq 2600$ y ¶ Astron. Alm. (epoch 2021.5) assigns MK temperature type A7p Sr Eu and does not assign an MK luminosity class ¶ $m_v$ , B–V values are for $\alpha$ Cir AB combined light ¶ slight variability of $\alpha$ CVn A (“ $\alpha^2$ CVn”) type, and additionally the brightest (slight) variable of the “rapidly oscillating Ap” type; V mag. range is 3.17–3.19, with period 6.826 min; (AAVSO(VSX) assessment in 2022: status flag = confirmed variable; variability classification symbol = “roAp+ACV”); magnetically an oblique rotator (4.4790 d, with field strength ~500× solar); <a href="#">2009MNRAS.396.1189B</a> discusses the rotation, two notably stable putative equatorial chemical-anomaly regions, and asteroseismology, with history and fresh WIRE+SAAO observations			
$\varepsilon$	Boo A	14 46.0+26 59	2.38 <sup>†</sup> 0.97 <sup>†</sup>	K0 II–III <sup>†</sup>	16 <sup>†</sup>	–1.6 200 <sup>†</sup>	0.044 288	–17 V	B: 4.81, 2.8”, PA 318°→347°, 1822→2018 orbit well over 1000 y ¶ $\varepsilon$ Boo B is of MK type A0 V, and is SB, with at least one component a rapid rotator ¶ $m_v$ , B–V values are for $\varepsilon$ Boo AB combined light; <a href="#">stars.astro.illinois.edu/sow/izar.html</a>			

$\beta$	UMi A+1P <sup>†</sup>	14 50.7+74 04	2.08 <sup>†</sup>	1.47 <sup>†</sup>	K4 III <sup>†</sup>	24.9	-0.9	131	0.035	289	+17 V	discusses difficulties in determination of the individual magnitudes and of the binary system's distance ¶ AAVSO(VSX) situation in 2022: no status flag (so not confirmed variable, not suspected variable, not confirmed non-variable), and a fortiori no variability classification symbol ¶ F.G.W. von Struve: "pulcherrima" ("the loveliest") useful for aligning small equatorial mount <b>Kochab</b> (since NCP, although not quite coincident with $\alpha$ UMi, does lie near the great-circle arc linking $\beta$ UMi with $\alpha$ UMi: <a href="http://arksky.org/Kochab.htm">arksky.org/Kochab.htm</a> [not <a href="http://darksky.org">darksky.org</a> ]) ¶ $m_v$ , B-V values are measured for $\beta$ UMi A, rather than for $\beta$ UMi AB combined light (however, greatly separated (~212")) $\beta$ UMi B would, because shining faintly at a mere mag. ~13, make only a minuscule contribution to the $\beta$ UMi AB combined light) ¶ AAVSO(VSX), as of 2022 Jul. 30, notes existence of NSV entry, finds that no AAVSO observations are available, offers the possible V-band range 2.02–2.08 (AAVSO(VSX) assessment: status flag = suspected variable; no variability classification symbol); further photometric study advisable? ¶ Fe underabundant, Ba possibly slightly overabundant ¶ <a href="#">2008A&amp;A...483L.43T</a> suggests (via <i>CORIOLIS-SMEI</i> ) two short-lived radial-pulsation mods ¶ <a href="#">2014A&amp;A...566A.67L</a> announces exoplanet B. 5.2, 231" (2012) <b>Zubenelgenubi</b> angular distance from $\alpha$ Lib B, which shares the proper motion of $\alpha$ Lib A, entails Aa,Ab–B distance $\geq$ 5500 au; if B and the A system are gravitationally bound, then their period is $\geq$ 200,000 y; alternative names for the $\alpha$ Lib Aa,Ab pairing and the single star $\alpha$ Lib B are $\alpha^2$ Lib and $\alpha^1$ Lib, respectively, with "1" signalling the fact that $\alpha^1$ Lib, lying to W of $\alpha^2$ Lib, although fainter than "2," is the earlier of the two in its crossing of the local meridian ¶ $\alpha$ Lib Aa, Ab are resp. mags. 3.30, 3.70; for this tight pairing (an SB as well as a resolved system), WDS documents just 4 measurements (from 2017), for angular separations perhaps $<$ 0.1" (distance between $\alpha$ Lib Aa and $\alpha$ Lib Ab may be a few tenths of 1 au); one of $\alpha$ Lib Aa,Ab is overabundant in some metals, perhaps due to the influence of its SB companion; $\alpha$ Lib B is likewise now resolved as $\alpha$ Lib Ba, $\alpha$ Lib Bb, at resp. mags. 4.30 and 7.70, with just 2 astrometry measurements (1999 and 2018, angular separations 0.4", 0.2" respectively); celestial-sphere neighbours $\alpha$ Lib C and $\alpha$ Lib E are faint; gravitationally bound neighbour D (at very small angular separation) is mag. 7.31 (only 3 astrometry measurements, 1991→1999) ¶ lunar occultations are possible, planetary occultations possible yet rare ¶ AAVSO(VSX) assessment in 2022: status flag = suspected variable; no variability classification symbol
$\alpha$	Lib Aa <sup>†</sup>	14 52.2–16 08	2.75	0.15	A3 III–IV <sup>†</sup>	43	0.9	76	0.126	237	-10 SB <sup>†</sup>	<b>Zubenelgenubi</b> angular distance from $\alpha$ Lib B, which shares the proper motion of $\alpha$ Lib A, entails Aa,Ab–B distance $\geq$ 5500 au; if B and the A system are gravitationally bound, then their period is $\geq$ 200,000 y; alternative names for the $\alpha$ Lib Aa,Ab pairing and the single star $\alpha$ Lib B are $\alpha^2$ Lib and $\alpha^1$ Lib, respectively, with "1" signalling the fact that $\alpha^1$ Lib, lying to W of $\alpha^2$ Lib, although fainter than "2," is the earlier of the two in its crossing of the local meridian ¶ $\alpha$ Lib Aa, Ab are resp. mags. 3.30, 3.70; for this tight pairing (an SB as well as a resolved system), WDS documents just 4 measurements (from 2017), for angular separations perhaps $<$ 0.1" (distance between $\alpha$ Lib Aa and $\alpha$ Lib Ab may be a few tenths of 1 au); one of $\alpha$ Lib Aa,Ab is overabundant in some metals, perhaps due to the influence of its SB companion; $\alpha$ Lib B is likewise now resolved as $\alpha$ Lib Ba, $\alpha$ Lib Bb, at resp. mags. 4.30 and 7.70, with just 2 astrometry measurements (1999 and 2018, angular separations 0.4", 0.2" respectively); celestial-sphere neighbours $\alpha$ Lib C and $\alpha$ Lib E are faint; gravitationally bound neighbour D (at very small angular separation) is mag. 7.31 (only 3 astrometry measurements, 1991→1999) ¶ lunar occultations are possible, planetary occultations possible yet rare ¶ AAVSO(VSX) assessment in 2022: status flag = suspected variable; no variability classification symbol
$\beta$	Lup	15 00.1–43 14	2.67 <sup>†</sup>	-0.23	B2 IV <sup>†</sup>	9	-2.7	380	0.054	222	0 SB	has been claimed to be slight ( $\beta$ Cep) var, salient period 0.232 d; however, AAVSO(VSX) in 2022 has no status flag (so not confirmed variable, not suspected variable, not confirmed non-variable), and a fortiori assigns no variability type symbol ¶ fast rotator ( $<$ 3.4 d) ¶ low metallicity
$\kappa$	Cen <u>Aa</u> <sup>†</sup>	15 00.7–42 12	3.13 <sup>†</sup>	-0.21 <sup>†</sup>	B2 V	9	-2.2	400	0.029	218	+8 SB	strictly a triple system, Aa+Ab+B; B mag. 11.5; AB 4", PA 84°→83°, 1926→2000, separation $\geq$ 470 au; $\geq$ 3000 y; Aa-to-Ab distance possibly ~10 au, period possibly ~10 y ( <a href="http://stars.astro.illinois.edu/sow/kappacen.html">stars.astro.illinois.edu/sow/kappacen.html</a> discusses various physical uncertainties),

$\beta$	Boo	15 02.8+40 18	3.51 <sup>†</sup>	0.96	G8 IIIa <sup>†</sup>	14.5	-0.7	230	0.049	234	-20	V?	<p>with mags. resp. 3.34, 4.71, and with angular separation 0.1"→0.1", 1991→2020</p> <p>¶ m<sub>v</sub>, B-V values are for <math>\kappa</math> Cen Aa,Ab,B combined light</p> <p>¶ line profiles vary; although the Aa+Ab binarity has made variability classification difficult, <math>\kappa</math> Cen Aa is classified by AAVSO(VSX), viewed 2021 Jan. 16 and 2022 Jul. 14, as a (slight) variable of the <math>\beta</math> Cep type, 3.13–3.14 in V, 2.2878 h (AAVSO(VSX) assessment: status flag = confirmed variable; variability classification symbol = "BCEP")</p> <p>Nekkar</p> <p>Ba 0.4, Fe -0.5</p> <p><a href="#">1995A&amp;A...296..509H</a> discusses the puzzling flare seen by <i>ROSAT</i> 1993 Aug. 08 (unusual for a lone M giant; it is possible, but seems unlikely, that flare came instead from an undetected M-dwarf companion; the mild Ba enhancement is, admittedly, consistent with presence of such a companion); slow rotator (~200 d)</p> <p>¶ AAVSO(VSX), as of 2022 Jul. 30, asserts variability, notes availability of 53 AAVSO observations, gives V-passband range 3.47–3.50, but is not yet able to assign a canonical variability type (AAVSO(VSX) assessment: status flag = confirmed variable; variability classification symbol = "VAR"); further photometric study advisable?</p>
$\sigma$	Lib	15 05.5–25 22	3.30v <sup>†</sup>	1.68	M2.5 III	11	-1.5	290	0.083	239	-4		<p>[THIS STAR ONLY IN ONLINE VERSION OF TABLE]</p> <p>semireg. var.: 3.2–3.46 in V, mean period 20 d Brachium</p> <p>(AAVSO(VSX) assessment in 2022: status flag = confirmed variable; variability classification symbol = "SRB")</p> <p>there is also rapid microvariability</p> <p>¶ highly evolved (on AGB, with dead carbon-oxygen core)</p>
$\zeta$	Lup A <sup>†</sup>	15 14.0–52 11	3.40	0.92	G8 III	~27.8	0.6	117	0.133	238	-10		<p>B: 6.74; 71.7" (2020), PA 249°→249°, 1826→2020</p> <p>A-to-B distance <math>\geq</math> 2600 au; shared proper motion suggests true binarity (period possibly <math>\geq</math> 68,000 y)</p> <p>¶ <math>\zeta</math> Lup A is in evolutionary terms on "Red Clump" (was Sun-like when still on MS, but helium flash now finished, core-helium fusion now underway)</p> <p>¶ AAVSO(VSX) situation in 2022: no status flag (so not confirmed variable, not suspected variable, not confirmed non-variable), and a fortiori no variability classification symbol</p> <p>a very wide true binary: B is mag. 7.84, 105" (2017)</p> <p>PA 84°→78°, 1780→2017, A-to-B distance <math>\geq</math> 3800 au, period 120,000 y (with shared proper motion indicating true binarity); the SB that is <math>\delta</math> Boo A is not as yet resolved, even in interferometry (so WDS cannot as yet write "δ Boo Aa," "δ Boo Ab")</p> <p>¶ m<sub>v</sub>, B-V values are for <math>\delta</math> Boo A, rather than for <math>\delta</math> Boo AB combined light; for <math>\delta</math> Boo B, GCPD gives m<sub>v</sub>, B-V values 7.84, 0.59</p> <p>¶ <math>\delta</math> Boo A is CN weak; <math>\delta</math> Boo B could be a subdwarf, consistently with the observed low metallicity of <math>\delta</math> Boo A</p> <p>¶ <math>\delta</math> Boo A is in evolutionary terms a "Red Clump" star (core-helium fusion now underway)</p> <p>¶ AAVSO(VSX), as of 2022 Jul. 30, notes existence of NSV entry and availability of 6 AAVSO observations, offers the possible V-passband range 3.44–3.50 (AAVSO(VSX) assessment: status flag = suspected variable; no variability classification symbol); further photometric study advisable?</p>
$\delta$	Boo A <sup>†</sup>	15 16.5+33 14	3.48 <sup>†</sup>	0.95 <sup>†</sup>	G8 III Fe-1 <sup>†</sup>	~26.8	0.6	122	0.140	143	-12	SB	<p>the SB that is <math>\delta</math> Boo A is not as yet resolved, even in interferometry (so WDS cannot as yet write "δ Boo Aa," "δ Boo Ab")</p> <p>¶ m<sub>v</sub>, B-V values are for <math>\delta</math> Boo A, rather than for <math>\delta</math> Boo AB combined light; for <math>\delta</math> Boo B, GCPD gives m<sub>v</sub>, B-V values 7.84, 0.59</p> <p>¶ <math>\delta</math> Boo A is CN weak; <math>\delta</math> Boo B could be a subdwarf, consistently with the observed low metallicity of <math>\delta</math> Boo A</p> <p>¶ <math>\delta</math> Boo A is in evolutionary terms a "Red Clump" star (core-helium fusion now underway)</p> <p>¶ AAVSO(VSX), as of 2022 Jul. 30, notes existence of NSV entry and availability of 6 AAVSO observations, offers the possible V-passband range 3.44–3.50 (AAVSO(VSX) assessment: status flag = suspected variable; no variability classification symbol); further photometric study advisable?</p>
$\beta$	Lib	15 18.3 -9 28	2.61 <sup>†</sup>	-0.11	B8 IIIIn	~17.6	-1.2	190	0.100	259	-35	SB	<p>Zubeneschamali</p> <p>AAVSO(VSX), as of 2022 Jul. 30, notes the existence of 32 AAVSO observations, asserts variability for the <math>\beta</math> Lib system, states V-passband range as 2.60–2.62, and is as yet unable to assign a conjectured canonical variability type</p> <p>(AAVSO(VSX) assessment in 2022: status flag = confirmed variable; variability classification symbol = "VAR");</p>

												the possibility of wide secular variations is suggested by the fact that Eratosthenes, resp. Ptolemy, asserted $\beta$ Lib to be brighter than, resp. equal to, $\alpha$ Sco; further photometric study advisable? ¶ rapid rotator ¶ E(B-V) = -0.02
$\gamma$	UMi	15 20.7+71 45	3.05 <sup>†</sup> 0.06	A3 III <sup>†</sup>	6.7 -2.9 490	0.025 315	-4 V					Pherkad a rapid rotator, and (despite being in MK type A, not B) said to be a variable shell star (cf <a href="#">2000A&amp;A...354..157H</a> ; BSC5: "shell possibly variable," H and CaII variable); AAVSO(VSX), however, viewed 2021 Jan. 16 and 2022 Jul. 14, classifies this as a low-amplitude variable in the $\delta$ Sct group, giving range 0.05 in V, salient period 3.43 h (AAVSO(VSX) assessment: status flag = confirmed variable; variability classification symbol = "DSCTC")
$\gamma$	TrA	15 21.1-68 46	2.88 0.01	A1 III <sup>n†</sup>	17.7 -0.9 184	0.074 244	-3 V					has been asserted to be chemically anomalous (Eu overabundance), and also, not quite consistently, has been classed as a rapid rotator (< 1.2 d) ¶ although we here give MK luminosity class III, class V has also been asserted; Astron. Alm. (epoch 2021.5) assigns MK type A1 III ¶ IR excess has been asserted (circumstellar disk?) ¶ AAVSO(VSX), as of 2022 Jul. 30, notes existence of NSV entry, but denies variability (AAVSO(VSX) assessment: status flag = confirmed non-variable; no variability classification symbol assigned (but should the symbol "CST" be assigned, the NSV entry notwithstanding?)
$\delta$	Lup	15 22.9-40 44	3.22 <sup>†</sup> -0.22	B1.5 IV <sup>n</sup>	4 -3.9 900	0.032 218	0 V?					rapid rotator (< 2.4 d) ¶ a (low-amplitude) variable of the $\beta$ Cep type, 3.2-3.24 in V (AAVSO(VSX), viewed 2021 Jan. 16 and 2022 Jul. 14; AAVSO(VSX) assessment has status flag = confirmed variable, variability classification symbol = "BCEP"), with a single period known, 3.971 h (cf <a href="#">2007MNRAS.377..645S</a> )
$\varepsilon$	Lup A <sub>a</sub> <sup>†</sup>	15 24.3-44 46	3.37 <sup>†</sup> -0.18 <sup>†</sup>	B2 IV-V	6 -2.6 500	-0.030 -230	+8 SB2					A: 3.56; B: 5.04, 0.1", PA 285°→53°, 1883→2019 orbit 737 y; in more detail, a (probable) hierarchical quadruple; although B experiences A as essentially a point mass, in fact A is SB, interferometrically resolved as $\varepsilon$ Lup Aa, $\varepsilon$ Lup Ab with a single 2010 measurement (mags. 3.60, 5.10, angular separation 0.1"), for which <a href="#">2005A&amp;A...440..249U</a> gives SB period 4.55970 d (classifying the primary as a suspect $\beta$ Cep variable and the secondary as a new $\beta$ Cep variable); experiencing AB, on the other hand, as essentially a point mass is the (probably) gravitationally bound C (mag. 9.10; 19"→26", PA 174°→168°, 1826→2020; AB-to-C distance $\geq$ 4100 au; if gravitationally bound, then period $\geq$ 60,000 y); in its stable kinematics, this putative hierarchical quadruple may be contrasted with the unstable, nonhierarchical $\theta$ Ori system, and in its detailed organization with the stable, hierarchical, but mere "double-double" $\varepsilon$ Lyr system; AAVSO(VSX) as viewed 2021 Aug. 11 and 2022 Jul. 14 indicates just slight variability for the $\varepsilon$ Lup system, 3.36-3.38 in V, period seemingly same as the SB period (AAVSO(VSX) assessment: status flag = confirmed variable; variability classification symbol = "HB+SPB") ¶ $m_v$ , B-V values are for $\varepsilon$ Lup Aa, Ab, B combined light possible slight var. (3.26-3.25 in V passband?) Edasich
$\iota$	Dra A+1P <sup>†</sup>	15 25.5+58 53	3.29 <sup>†</sup> 1.17	K2 III <sup>†</sup>	32.2 0.8 101	0.019 334	-11					AAVSO(VSX), as of 2022 Jul. 30, notes existence of NSV entry, finds no record of AAVSO observations (AAVSO(VSX) assessment:

$\alpha$	CrB	15 35.7+26 38	2.23v <sup>†</sup> -0.02	A0 IV (composite) <sup>†</sup> 43	0.4	75	0.150	127	+2 SB <sup>†</sup>	<p>status flag = suspected variable; no variability classification symbol); further photometric study advisable? ¶ <a href="#">2002ApJ...576..478F</a> announces substellar-mass companion and discusses possibility of transits; this is the first discovery of a planet or brown dwarf (IAU name: Hypatia) orbiting a star that has finished stable core-hydrogen fusion; <a href="#">exoplanet.eu/catalog/HIP%2075458_b</a> may from time to time have updates; its substellar companion notwithstanding, <math>\iota</math> Dra has metallicity only slightly greater than solar ¶ since the <math>\iota</math> Dra A substellar-mass companion is known through radial-velocity work, and has not yet been resolved (not even interferometrically), WDS is not as yet able to write "<math>\iota</math> Dra Aa," "<math>\iota</math> Dra Ab"; the widely separated celestial-sphere neighbour catalogued in WDS as <math>\iota</math> Dra B (mag. 8.87, 255"→253", PA 50°→51°, 1879→2020) is not gravitationally bound to the <math>\iota</math> Dra A system) ecl.: 2.21–2.32 in B band, 17 d <b>Alphecca</b> (more precisely, from AAVSO(VSX) as viewed 2021 Jan. 16 and 2022 Jul. 14, 17.359907 d): this SB is a detached binary, of <math>\beta</math> Per type (AAVSO(VSX) assessment: status flag = confirmed variable; variability classification symbol = "<b>EA/DM</b>"); distance between components 0.13 au min; as with <math>\beta</math> Per, so also with <math>\alpha</math> CrB, instrumental photometry reveals both the primary and the secondary eclipse; components have not been interferometrically resolved (so WDS-conformant designation is still "<math>\alpha</math> CrB," not "<math>\alpha</math> CrB A" and "<math>\alpha</math> CrB B") ¶ individual MK types are difficult: primary possibly A0 V, secondary possibly G5 ¶ primary has IR excess (debris disk?) ¶ secondary is X-ray visible and is a rather rapid rotator (~9 d or ~7 d or less, so not tidally locked) ¶ non-IAU name Gemma denotes <math>\alpha</math> CrB as "gem of the Northern Crown" A: 3.0; B: 4.4; similar spectra 0.8" (2019) PA 94°→275°, 1835→2019; maximum angular separation 1980, minimum angular separation 2075; orbit 190 y: <math>\gamma</math> Lup AB orbit is seen nearly edge-on; separation 41 au min, 128 au max, 84.5 au average; <a href="#">stars.astro.illinois.edu/sow/gammalup.html</a> has an orbit map, showing that observational coverage is imperfect (green for micrometry (with large error bars), violet for photography, blue for interferometry); <math>\gamma</math> Lup A is itself SB (2.801 d, unresolved), making this a hierarchical triple system, with the primary in the <math>\gamma</math> Lup A pairing a fast rotator (&lt; 1 d, so not tidally locked) ¶ BSC5 asserts expanding circumstellar shell, and (citing 1987 Vainu Bappu spectra) notes emission peaks in H<math>\alpha</math> profiles, says possibly in transition from B to Be ¶ m<sub>v</sub>, B–V values are for <math>\gamma</math> Lup AB combined light ¶ the <math>\gamma</math> Lup system harbours slight variability, range 2.77–2.79 in V, with AAVSO(VSX) giving period 2.85 d (equalling the period of the SB?), light variations involve "reflection" (irradiation, causing re-radiation); AAVSO(VSX) assessment in 2022 has status flag = confirmed variable; variability classification symbol = "<b>R</b>" slight semiregular variable? (range ~0.2 in V?) Unukalhai AAVSO(VSX), as of 2022 Jul. 30, notes the existence of NSV entry and the availability of 60 AAVSO observations (AAVSO(VSX) assessment: status flag = suspected variable; variability classification symbol = "<b>SR</b>:"); further photometric study advisable?</p>	
$\gamma$	Lup $\Delta$ <sup>†</sup>	15 36.7–41 15	2.78 <sup>†</sup> -0.21 <sup>†</sup>	B2 IVn <sup>†</sup>	8	-2.8	400	-0.030	~212	+2 V	<p>status flag = suspected variable; no variability classification symbol); further photometric study advisable? ¶ <a href="#">2002ApJ...576..478F</a> announces substellar-mass companion and discusses possibility of transits; this is the first discovery of a planet or brown dwarf (IAU name: Hypatia) orbiting a star that has finished stable core-hydrogen fusion; <a href="#">exoplanet.eu/catalog/HIP%2075458_b</a> may from time to time have updates; its substellar companion notwithstanding, <math>\iota</math> Dra has metallicity only slightly greater than solar ¶ since the <math>\iota</math> Dra A substellar-mass companion is known through radial-velocity work, and has not yet been resolved (not even interferometrically), WDS is not as yet able to write "<math>\iota</math> Dra Aa," "<math>\iota</math> Dra Ab"; the widely separated celestial-sphere neighbour catalogued in WDS as <math>\iota</math> Dra B (mag. 8.87, 255"→253", PA 50°→51°, 1879→2020) is not gravitationally bound to the <math>\iota</math> Dra A system) ecl.: 2.21–2.32 in B band, 17 d <b>Alphecca</b> (more precisely, from AAVSO(VSX) as viewed 2021 Jan. 16 and 2022 Jul. 14, 17.359907 d): this SB is a detached binary, of <math>\beta</math> Per type (AAVSO(VSX) assessment: status flag = confirmed variable; variability classification symbol = "<b>EA/DM</b>"); distance between components 0.13 au min; as with <math>\beta</math> Per, so also with <math>\alpha</math> CrB, instrumental photometry reveals both the primary and the secondary eclipse; components have not been interferometrically resolved (so WDS-conformant designation is still "<math>\alpha</math> CrB," not "<math>\alpha</math> CrB A" and "<math>\alpha</math> CrB B") ¶ individual MK types are difficult: primary possibly A0 V, secondary possibly G5 ¶ primary has IR excess (debris disk?) ¶ secondary is X-ray visible and is a rather rapid rotator (~9 d or ~7 d or less, so not tidally locked) ¶ non-IAU name Gemma denotes <math>\alpha</math> CrB as "gem of the Northern Crown" A: 3.0; B: 4.4; similar spectra 0.8" (2019) PA 94°→275°, 1835→2019; maximum angular separation 1980, minimum angular separation 2075; orbit 190 y: <math>\gamma</math> Lup AB orbit is seen nearly edge-on; separation 41 au min, 128 au max, 84.5 au average; <a href="#">stars.astro.illinois.edu/sow/gammalup.html</a> has an orbit map, showing that observational coverage is imperfect (green for micrometry (with large error bars), violet for photography, blue for interferometry); <math>\gamma</math> Lup A is itself SB (2.801 d, unresolved), making this a hierarchical triple system, with the primary in the <math>\gamma</math> Lup A pairing a fast rotator (&lt; 1 d, so not tidally locked) ¶ BSC5 asserts expanding circumstellar shell, and (citing 1987 Vainu Bappu spectra) notes emission peaks in H<math>\alpha</math> profiles, says possibly in transition from B to Be ¶ m<sub>v</sub>, B–V values are for <math>\gamma</math> Lup AB combined light ¶ the <math>\gamma</math> Lup system harbours slight variability, range 2.77–2.79 in V, with AAVSO(VSX) giving period 2.85 d (equalling the period of the SB?), light variations involve "reflection" (irradiation, causing re-radiation); AAVSO(VSX) assessment in 2022 has status flag = confirmed variable; variability classification symbol = "<b>R</b>" slight semiregular variable? (range ~0.2 in V?) Unukalhai AAVSO(VSX), as of 2022 Jul. 30, notes the existence of NSV entry and the availability of 60 AAVSO observations (AAVSO(VSX) assessment: status flag = suspected variable; variability classification symbol = "<b>SR</b>:"); further photometric study advisable?</p>
$\alpha$	Ser A	15 45.4 +6 21	2.64 <sup>†</sup> 1.17	K2 IIIb CN1 <sup>†</sup>	44	0.9	74	0.141	71	+3 V?	<p>status flag = suspected variable; no variability classification symbol); further photometric study advisable? ¶ <a href="#">2002ApJ...576..478F</a> announces substellar-mass companion and discusses possibility of transits; this is the first discovery of a planet or brown dwarf (IAU name: Hypatia) orbiting a star that has finished stable core-hydrogen fusion; <a href="#">exoplanet.eu/catalog/HIP%2075458_b</a> may from time to time have updates; its substellar companion notwithstanding, <math>\iota</math> Dra has metallicity only slightly greater than solar ¶ since the <math>\iota</math> Dra A substellar-mass companion is known through radial-velocity work, and has not yet been resolved (not even interferometrically), WDS is not as yet able to write "<math>\iota</math> Dra Aa," "<math>\iota</math> Dra Ab"; the widely separated celestial-sphere neighbour catalogued in WDS as <math>\iota</math> Dra B (mag. 8.87, 255"→253", PA 50°→51°, 1879→2020) is not gravitationally bound to the <math>\iota</math> Dra A system) ecl.: 2.21–2.32 in B band, 17 d <b>Alphecca</b> (more precisely, from AAVSO(VSX) as viewed 2021 Jan. 16 and 2022 Jul. 14, 17.359907 d): this SB is a detached binary, of <math>\beta</math> Per type (AAVSO(VSX) assessment: status flag = confirmed variable; variability classification symbol = "<b>EA/DM</b>"); distance between components 0.13 au min; as with <math>\beta</math> Per, so also with <math>\alpha</math> CrB, instrumental photometry reveals both the primary and the secondary eclipse; components have not been interferometrically resolved (so WDS-conformant designation is still "<math>\alpha</math> CrB," not "<math>\alpha</math> CrB A" and "<math>\alpha</math> CrB B") ¶ individual MK types are difficult: primary possibly A0 V, secondary possibly G5 ¶ primary has IR excess (debris disk?) ¶ secondary is X-ray visible and is a rather rapid rotator (~9 d or ~7 d or less, so not tidally locked) ¶ non-IAU name Gemma denotes <math>\alpha</math> CrB as "gem of the Northern Crown" A: 3.0; B: 4.4; similar spectra 0.8" (2019) PA 94°→275°, 1835→2019; maximum angular separation 1980, minimum angular separation 2075; orbit 190 y: <math>\gamma</math> Lup AB orbit is seen nearly edge-on; separation 41 au min, 128 au max, 84.5 au average; <a href="#">stars.astro.illinois.edu/sow/gammalup.html</a> has an orbit map, showing that observational coverage is imperfect (green for micrometry (with large error bars), violet for photography, blue for interferometry); <math>\gamma</math> Lup A is itself SB (2.801 d, unresolved), making this a hierarchical triple system, with the primary in the <math>\gamma</math> Lup A pairing a fast rotator (&lt; 1 d, so not tidally locked) ¶ BSC5 asserts expanding circumstellar shell, and (citing 1987 Vainu Bappu spectra) notes emission peaks in H<math>\alpha</math> profiles, says possibly in transition from B to Be ¶ m<sub>v</sub>, B–V values are for <math>\gamma</math> Lup AB combined light ¶ the <math>\gamma</math> Lup system harbours slight variability, range 2.77–2.79 in V, with AAVSO(VSX) giving period 2.85 d (equalling the period of the SB?), light variations involve "reflection" (irradiation, causing re-radiation); AAVSO(VSX) assessment in 2022 has status flag = confirmed variable; variability classification symbol = "<b>R</b>" slight semiregular variable? (range ~0.2 in V?) Unukalhai AAVSO(VSX), as of 2022 Jul. 30, notes the existence of NSV entry and the availability of 60 AAVSO observations (AAVSO(VSX) assessment: status flag = suspected variable; variability classification symbol = "<b>SR</b>:"); further photometric study advisable?</p>

$\mu$	Ser <u>A</u>	15 50.8 –3 30	3.54 –0.04	A0 III	19	0.0	170	0.104	255	–9 SB <sup>†</sup>	<p>¶ a “strong-lined giant” (although [Fe/H] metallicity is not very much above solar)</p> <p>¶ a modest X-ray source</p> <p>¶ has borne also the (not IAU-official) name Cor Serpentis (“Heart of the Serpent”), despite being the principal luminary of Serpens Caput (“Serpent Head”)</p> <p>binary resolved with speckle interferometry, and subsequently (2010NewA...15.324G) analyzed with astrometry: B is mag. 5.39, 0.2” → 0.4”, 1991 → 2018; 2010NewA...15.324G offers an orbital solution, with period 36±2 y, <math>e=0.4\pm 0.3</math>; these authors remark that the low precision of their orbit-based mass determinations leaves various possibilities open regarding the nature of <math>\mu</math> Ser B (“A or F dwarf, subgiant, giant or even a pair of late-type dwarfs”)</p> <p>¶ AAVSO situation in 2022: no status flag (so not confirmed variable, not suspected variable, not confirmed non-variable), and a fortiori no variability classification symbol</p> <p>[THIS STAR ONLY IN ONLINE VERSION OF TABLE]</p>
$\beta$	TrA A	15 57.2–63 30	2.84 <sup>†</sup> 0.30 <sup>†</sup>	F0 IV <sup>†</sup>	~80.8	2.4	40.4	0.444	205	0	<p><i>Spitzer Space Telescope</i> finds IR excess (debris disk?)</p> <p>¶ rapid rotator (slightly &lt; 1 d), with detectable magnetic field</p> <p>¶ metals vary widely (some overabundant, some underabundant)</p> <p>¶ <math>m_v</math>, B–V values are for <math>\beta</math> TrA A (but these can differ only negligibly from the corresponding values for <math>\beta</math> TrA AB combined light, since <math>\beta</math> TrA B (widely separated, at ~152”), is faint, with GCPD giving <math>m_v</math>, B–V as 13.22, 0.83)</p> <p>¶ AAVSO(VSX) situation in 2022: no status flag (so not confirmed variable, not suspected variable, not confirmed non-variable), and a fortiori no variability classification symbol</p>
$\pi$	Sco <u>Aa</u>	16 00.3–26 11	2.89 <sup>†</sup> –0.19 <sup>†</sup>	B1 V <sup>†</sup>	6	–3.4	600	0.029	203	–3 SB2	<p>Aa,Ab ecl.(?) SB; mags. 3.4, 4.5; 1.57d; 2.88–2.91 in V Fang (more precisely, from AAVSO(VSX) viewed 2021 Dec. 03 and 2022 Jul. 14, 1.570103 d [but a published orbital solution gives instead 1.5700925 d], WDS asserts, and AAVSO(VSX) denies, that orbit is so close to edge-on as to be eclipsing) (AAVSO(VSX) assessment: status flag = confirmed variable; variability classification symbol = “<b>ELL</b>”); if not eclipsing, then the observed slight variability is the effect of ellipsoidally distorted stars presenting different surface areas to the photometer at different stages in their mutual orbit); orbit is circular or nearly circular (two published orbital solutions disagree slightly, asserting <math>e=0</math>, <math>e=0.15</math>), possibly with tidal locking, Aa-to-Ab distance possibly ~0.07 au; the binarity has been detected also via occultation; WDS documents just one Aa,Ab measurement (in the year 2000, with angular separation 2”); system has been said, but not at AAVSO(VSX), to be of <math>\beta</math> Lyr type; AAVSO observations archive, as viewed 2021 Dec. 3, indicates a longstanding shortage of photometry (and Kaler at <a href="http://stars.astro.illinois.edu/sow/pisco.html">stars.astro.illinois.edu/sow/pisco.html</a> additionally discusses some difficulties in astrophysical modelling); further photometric study advisable?</p> <p>¶ <math>m_v</math>, B–V values are for <math>\pi</math> Sco Aa,Ab combined light</p> <p>¶ E(B–V) = +0.08</p>
T	CrB A	16 00.5+25 51	9.90v <sup>†</sup> 1.40 <sup>†</sup>	M3 III <sup>†</sup>	—	0.6	2500?	0.011	329	–29 SB	<p>recurrent nova 1866&amp;1946 mags 3&amp;2; V=9.7(2022 Jul. 12)</p> <p>only ten galactic recurrent novae are currently known (2010ApJS...187..275S; these are by definition novae known to recur, and yet lacking the short periods of dwarf novae)</p> <p>¶ T CrB A partner in the recurrent-nova activity, T CrB B, is WD with hot circumstellar accretion (dominating the aggregate T Cr AB signal in UV) of MK type Bep, orbit 227.5 d or 227.6 d,</p>

												<p>A-to-B distance ~0.5 au; angular separation has been measured only twice (in 1946 (considered doubtful by WDS) and 2010, as 0.3" and 0.7" respectively)</p> <p>¶ <math>m_v</math>, B-V values are for T CrB AB combined light</p> <p>¶ long documented in Handbook as mag. 10.08, T CrB AB (combined light) brightened from February 2015, attaining ~9.2 in April 2015 (with V mag. 9.555 and 9.655, on the other hand, are reported in general AAVSO database (not AAVSO(VSX) database) for 2022 Feb. 15, 2022 Jul. 12); AAVSO(VSX), while noting the recurrent nova status, additionally assigns variability type "ELL" (non-eclipsing binary, with light variations due to ellipsoid-photosphere geometry), at period of the SB (AAVSO(VSX) assessment: status flag = confirmed variable; variability classification symbol = "NR+ELL");</p> <p>Bob King</p> <p>in <i>Sky&amp;Telescope</i> 2016 Apr. 20 gives recent history, and AAVSO has a backgrounder at <a href="http://www.aavso.org/t-crb">www.aavso.org/t-crb</a>;</p> <p>next eruption 2026, or earlier?</p> <p>B: 7.87, 14.8", PA 22°→19°, 1834→2020 orbit ≥ 26,000 y;</p> <p>a hierarchical system, with remote outlier D at angular distance 135" (separation ≥ 18,000 au, period ≥ 750,000 y), with D experiencing the AB pair as essentially a point mass; η Lup C is not part of this (triple) system, C's angular proximity to AB being a mere line-of-sight coincidence (mag. 9.39; 60"→115", PA 245°→248°, 1825→2015)</p> <p>¶ <math>m_v</math>, B-V values are for η Lup AB combined light; GCPD additionally gives, for η Lup B, 7.87, 0.16</p> <p>¶ although η Lup A is a rapid rotator (&lt; 1.1 d), there is no evidence of a circumstellar disk, and in particular there seems to be no documentation of "Be phenomenon" spectral behaviour; AAVSO(VSX), as of 2022 Jul. 30, notes existence of NSV entry and unavailability of AAVSO observations, assigns possible variability range as "3.41-?" in V, and indicates that</p> <p>the η Lup system presents possibly-yet-not-certainly constant light</p> <p>(AAVSO(VSX) assessment: status flag = suspected variable; variability classification symbol = "CST:");</p> <p>further photometric study advisable?</p>
η	Lup A <sup>†</sup>	16 01.7-38 28	3.41 <sup>†</sup> -0.23 <sup>†</sup>	B2.5 IVn <sup>†</sup>	7	-2.2	440	0.033	211	+8 V		
δ	Sco A <sup>†</sup>	16 01.7-22 41	2.32v <sup>†</sup> -0.12	B0.5 Ve <sup>†</sup>	7	-3.6	440	~0.037	~196	-7		
												<p>¶ η Lup B is chemically peculiar</p> <p>pro-am photm., spectr. data needed esp. 2022-May Dschubba (when this binary passes through periastron): the δ Sco AB interferometrically resolvable SB has been measured since 1973, although binarity was reported from lunar occultation as early as 1901; period is 10.8 y; previous recent periastra were in 1990, 2000, and 2011;</p> <p>orbit is discussed in. e.g. <a href="#">2012ApJ...757...29C</a> (slightly refining the orbital solution of <a href="#">2011ApJ...729L...5T</a>);</p> <p>orbit is remarkable for its extreme elongation (<math>e = 0.94</math>; for most Be binaries with a non-degenerate secondary component, <math>e \approx 0</math>); the suggestion that δ Sco A is itself a tight binary, with period ~20 d, is not now generally favoured (<a href="#">2013ApJ...766..119M</a> argues against the suggestion); connected with the unexpectedly high-eccentricity AB orbit, however, might be some as-yet-undetected distant orbiter, with period ~200 y, perhaps participating in a Lidov-Kozai interaction (<a href="#">2013ApJ...766..119M</a>, and additionally cf. <a href="http://en.wikipedia.org/wiki/Kozai_mechanism">en.wikipedia.org/wiki/Kozai_mechanism</a>);</p> <p>AB angular separation can become as great as ~200 mas, but at periastron diminishes to 5.9 mas, with AB physical distance diminishing to within 0.8 au (a distance ~25x the radius of δ Sco A, so tidal interaction, perhaps even the generation of a tidal trail of ejecta, is to be expected at periastron); orbital plane is inclined only rather gently to the plane of the sky</p>



([2020ApJ...890...86S](#) gives the angle as  $38^\circ$ ), making this binary far from eclipsing

¶ important instance of “Be phenomenon,” offering opportunity to examine a recent disk-building event:  $\delta$  Sco system seemed unremarkable in much of the 20th century, and was even taken as a B0 IV MK standard; [1993A&A...274..870C](#), however, reported Be phenomenon, from spectroscopy at or near the 1990 periastron; at or near the periastron of 2000, Be-phenomenon behaviour in  $\delta$  Sco A became for the first time strongly evident, with pronounced H $\alpha$  emission in spectroscopy, and also with brightening in photometry; the system faded somewhat in 2005, both in V band and in IR, while H $\alpha$  equivalent width (a signature of material toward the outer reaches of the Be-phenomenon disk around  $\delta$  Sco A) increased; the system again brightened in V passband in 2010 (a signature of material being added to the inner reaches of the disk), and stayed bright, with minor V fluctuations, through the 2011 periastron (and for at least some of this period, notably for 2009 through 2012, cyclic photometric variability was observed on timescales of  $\sim 60$  d to  $\sim 100$  d (similar behaviour had been found also for 2000 through 2002; but the orbital periods of the inner and outer portions of the disk are on the order of 0.5 d and 1.5 d, and so are on a different timescale), consistently with a variable rate of mass transfer upward to the disk out of the  $\delta$  Sco A photosphere); the speed of disk growth seems unusually high in the general population of Be stars, where an episode of growth can take decades, and yet rapid rotator  $\delta$  Sco A does not appear to be rotating so rapidly as to diminish effective photosphere gravity at equator down to  $\sim 0$ ; on the modelling of [2020ApJ...890...86S](#), the Be-phenomenon H $\alpha$ -emitting portion of the disk of equatorially ejected gas around  $\delta$  Sco A was of radius  $10 R^*$  in 2000,  $14 R^*$  in 2002,  $11 R^*$  in 2007 (there was a temporary partial dissipation of the disk in 2005),  $46 R^*$  by 2018 (the temporary partial dissipation was followed on this modelling by a period of variability from 2005 to 2009, and by a disk-growth process from 2010 to 2011, with a rather steady state attained from 2011 to 2018 or beyond); [2020ApJ...890...86S](#) remains agnostic on the question whether gravitational perturbations, especially at the outer reaches of the disk, have affected dissipation and growth (for instance, through tidal effects, including the tidal locking, at periastron, of a local density enhancement, such as a spot or a spiral wave? with disk possibly even overflowing  $\delta$  Sco A Roche lobe at periastron, yielding mass transfer to  $\delta$  Sco B [although mass transfer is rejected by a CHARA disk-imaging team, at [2012ApJ...757...29C](#)?]); it is to address this question that photometry and spectroscopy are sought, especially around the 2022 May periastron, both from professionals and from amateurs (cf pro-am 2022-campaign request at [www.aavso.org/delta-sco-campaign](http://www.aavso.org/delta-sco-campaign); AAVSO additionally has a circa-2011 background briefing, with emphasis on the photometry, at [www.aavso.org/vsots\\_delsco](http://www.aavso.org/vsots_delsco); [2013ApJ...766..119M](#) describes pro-am spectroscopy contributions at the 2011 periastron, involving on the amateur side nearly 20 observers from Australia, France, Germany, Portugal, Spain, and the USA); AAVSO(VSX) assigns  $\gamma$  Cas-type, and yet not  $\lambda$  Eri-type, variability, with range 1.59–2.32 in V (AAVSO(VSX) assessment of  $\delta$  Sco system in 2022: status flag = confirmed variable; variability classification symbol = “GCAS”)

¶ our apparent and absolute magnitudes do not reflect the post-2000 combined light of  $\delta$  Sco AB: [www.aavso.org/delta-scorpi](http://www.aavso.org/delta-scorpi) has recent forum discussion, notably on choice of comparison stars for visual photometric estimates; 4 typical AAVSO V-filter photometry reports, from one and the same observer, are 2020 June 20 mag. 1.86, 2020 June 26 mag. 1.81, 2021 June 26 mag. 1.82, and 2021 Jul. 5 mag. 1.78;

$\beta$	Sco Aa <sup>†</sup>	16 06.8–19 52	2.56 <sup>†</sup> –0.07 <sup>†</sup>	B0.5 V	8	–2.9	400	0.025	192	–1 SB	<p>for 2022 Jul. 01 through 14, AAVSO lacks V-filter reports, but does have three visual observations, from two observers, at mag. 2.0 (suggesting, therefore, very modest variability in V at present, a little above the naked-eye threshold); <math>\delta</math> Sco B (not known to be variable) is reported by WDS as of mag. 4.62</p> <p>¶ we follow <a href="#">2020ApJ...890...86S</a> in assigning MK type B0.5 V, which we take to be appropriate for the combined binary-system light (but elsewhere in the literature, a slightly different MK type is assigned); <a href="#">2020ApJ...890...86S</a> suggests B2 V as an MK basis for modelling <math>\delta</math> Sco B</p> <p>¶ we follow <a href="#">2020ApJ...890...86S</a> in assigning, consistently with our policy for using rounding-off to reflect uncertainties, D = 440 ly, and on this basis asserting <math>\pi</math> to be, with reasonable rounding-off, 7 mas</p> <p>¶ <math>\delta</math> Sco AB (and its remoter gravitationally bound third star, if there is such a companion) may possibly be a low-velocity runaway system with ISM bow shock</p> <p>¶ E(B–V)=+0.16</p> <p>Aa: 2.9; B: 10.6, 0.3" (2019); C: 4.52, 13" Acrab (AC astrometry in more detail: 14"→13", PA 25°→20°, 1779→2019); in gross terms a visual binary (as AC), with A-to-C distance <math>\geq</math> 2200 au, period &gt; 16,000 y, but in fact putatively a sextuplet; <a href="https://en.wikipedia.org/wiki/Beta_Scorpii">en.wikipedia.org/wiki/Beta_Scorpii</a> summarizes the sextuplet hierarchy in a diagram (Aa with Ab (6.82 d), and B experiencing Aa+Ab as essentially a point mass (610 y); E as an as a not-yet-resolved SB (although Wikipedia in an informal spirit writes "Ea" and "Eb," WDS, whose terminology we in this Handbook article take as normative, cannot as yet do so; period of this unresolved SB is 10.7 d), and C experiencing the binary system that is as yet just (in WDS-formal terms) "<math>\beta</math> Sco E" as essentially a point mass (39 y, <math>\beta</math> Sco E combined light mag. 6.60); the B+(Aa,Ab) triple is in a wide, &gt; 16,000-y orbit with the C+E triple, around the centre of mass shared by this pair of triples, thereby delivering the gross visual-binary phenomenology studied as <math>\beta</math> Sco AC since 1779; the Aa,Ab angular separation is too small to yield a measure for WDS; the CE separation is tight, measured as 0.1" in 2019; the entire sextuplet has an outlying celestial-sphere neighbour, physically unrelated to the sextuplet, WDS-catalogued as <math>\beta</math> Sco D (mag. 7.5; 520"→518", PA 31°→30°, 1860→1998)</p> <p>¶ <math>m_v</math>, B–V values are for <math>\beta</math> Sco Aa,Ab,B combined light; for <math>\beta</math> Sco C, GCPD additionally appears to give <math>m_v</math>, B–V values 4.91, –0.02</p> <p>¶ AAVSO(VSX), as of 2022 Jul. 30, notes the availability of a single AAVSO observation and offers possible V-passband range 2.61–2.67 (AAVSO(VSX) assessment: status flag = suspected variable; no variability classification symbol); further photometric study advisable?</p> <p>¶ lunar occultations possible, planetary occultations possible yet rare (1971 May 14 occultation by Jovian satellite Io)</p>
$\delta$	Oph A	16 15.6 –3 45	2.73 <sup>†</sup> 1.58	M1 III <sup>†</sup>	–19.1	–0.9	171	0.150	198	–20 V	<p>¶ the name Graffias is not IAU-official slight var.?. (2.72–2.75 in V passband?) Yed Prior<sup>†</sup></p> <p>AAVSO(VSX), as of 2022 Jul. 30, notes the existence of NSV entry and the availability of a single AAVSO observation (AAVSO(VSX) assessment: status flag = suspected variable; no variability classification symbol); <a href="#">1992IBVS.3792....1P</a> finds no variability, but says that since NSV V-passband amplitude is just 0.03 mag., variability cannot be excluded (further photometric study advisable?)</p> <p>¶ slow rotator</p>

													¶ high metallicity ¶ although $\delta$ Oph has finished core hydrogen fusion, its exact evolutionary state is uncertain (cf <a href="http://stars.astro.illinois.edu/sow/yedprior.html">stars.astro.illinois.edu/sow/yedprior.html</a> ) ¶ Astron. Alm. (epoch 2021.5) assigns MK type M0.5 III ¶ naked-eye neighbour Yed Posterior is a mere optical companion, too greatly separated in space for true binarity; the “prior” and “posterior” in the traditional, and as of 2016 IAU-official, names denote the order in that these two (physically unrelated) stars cross the local meridian
$\epsilon$	Oph A	16 19.6 -4 45	3.23	0.97	G9.5 IIIb <sup>†</sup>	31	0.7	106	0.093	64	-10 V		Yed Posterior cyanogen and carbon notably underabundant, suggesting that $\epsilon$ Oph is an interloper from outside the galactic thin disk; Astron. Alm. (epoch 2021.5) assigns MK type G9.5 IIIb Fe-0.5 ¶ slight variability, with AAVSO(VSX) supplying range of 0.003 mag. in V (AAVSO(VSX) assessment in 2022: status flag = confirmed variable; variability classification symbol = “PULS”); <a href="#">2008A&amp;A...478.497K</a> , using <i>MOST</i> mission data, finds pulsation modes favourable to astroseismology slight var.: 2.86-2.94, 0.25 d; B: 8.4, B9 V, 20.5” (2019) Alniyat (AB astrometry in more detail: 22”→20”, PA 270°→269°, 1783→2019); recent studies, including lunar occultation measures, show $\sigma$ Sco to be a quadruple system, with $\sigma$ Sco Aa1,Aa2 in fact SB (33.0 d; considered by WDS to be successfully resolved, although only one PA measurement, from 2010, is documented in WDS), and with the entire 3-star $\sigma$ Sco A configuration (where the (Aa1+Aa2), Ab period is > 100 y) in some slow, wide orbit with $\sigma$ Sco B; orbital solution has been published for Aa1,Aa2, but not for Aa,Ab; <a href="#">2007MNRAS.380.1276N</a> announces interferometric solution for the SB orbit, proposing for primary and secondary the respective MK types B1 III, B1 V; in the SB pair, the primary is a variable of the $\beta$ Cep type (AAVSO(VSX), as of 2022 Jul. 14, gives V-mag. range 2.86-2.94, salient period 5.9241 h, with the following assessment: status flag = confirmed variable; variability classification symbol = “BCEP”); <a href="#">1992A&amp;A...261..203P</a> discusses period changes) ¶ $m_v$ , B-V values are for $\sigma$ Sco Aa1,Aa2,Ab combined light ¶ photography shows $\sigma$ Sco to be embedded in diffuse nebula
$\sigma$	Sco <u>Aa1</u> <sup>†</sup>	16 22.6-25 39	2.87 <sup>†</sup>	0.14 <sup>†</sup>	B1 III <sup>†</sup>	5	-3.7	700	0.019	213	+3 SB <sup>†</sup>		
$\eta$	Dra A	16 24.3 +61 28	2.73 <sup>†</sup>	0.91 <sup>†</sup>	G8 IIIab	35.4	0.5	92	0.059	343	-14 SB <sup>†</sup>		¶ E(B-V) = +0.4 (pronounced reddening) B: 8.2, 4.4”, PA 150°→143°, 1843→2015 orbit $\geq$ 1000 y, separation $\geq$ 140 au ¶ a “Red Clump” resident (evolved, presently stable, performing core-helium fusion) ¶ believed to be a slow rotator (~400 d) ¶ a modest X-ray source ¶ AAVSO(VSX), as of 2022 Jul. 30, notes the existence of NSV entry and the absence of AAVSO observations, and gives the possible V-passband range 2.70-2.74 (AAVSO(VSX) assessment: status flag = suspected variable; no variability classification symbol); further photometric study advisable? ¶ $m_v$ , B-V values are for $\eta$ Dra AB combined light ¶ the $\eta$ Dra AB pair has a reasonably bright outlying celestial-sphere neighbour, WDS-catalogued as $\eta$ Dra C: mag. 8.10, angular separation from $\eta$ Dra A ~565”, PA ~240°, angular separation and PA nearly constant 1866→2003 ¶ near the radiant of the $\eta$ Draconids meteor shower semireg. var.: 0.75-1.21, 5.97 y; B: 5.40, 2.8” (2019) <b>Antares</b> <sup>†</sup> AB astrometry: PA 273°→276°,
$\alpha$	Sco <u>A</u> <sup>†</sup>	16 30.9 -26 29	0.99v <sup>†</sup>	1.83	M1.5 Iab <sup>†</sup>	6	-5.1	600	0.026	207	-26 SB		

												1847→2016; orbit 2500 y? the large magnitude difference makes this a difficult binary (the binarity was discovered only in 1819, through controverted lunar-occultation work ( <a href="#">1972BAA...82..43IK</a> describes the 18.6-year 1940-through-2050 cycle of lunar occultation possibilities); the binarity was established beyond controversy still later, in the 1840s) ¶ AAVSO(VSX): semireg. (with some discussion of period (5.97 y); cf also <a href="#">2013AJ...145..38P</a> , where a true period is found for radial-velocity variations, and the detected variation is judged to be more likely of pulsational than of orbital origin (AAVSO(VSX) assessment in 2022: status flag = confirmed variable, variability classification symbol = “SRC,” V mag. 0.75–1.21; but the variability has also been called, outside AAVSO(VSX), irregular); <a href="#">2018AuJAn...29..89H</a> reports that variability was observed by, and incorporated into the oral tradition of, aboriginals in southern Australia; asserted by Eratosthenes to be fainter than β Lib, and by Ptolemy to equal β Lib ¶ Astron. Alm. (epoch 2021.5) assigns MK type M1.5 Iab–Ib ¶ radius has been studied interferometrically and via lunar occultations (up to 3.4 au; however, even apart from the problem of pulsation, radius determination of highly evolved red stars is wavelength-dependent); one of the two first-mag. supergiants (the other being α Ori Aa [Betelgeuse]) ¶ significant stellar wind, with mass loss almost 1e-6 M <sub>⊙</sub> /y, within which α Sco B has created a locally ionized region ¶ the most massive member of the Sco-Cen Association (the nearest OB association) ¶ B shares in the proper motion of A, indicating true binarity: AB separation is ≥ 530 au, period possibly ~1200 y ¶ location (within zodiac) makes the classical Greek name for “rival of Mars” appropriate not only as regards naked-eye colour but also as regards sky geometry ¶ <a href="#">1972BAA...82..43IK</a> describes the 18.6-year 1940-through-2050 cycle of lunar occultation possibilities possible slight var. (2.76–2.81 in V?) AAVSO(VSX), as of 2022 Jul. 30, notes the existence of NSV entry and the nonexistence of AAVSO observations (AAVSO(VSX) assessment: status flag = suspected variable; no variability classification symbol); further photometric study advisable? ¶ SB period computed 1908, and again 2008, in both cases ~410 d; <a href="#">1977ApJ...214L..79B</a> announces speckle-interferometry resolution of the β Her Aa,Ab SB, with angular separation 43 mas; WDS documents 4 astrometric measurements of the Aa,Ab SB as a visual binary, 1975→1984; Aa,Ab possesses, and AB lacks, a published orbit ¶ m <sub>v</sub> , B–V values are for β Her Aa,Ab combined light ¶ X-ray emission from the SB primary indicates magnetic activity ¶ Astron. Alm. (epoch 2021.5) assigns MK type G7 IIIa Fe–0.5 ¶ Kaler, noting that primary has N enhanced relative to C, says in his overall summation “a very normal star for its state of age” ¶ “Kornephoros” = Gk “club-bearer,” in reference to the weapon of Hercules (compare α Her, which in the pictorial-atlas tradition, marks the hero’s head)
β	Her <a href="#">Aa</a>	16 31.2 +21 26	2.78 <sup>†</sup>	0.93 <sup>†</sup>	G7 IIIa <sup>†</sup>	23	–0.4	140	0.100	261	–26 SB <sup>†</sup>	Kornephoros <sup>†</sup>

$\tau$	Sco	16 37.3 –28 16	2.82 –0.25	B0 V <sup>†</sup>	7	–3.0 500	0.025 203	+2 V	Paikauhale <sup>†</sup>
									<p>intrinsically more luminous than <math>\sigma</math> Sco, but more heavily obscured by ISM</p> <p>¶ anomalous in its UV lines (P Cyg profile)</p> <p>¶ O and Fe are underabundant</p> <p>¶ <a href="#">2006MNRAS.370..629D</a> discusses <math>\tau</math> Sco magnetic topology (poloidal, with also a warped toroidal component of modest strength), including both its origin (more likely a fossil field from the star's (recent) birth than a dynamo effect) and its connection with winds and with the observed hard X-ray emission; the authors note that the topology is stable over the 1.5-y period of their observations (in contrast with a strongly differential-rotation star, such as Sun); in additionally announcing a (refined) rotation period of 41.033 d, the authors comment, “the second-slowest rotator so far known among high-mass stars”</p> <p>¶ Kaler: “among the most-observed stars in the sky”; however, AAVSO(VSX) situation in 2022 is as follows: no status flag (so not confirmed variable, not suspected variable, not confirmed non-variable), and a fortiori no variability classification symbol</p> <p>¶ E(B-V)=+0.06</p> <p>¶ the <math>\tau</math> Sco name Paikauhale was IAU-approved in 2018 Aug. 10; the not-IAU-official “Al Niyat,” or “the arteries of the Heart,” on the other hand, denotes <math>\sigma</math> Sco and <math>\tau</math> Sco jointly, as flanking <math>\alpha</math> Sco</p>
$\zeta$	Oph	16 38.5 –10 37	2.57 <sup>†</sup> 0.02	O9.5 Vne <sup>†</sup>	9	–2.7 370	0.029 <sup>†</sup> 32	–15 V <sup>†</sup>	<p>the nearest O-type star (and consistently with this extreme temperature, resident in an H II region)</p> <p>¶ unusual in being an “Oe,” i.e. an O-star instance of the “Be phenomenon”</p> <p>¶ “runaway star” (consistently with this extreme speed-relative-to-LSR, forming bow shock in ISM), perhaps formerly the secondary in a binary pair whose primary perished in a supernova; <a href="#">2011AN....332..147H</a> confirms magnetic field, discusses X-ray properties, suggests PSR B1919+10 as remnant of the hypothesized defunct companion</p> <p>¶ line of sight to <math>\zeta</math> Oph is one of the most used in spectroscopic studies of ISM</p> <p>¶ <a href="#">2014MNRAS.440.1674H</a> is a recent discussion of variability, from radial and non-radial pulsation modes; AAVSO(VSX), assigning magnitude range 2.56–2.58 in V and period 4.6 h, follows GCVS in treating <math>\zeta</math> Oph as a variable with Be-phenomenon behaviour, and yet lacking the history of outbursts found in the <math>\gamma</math> Cas class (AAVSO(VSX) assessment in 2022: status flag = confirmed variable; variability classification symbol = “BE”);</p> <p><math>\zeta</math> Oph is, on the other hand, classified as <math>\gamma</math> Cas-variable (and is termed a shell star) in BSC5; still elsewhere, <math>\zeta</math> Oph has been treated as a prototype for the “<math>\zeta</math> Oph variables”</p> <p>¶ E(B-V)=+0.32 (pronounced reddening; if ISM were not present, <math>\zeta</math> Oph would reach nearly first mag.)</p> <p>¶ recapitulations of recent <math>\zeta</math> Oph studies include <a href="#">2012MNRAS.427L..50G</a> (MK classification problem, also mass-loss rate in context of “weak-wind problem”),</p>

ζ	Her <u>A</u> <sup>†</sup>	16 42.2+31 34	2.81 <sup>†</sup>	0.64 <sup>†</sup>	G1 IV <sup>†</sup>	93	2.7	35	~0.575	~307	-70 <sup>†</sup> SB	<p><a href="#">2014MNRAS.440.1674H</a> (rotation, pulsation, Ha emission episodes, inferred circumstellar decretion disk, satellite-based photometry), <a href="#">2015ApJ...800.132C</a> (distance, age, mass, effective temperature, bow shock in ISM, ... ); additionally, <a href="#">2012A&amp;A...543A.56D</a> is among the papers describing not only the specific interaction of ζ Oph with ISM, but also the quite general ISM bow-shock topic (noting inter alia that not all runaway stars produce bow shocks)</p> <p>B: 5.40, G7 V, 1.6", PA 110° (2019), orbit 34.45 y orbit well studied since F.G.W. von Struve 1826 micrometry (however, it was Herschel, not von Struve, who discovered the binarity); separation 8 au min, 21 au max, 15 au average, 34.45 y; considered one of the few binaries in which ratio of B mass to sum of A and B masses can be studied both via traditional (non-interferometric) astrometry and via spectroscopy; WDS indicates, however, that an inner binary with orbit ~12 y has been suspected repeatedly, and that the inner-binary component has been detected in IR speckle interferometry (but WDS, at any rate as of 2021 Dec. 13, continues to write simply "ζ Her A," without as yet distinguishing between ζ Her Aa and ζ Her Ab)</p> <p>¶ m<sub>v</sub>, B–V values are for ζ Her AB combined light</p> <p>¶ Astron. Alm. (epoch 2021.5) assigns MK type G0 IV</p> <p>¶ ζ Her A is unusual in its evolutionary phase, being in the Hertzsprung Gap (and so in rapid evolutionary transition)</p> <p>¶ AAVSO(VSX), as of 2022 Jul. 30, notes existence of NSV entry and nonexistence of AAVSO observations, and assigns V-passband range 2.78–2.85 (AAVSO(VSX) assessment: status flag = suspected variable; no variability classification symbol); further photometric study advisable?</p> <p>¶ <a href="#">2001A&amp;A...379..245M</a> summarizes previous work, presents detailed physical modelling for A and B, and discusses asteroseismology, remarking in conclusion that "among the binaries to be calibrated with some confidence, ζ Herculis is one of the most interesting owing to the difference of evolutionary state of components"</p> <p>¶ high velocity relative to Sun possible var., type unknown, 3.47–3.61 in V passband?</p> <p>AAVSO(VSX), as of 2022 Jul. 30, flags the η Her system as possibly harbouring variability, noting the existence of NSV entry and the existence of 2 AAVSO observations (AAVSO(VSX) assessment: status flag = suspected variable; no variability classification symbol); further photometric study advisable?</p> <p>¶ the outlying (116", 84") and faint (mag. 11.7, 13.9) celestial-sphere neighbours η Her B and η Her C aside, a close (0.3") celestial-sphere neighbour of η Her A was suspected in 1842, without subsequent detection</p> <p>¶ in evolutionary terms a resident of the "Red Clump" (fusing helium in stable core)</p> <p>¶ Astron. Alm. (epoch 2021.5) assigns MK type G7 III Fe–1</p> <p>¶ Fe is notably underabundant</p>
η	Her A	16 43.7+38 53	3.50 <sup>†</sup>	0.92	G7.5 IIIb Fe–1 <sup>†</sup>	30.0	0.9	109	0.092	157	+8 V?	<p>AAVSO(VSX), as of 2022 Jul. 30, flags the η Her system as possibly harbouring variability, noting the existence of NSV entry and the existence of 2 AAVSO observations (AAVSO(VSX) assessment: status flag = suspected variable; no variability classification symbol); further photometric study advisable?</p> <p>¶ the outlying (116", 84") and faint (mag. 11.7, 13.9) celestial-sphere neighbours η Her B and η Her C aside, a close (0.3") celestial-sphere neighbour of η Her A was suspected in 1842, without subsequent detection</p> <p>¶ in evolutionary terms a resident of the "Red Clump" (fusing helium in stable core)</p> <p>¶ Astron. Alm. (epoch 2021.5) assigns MK type G7 III Fe–1</p> <p>¶ Fe is notably underabundant</p>
α	TrA A	16 51.2–69 04	1.91	1.45	K2 IIb–IIIa <sup>†</sup>	~8.4	-3.5	390	0.036	150	-3	<p><b>Atria</b></p> <p>anomalous for its MK type, with flares and X-ray emission, perhaps from as-yet-undetected magnetically active companion (a companion would indeed be indicated by the claimed "barium star" status of α TrA;</p> <p><a href="#">stars.astro.illinois.edu/sow/atria.html</a>, in discussing the possibility of a companion, also remarks,</p>

$\epsilon$	Sco	16 51.7–34 20	2.29	1.15	K2 III	51	0.8	64	0.666 <sup>†</sup>	247	-3 <sup>†</sup>	<p>however, “the classic ‘hybrid star,’ a giant that shows evidence for blowing a cool wind from its surface, yet having a hot surrounding magnetic corona at the same time”; <a href="https://ntrs.nasa.gov/archive/nasa/casi.ntrs.nasa.gov/20040086627.pdf">ntrs.nasa.gov/archive/nasa/casi.ntrs.nasa.gov/20040086627.pdf</a> further discusses both <math>\alpha</math> TrA and <math>\beta</math> Dra, as (solitary) stars, which are in this particular posited sense “hybrid”;</p> <p>the faint (mag. 11.4) and outlying (angular separation 92”) celestial-sphere neighbour <math>\alpha</math> TrA B cannot be the postulated flaring and X-ray-bright companion</p> <p>¶ AAVSO(VSX) situation in 2022: no status flag (so not confirmed variable, not suspected variable, not confirmed non-variable), and a fortiori no variability classification flag var.? (2.24–2.35 in V?)</p> <p>AAVSO(VSX), as of 2022 Jul. 30,</p> <p>notes existence of NSV entry, finds no record of AAVSO observations (AAVSO(VSX) assessment: status flag = suspected variable; no variability classification symbol); further photometric study advisable?</p> <p>¶ slow rotator (possibly even 1.3 y)</p> <p>¶ evolved, and yet not a clump star; <a href="https://stars.astro.illinois.edu/sow/epssco.html">stars.astro.illinois.edu/sow/epssco.html</a> discusses the uncertainty in evolutionary stage (brightening, with He core as yet awaiting ignition? dimming, with He core fusion in progress? or brightening, with dead C-and-O core, He-core fusion now over?)</p> <p>¶ high velocity relative to Sun indicates origin outside the galactic thin disk (and metal underabundances are consistent with such an origin)</p>	Larawag
$\mu^1$	Sco A	16 53.5–38 05	3.04v <sup>†</sup> -0.21		B1.5 IVn	7	-2.9	500	0.024 <sup>†</sup>	206–25 <sup>†</sup>	SB2 <sup>†</sup>	<p>ecl.: 2.94–3.22, 1.4463 d</p> <p>(more precisely, in AAVSO(VSX) as viewed 2021 Dec. 14 and again 2022 Jul. 14, 1.44626907 d); 2 published solutions for the orbit give, respectively, <math>e=0.019</math>, <math>e=0.0</math>; semidetached, partially eclipsing binary system, with mass transfer, resembling <math>\beta</math> Lyr in its never-constant light and in exhibiting both primary and secondary minima</p> <p>(AAVSO(VSX) assessment: status flag = confirmed variable; variability classification symbol = “<b>EB/SD</b>”); <a href="https://doi.org/10.1093/mnras/stz398">1948MNRAS.108..398S</a> gives the light curve, and also discusses early observational history (this is the third eclipsing SB discovery in astronomy [made by Bailey, 1896]); distance between components is <math>\sim 0.07</math> au; since the SB is not as yet resolved, even interferometrically, WDS is not as yet able to write “<math>\mu^1</math> Sco Aa,” “<math>\mu^1</math> Sco Ab”</p> <p>¶ <math>\mu^1</math> Sco has celestial-sphere neighbor <math>\mu^2</math> Sco, at mag. 3.6 (so just barely fainter than our chosen “Brightest Stars” magnitude cutoff), at angular separation <math>\sim 5.8'</math> (Mizar-Alcor angular separation is just under 12”; normal naked-eye resolution is taken in ophthalmology, under at any rate some reasonable selection of consulting-room eye chart, to be <math>\sim 1'</math>); <math>\mu^1</math> Sco and <math>\mu^2</math> Sco are the “Little Cat’s Eyes,” as distinct from the “Cat’s Eyes” which are <math>\lambda</math> Sco and <math>\nu</math> Sco; the IAU-official name “Xamidimura” applies just to the primary in the SB which is the right, unresolved, two-star <math>\mu^1</math> Sco system; for the Khoekhoem nomadic pastoralists of SW Africa, on the other hand, “Xami di mura” is “eyes of the lion,” as a designation for the naked-eye challenge pair <math>\mu^1</math> Sco, <math>\mu^2</math> Sco; <math>\mu^1</math> Sco and <math>\mu^2</math> Sco are not gravitationally bound, although both belong to the (gravitationally unbound) “Upper Sco” subgroup of the Sco-Cen Association</p> <p>¶ <math>\mu^1</math> Sco and <math>\mu^2</math> Sco are not gravitationally bound, although both belong to the (gravitationally unbound) “Upper Sco” subgroup of the Sco-Cen Association</p> <p>¶ a little confusingly, <math>\mu^2</math> Sco is also formally</p>	Xamidimura

												<p>“<math>\mu^1</math> Sco H” (<math>\mu^1</math> Sco AH astrometry: 333”→347”, PA 71°→72°, 1752→2015); additionally, the unresolved <math>\mu^1</math> Sco A SB has two other WDS-documented celestial-sphere neighbours, both reasonably bright, <math>\mu^1</math> Sco B (mag. 8.9, 8.9”→9.2”, PA 211°→210°, 1999→2000) and <math>\mu^1</math> Sco G (mag. 9.4, 81”→81”, PA 257°→257°, 1935→2016)</p>		
$\kappa$	Oph	16 58.8 +9 20	3.20 <sup>†</sup>	1.16	K2 III	36	1.0	91	0.292 <sup>†</sup>	268	-56	V <sup>†</sup>	<p>slow rotator (possibly as slow as 1.6 y)  <math>\P</math> historical assertion of variability may be due to a confusion between <math>\kappa</math> Oph and <math>\chi</math> Oph; completely apart from this historical problem, however, <a href="#">2001BaltA...10.593A</a> discusses the possible variability both of <math>\kappa</math> Oph and of other Red Clump stars; AAVSO(VSX) for its part asserts constant light (AAVSO(VSX) assessment in 2022: status flag = confirmed non-variable; variability classification symbol = “CST”)  <math>\P</math> high velocity relative to Sun suggests origin outside the galactic thin disk</p>	
$\zeta$	Ara	17 00.6–56 01	3.12	1.60	K4 III	7	-2.7	490	0.041	206		-6	<p>one of the rather rare instances of a giant excessively bright in far IR  <a href="#">(1997A&amp;A...323.513P</a> suggests that such giants are more likely to be radiating their IR excess from circumstellar debris disks than from winds, and so are to be considered evolved-star analogues of the unevolved (and IR-bright) <math>\alpha</math> Lyr  <math>\P</math> AAVSO(VSX) situation in 2022: no status flag (so not confirmed variable, not suspected variable, not confirmed non-variable), and a fortiori no variability classification symbol</p>	
$\zeta$	Dra $\Delta$	17 08.9+65 41	3.17	-0.12	B6 III <sup>†</sup>	10	-1.8	330	0.028	314		-17	V	<p>Aldhibah  <math>\zeta</math> Dra A and <math>\zeta</math> Dra B are mags. 3.2, 4.2 respectively; a difficult binary, resolved interferometrically, but as of at any rate 2021 Dec. 14 with just 10 astrometry measurements (AB 0.0”→0.1”, 1981→1994); on the preliminary orbital solution offered in <a href="#">1998A&amp;AS...133..149M</a>, the orbital plane coincides with the plane of the sky, and the orbit is circular, with radius 67 mas; <a href="#">1998A&amp;AS...133..149M</a> suggests, tentatively, that <math>\zeta</math> Dra A and <math>\zeta</math> Dra B are “a pair of giants”  <math>\P</math> given the recent formation of the <math>\zeta</math> Dra system, Fe is anomalously underabundant  <math>\P</math> AAVSO(VSX) situation in 2022: no status flag (so not confirmed variable, not suspected variable, not confirmed non-variable), and a fortiori no variability classification symbol  <math>\P</math> E(B-V) =+0.03</p>
$\eta$	Oph $\Delta$ B <sup>†</sup>	17 11.7–15 45	2.43	0.06	A1 IV + A1 IV? <sup>†</sup>	37	0.3	90	-0.107	~22		-1	SB	<p>A: 3.0; B: 3.3, A3 V, 0.5” (2019), orbit 87.6 y <b>Sabik<sup>†</sup></b>  highly eccentric orbit: separation 2 au min, 65 au max  <math>\P</math> under IAU rules, “Sabik” designates <math>\eta</math> Oph A, not <math>\eta</math> Oph B  <math>\P</math> <math>m_v</math>, B-V values are for <math>\eta</math> Oph AB combined light  <math>\P</math> our present assignment of MK types (confident for <math>\eta</math> Oph A, tentative for <math>\eta</math> Oph B) is from the literature; our Handbook predecessor R.F. Garrison, however, himself favoured “A2.5 Va,” perhaps for the AB composite; Astron. Alm. (epoch 2021.5), perhaps again for the AB composite, assigns MK type “A2 Va<sup>+</sup> (Sr)”  <math>\P</math> it is possible that A, or B, or both A and B, are superabundant in metals  <math>\P</math> AAVSO(VSX) situation in 2022: no status flag (so not confirmed variable, not suspected variable, not confirmed non-variable), and a fortiori no variability classification symbol</p>
$\eta$	Sco A	17 13.8–43 16	3.33	0.40	F5 IV <sup>†</sup>	~44.4	1.6	73	0.290	175		-28		<p>possible variability of unknown type  AAVSO(VSX), as of 2022 Jul. 30, notes the existence of NSV entry and the nonexistence of AAVSO observations, states possible V-passband range as “3.31-?”  (AAVSO(VSX) assessment: status flag = suspected variable; no variability classification symbol); further photometric study advisable?</p>



$\alpha$  Her Aa<sup>†</sup> 17 15.7+14 22 3.06<sup>†</sup> 1.45<sup>†</sup> M5 Ib-II 9 -2.4 400 0.032 347 -33 V

¶ we now take MK type (slightly evolved beyond stable core-hydrogen fusion, because in luminosity class “IV”) from NASA NStars work summarized at [2006AJ...132..161G](#) (with Garrison the third author); Garrison had himself previously, in this Handbook table, proposed the dwarf MK type “F2 V:p(Cr)”; the intricacy of his previous type hints at difficulties in classification, and indeed even “dwarf barium star” has been asserted elsewhere; Astron. Alm. (epoch 2021.5) assigns the same MK type as legacy-Garrison

¶ rapid rotator (< 1 d); the observed X-ray emission is consistent with magnetic effects (including coronal heating?) stemming from rapid rotation semireg. var.: 2.73–3.60 in V; B: 5.4, 5” (2019) Rasalgethi<sup>†</sup> the second-nearest AGB star (the nearest being the more dramatic visual variable  $\alpha$  Ceta Aa (Mira))

¶ WDS catalogues  $\alpha$  Her Aa,Ab, but with the caveat that the asserted duplicity may not be real (only 3 measurements are available (0.2”→0.2”, 1986→1991), and attempts to resolve the asserted binary failed over the period 1985→1997, even with speckle interferometry at BTA-6; a period of ~10 y has been suspected if the Aa,Ab pairing is real); the IAU-official name Rasalgethi applies to  $\alpha$  Her Aa if the Aa,Ab pairing is real, and to  $\alpha$  Her A otherwise; although we use WDS nomenclature (for this as for all stars in this table), the literature also (e.g. [1993A&A...274..838T](#), [2013AJ...146..148M](#)) uses an alternative terminology, on which Rasalgethi is designated “ $\alpha^1$  Her A,” and the two as-yet-unresolved SB components are designated “ $\alpha^2$  Her A” and “ $\alpha^2$  Her B”; the as-yet-unresolved SB (MK types G8 III and A9 IV–V, period 51.578 d, distance between the unresolved components ~0.4 au) which both for WDS and for us is designated  $\alpha$  Her B is mag. 5.4 (9”→5”, PA 117°→103°, 1777→2020); AB orbital solution asserts period 3600 y; this makes the  $\alpha$  Her system at least a (kinematically stable, hierarchically organized) triple; the faint celestial-sphere neighbours  $\alpha$  Her C (mag. 15.5) and  $\alpha$  Her D (mag. 11.1), on the other hand, are not part of the system

¶  $m_v$ , B–V values are for  $\alpha$  Her Aa,Ab,B combined light

¶ the mass of Rasalgethi has been controverted (mass as high as ~15  $M_{\odot}$ , putting Rasalgethi into the same mass league as (admittedly less evolved) Betelgeuse and Aldebaran; or, rather, as low as ~2  $M_{\odot}$ , putting Rasalgethi into the same mass league as its AGB colleague Mira?); [2013AJ...146..148M](#), proceeding from the surely safe assumption that the three known stars in the  $\alpha$  Her AB system are of the same age (being surely born in a single ISM cloud condensation event), and taking the system age from the MS star that is the secondary in the  $\alpha$  Her unresolved SB (age is more safely determined from an MS star than from an evolved star), and using photometry sensitive to TiO to obtain a fluctuating effective temperature for Rasalgethi (and thereby, via the Stefan-Boltzmann law, fluctuating luminosity and fluctuating radius) via a 55-track grid of evolutionary models (with each track tracing evolution stepwise until the depletion of core helium that marks a star’s arrival on AGB) deduces that the Rasalgethi mass is in the range [2.175  $M_{\odot}$ , 3.250  $M_{\odot}$ ]; the authors remarks that “very few AGB stars have reliable ages and masses”; the luminosity and effective temperature deduced for Rasalgethi agree to within uncertainties with the Rasalgethi interferometry results of [2004A&A...418..675P](#)

¶ in the classification of AAVSO(VSX), Rasalgethi is a “semi-regular late-type giant,” with V-mag. range 2.73–3.60, period 125.6 d (AAVSO(VSX) assessment in 2022: status flag = confirmed variable; variability classification symbol = “SRB”); [2010Ap&SS.328..113M](#) reports “up to seven” pulsation modes, with one period of ~1343 d (a radial pulsation), other periods on the order of ~125 d (and cf. also light curve in Fig. 5 of [2001PASP...113..983P](#)); two typical measurements in V band, from AAVSO database, from one and the same observer, are

$\pi$	Her	17 15.9+36 47	3.16 <sup>†</sup>	1.44	K3 IIab <sup>†</sup>	8.7	-2.2	380	0.027	276	-26 V?	<p>mag. 3.37 (2021 June 27) and mag. 3.07 (2021 Sep. 29)</p> <p>¶ Rasalgethi radius variation, associated with the photometric variation, is assessed in <a href="#">2013AJ...146..148M</a> as 264 R<sub>⊙</sub> min, 303 R<sub>⊙</sub> max;</p> <p><a href="#">en.wikipedia.org/wiki/List_of_largest_stars</a> shows ranking of Rasalgethi in the overall known cosmic population of highly evolved stars</p> <p>¶ <a href="#">1956ApJ...123..210D</a> initiated the successful analysis of mass loss from Rasalgethi (mass loss is to be expected, in an AGB star, and in the case the consequent circumstellar material is so copious as to extend into our line of sight to <math>\alpha</math> Her B); <a href="#">1956ApJ...123..210D</a> proceeds by examining the radial velocity of the Rasalgethi ejecta (as seen in absorption, as very cold gas, in the spectrum of the <math>\alpha</math> Her B brighter component, and (crucially) not sharing in the ~52 d fast SB orbital motion of that component); <a href="#">2007ApJ...658L.103T</a> finds interferometric evidence for an exceptionally violent episode of mass loss, comprising about 10e-6 M<sub>⊙</sub> (cp. Mira, for which we take as usual mass-loss rate ~2.5e-7 M<sub>⊙</sub>/y), with the ejected material (condensing as dust) flowing outward at exceptionally high speeds, possibly ≥ 72 km/s, and with no similarly drastic mass loss until at least the conclusion of the study (~2003); this suggests to us, as the Handbook team, that ongoing photometric study is advisable</p> <p>¶ in the pictorial-atlas tradition, <math>\alpha</math> Her marks the head of hero Hercules (with <math>\beta</math> Her marking his club; for summer-evening observers in the northern hemisphere, the hero is to be visualized inverted, with feet high in the sky, club and head lower; indeed the IAU-official Arabic name “Rasalgethi” derives from the Arabic for “the kneeler’s head”</p> <p>variability just below visual threshold? (3.07–3.16 in V?)</p> <p>AAVSO(VSX), as of 2022 Jul. 30, notes existence of NSV entry and nonexistence of AAVSO observations (AAVSO(VSX) assessment: status flag = suspected variable; no variability classification symbol) (further photometric study advisable?); the gross photometry aside, low-amplitude photometric variations with low-amplitude radial-velocity variations, 613 d, perhaps favour the hypothesis of non-radial pulsation over the competing hypotheses of an undetected low-mass companion and of rotation with starspots</p>
$\delta$	Her Aa <sup>†</sup>	17 16.0+24 49	3.12 <sup>†</sup>	0.08 <sup>†</sup>	A1 Vann	43.4	1.3	75	-0.158	-188	-40 SB <sup>†</sup>	<p>¶ Astron. Alm. (epoch 2021.5) assigns MK type K3 II B: mag. 8.3, 14” (2020) is mere optical companion Sarin <math>\delta</math> Her A, being SB (and also resolved as a binary in interferometry, with angular separation 60 mas; inter-component distance ≥ 1.45 au, period ≥ 335 d; WDS documents just 5 astrometry measurements [0.1”→0.1”, 1978→1989]), is strictly <math>\delta</math> Her Aa,Ab</p> <p>¶ m<sub>v</sub>, B–V values are for <math>\delta</math> Her Aa,Ab combined light</p> <p>¶ <math>\delta</math> Her Aa is a fast rotator (&lt; 9 h)</p> <p>¶ as with <math>\delta</math> Her B, so also <math>\delta</math> Her C and <math>\delta</math> Her D, at respective angular separations 174” (2013) and 192” (2009), are most likely mere optical companions</p> <p>¶ AAVSO(VSX), as of 2022 Jul. 30, notes existence of NSV entry, but flags the <math>\delta</math> Her system as confirmed non-variable (without assigning a variability classification symbol; should the symbol “CST” be assigned?)</p>
$\theta$	Oph A <sup>†</sup>	17 23.5–25 01	3.26 <sup>†</sup>	-0.22	B2 IV	~7.5	-2.4	440	0.025	197	-2 SB <sup>†</sup>	<p>slight var., <math>\beta</math> Cep type, 3.26–3.29 in V, 3.37 h</p> <p>¶ <math>\theta</math> Oph A is classified as an unresolved SB, with period variously stated in secondary literature as 11.44 d (Kaler) and 56.71 d (Wikipedia), and with inter-component distance proposed by Kaler as ~0.25 au;</p> <p><math>\theta</math> Oph B is mag. 6.2, 0.2”→0.1”, 1992→2020, with 10 astrometry measurements documented in WDS; lunar occultations occur, and indeed a lunar occultation event might possibly (Kaler, <a href="#">stars.astro.illinois.edu/sow/thetaoph.html</a>) have split the SB that is <math>\theta</math> Oph A</p> <p>¶ according to AAVSO(VSX) as viewed 2022 March 3,</p>

												and again 2022 Jul. 14, the $\theta$ Oph assemblage presents slight variability of both the $\beta$ Cep type and “slowly pulsating B” type, with period 3.37267 h (AAVSO(VSX) assessment: status flag = confirmed variable; variability classification symbol = “ <b>BCEP+SPB</b> ”)
$\beta$	Ara	17 27.3–55 33	2.84 1.46	K3 Ib–IIa <sup>†</sup>	5	–3.6 600	0.027 199	0				slow rotator (possibly as much as 2.33 y) ¶ high metallicity ¶ AAVSO(VSX) situation in 2022: no status flag (so not confirmed variable, not suspected variable, not confirmed non-variable), and a fortiori no variability classification symbol ¶ not gravitationally bound to $\gamma$ Ara AB
$\gamma$	Ara A	17 27.4–56 24	3.33 <sup>†</sup> –0.14 <sup>†</sup>	B1 Ib	~2.9	–4.4 1100	0.016 182	–3 V				broad lines for Ib ¶ $\gamma$ Ara A is rapid rotator (both “~4.8 d” and “< 2.5 d” have been asserted, and yet rapid rotation is unusual for the (evolved) $\gamma$ Ara A luminosity class) ¶ <a href="#">1997A&amp;A...318..157P</a> finds via <i>IUE</i> spectroscopy that, consistently with this rapid rotation, the stellar wind of $\gamma$ Ara A may be equatorially enhanced (and more generally, that the wind is variable, and is structured with two components, its structure being not typical of stars in this portion of the HR diagram) ¶ $\gamma$ Ara AB is not gravitationally bound to $\beta$ Ara; $\gamma$ Ara B is faint (mag. 10.21: 18”→18”, PA 324°→326°, 1835→2016) ¶ $m_v$ , B–V values are for $\gamma$ Ara A (not for $\gamma$ Ara AB combined light) ¶ AAVSO(VSX) situation in 2022: no status flag (so not confirmed variable, not suspected variable, not confirmed non-variable), and a fortiori no variability classification symbol ¶ E(B–V) = +0.08
$\beta$	Dra A	17 31.0+52 17	2.80 <sup>†</sup> 0.97 <sup>†</sup>	G2 Ib–IIa	8.6	–2.5 380	0.020 308	–20 V				Rastaban in evolutionary terms, $\beta$ Dra A is somewhat unusual, as a yellow more-than-giant (having been a stable core-hydrogen fuser just 0.5 My ago, the star is in transition to being redder, and of still larger radius) ¶ it is also odd that $\beta$ Dra A, while lying in the HR diagram IS, has not been observed to pulsate; however, AAVSO(VSX), as of 2022 Jul. 30, documents Gabriel Cristian Neagu as discovering slight variability, range 2.780–2.794 in V, finding as yet no AAVSO observations (AAVSO(VSX) assessment in 2022: status flag = confirmed variable; variability classification symbol = “ <b>VAR</b> ”); further photometric study advisable? ¶ $m_v$ , B–V values are for $\beta$ Dra AB combined light (where, however, the contribution from $\beta$ Dra B, at mag. –14, is minuscule)
$\upsilon$	Sco	17 32.4–37 19	2.70 –0.23	B2 IV <sup>†</sup>	6	–3.5 600	0.030 185	+8 SB				Lesath although we here give spectral type B, type Be has also been asserted ¶ $\upsilon$ Sco and $\lambda$ Sco are not gravitationally bound (although both belong to the (gravitationally unbound) Sco-Cen OB association, and have as an optical double been called the “Cat’s Eyes”) ¶ AAVSO(VSX) situation in 2022: no status flag (so not confirmed variable, not suspected variable, not confirmed non-variable), and a fortiori no variability classification flag ¶ E(B–V) = +0.02
$\alpha$	Ara A	17 33.7–49 54	2.94 <sup>v†</sup> –0.17 <sup>†</sup>	B2 Vne <sup>†</sup>	12 <sup>†</sup>	–1.7 300 <sup>†</sup>	0.075 206	0 SB				var. of $\lambda$ Eri and $\gamma$ Cas types, 2.73–3.00 in V an instance of the “Be phenomenon,” with (since the star, with its equatorial ejecta, is seen nearly equator-on) “shell” spectrum: <a href="#">2007A&amp;A...464...59M</a> says, “For the first time, we obtain the clear evidence that the [equatorial ejecta] disk is in Keplerian rotation, closing a debate that has continued since the discovery of the first Be star $\gamma$ Cas by Father Secchi”; on the authors’ modelling, $\alpha$ Ara is rotating



													<p>¶ <math>\alpha</math> Oph A is a fast rotator (oblateness has been imaged interferometrically), seen nearly equator-on</p> <p>¶ B is mag. 5.0</p> <p>¶ the SB system <math>\alpha</math> Oph Ab, although tight, has been observed interferometrically, with 17 measurements 1982→2018; the binary system has become better understood with the recent, <a href="#">2011ApJ...726..104H</a>, determination of masses and orbit geometry, through coronagraph and adoptive optics (period 3148.4 d, angular distance at periastron passage ~50 mas; the now-achieved determination of masses in this particular system has implications for astrophysics generally, since it potentially facilitates the refining of numerical models for rapidly rotating hot stars)</p> <p>¶ asteroseismology mission <i>MOST</i> has identified ~50 pulsational modes in <math>\alpha</math> Oph A</p>
$\xi$	Ser Aa <sup>†</sup>	17 38.9–15 25	3.54 <sup>†</sup>	0.26	F0 IIIb <sup>†</sup>	31	1.0	105	0.073	215	-43	SB	<p>hierarchically organized triple system, comprising the resolved (with just one measurement, in 1987: 0.3") single-lined SB that is <math>\xi</math> Ser Aa and <math>\xi</math> Ser Ab, experienced as essentially a point mass by the outlying <math>\xi</math> Ser B; period of the single-lined SB Aa,Ab is 2.29 d; <math>\xi</math> Ser B is faint, at mag. 13.0, with AB period possibly ~15,000 y; there is additionally a faint celestial-sphere neighbour <math>\xi</math> Ser C, at mag. 13.8</p> <p>¶ <math>\xi</math> Ser Aa has been asserted to be very slightly hotter than Garrison's "F0 IIIb," and moreover to be chemically peculiar, being on this (more recent?) determination of MK type A9 IIIp Sr; additionally, AAVSO(VSX), as of 2022 Jul. 30, notes the existence of NSV entry and the nonexistence of AAVSO observations, suggesting <math>\delta</math> Sct variability, suggesting possible V-passband range 3.52–3.54 (AAVSO(VSX) assessment: status flag = suspected variable; variability classification symbol = "DSCT:"); (further photometric study advisable?); Kaler comments at <a href="#">stars.astro.illinois.edu/sow/xiser.html</a>: "the star /.../ remains cryptic"</p> <p>[THIS STAR ONLY IN ONLINE VERSION OF TABLE]</p>
$\theta$	Sco A <sup>†</sup>	17 39.0–43 01	1.86 <sup>†</sup>	0.40	F1 III	~11	-3.0	300	0.006	119	+1		<p>rapid rotator, in the sense that <math>v \sin i</math> is (according to <a href="#">2005yCat.3244...0G</a>) 125.0 km/s; since, however, <math>\theta</math> Sco is a (rapidly evolving) giant, its high <math>v \sin i</math> may correspond to a not-spectacularly short rotation period, of up to 10 d; if, as asserted in literature, <math>\theta</math> Sco truly is a rapid rotator, it will resemble <math>\beta</math> Cas A (a rapid rotator that, strikingly, has evolved beyond the MS)</p> <p>¶ <math>\theta</math> Sco B is catalogued by WDS as mag. 5.4, with <math>\theta</math> Sco AB 6"→6", PA 322°→315°, 1896→1991, but with the caveat (in WDS Notes) that <math>\theta</math> Sco B information may be erroneous (grid-step error in <i>HIPPARCOS</i> data reduction? erroneous work in 1896?)</p> <p>¶ AAVSO(VSX), as of 2022 Jul. 30, notes the existence of NSV entry, fails to find a record of AAVSO observations, and offers possible V-passband range 1.84–1.88 (AAVSO(VSX) assessment: status flag = suspected variable; no variability classification symbol); further photometric study advisable?</p> <p>¶ although it is the Sumerian name Sargas that is IAU-official as of 2016 Aug. 21, <math>\theta</math> Sco, like <math>\kappa</math> Sco, has also been known under the different, not IAU-official, name Girtab (originally applied by the Sumerians to an entire asterism)</p>
$\kappa$	Sco <sup>†</sup>	17 44.1–39 02	2.41 <sup>†</sup>	-0.23	B1.5 III	7	-3.5	480	0.026	193	-14	SB2	<p>slight var., <math>\beta</math> Cep type, 2.41–2.42 in V, 4.80 h</p> <p>¶ since the SB has not yet been resolved, even</p>

$\beta$	Oph	17 44.6 +4 34	2.77 <sup>†</sup>	1.17	K2 III <sup>†</sup>	~39.8	0.8	82	0.165	345	-12 V	<p>interferometrically, WDS cannot yet write “<math>\kappa</math> Sco A” and “<math>\kappa</math> Sco B”; the SB has orbital period 195.65 d, with inter-component distance 1.7 au</p> <p>¶ the SB primary is a rapid rotator (1.9 d), in addition to being a variable of <math>\beta</math> Cep type (0.1998303 d in AAVSO(VSX) as viewed 2021 Jan. 16 and 2022 Mar. 3 (AAVSO(VSX) assessment in 2022: status flag = confirmed variable; variability classification symbol = “BCEP”); <a href="#">1975MNRAS.173..709L</a> gives some photometry, confirming a beat period</p> <p>¶ <math>\kappa</math> Sco, as a single naked-eye object, has (like <math>\theta</math> Sco) been known under the different, not IAU-official, name Girtab (originally applied by the Sumerians to an entire asterism)</p> <p>¶ Astron. Alm. (epoch 2021.5) assigns MK type K2 III CN 0.5</p> <p>¶ <a href="#">1996ApJ...468..391H</a> finds multiple pulsation periods, in behaviour paralleling <math>\alpha</math> Boo (“it may well be that these [two] stars represent a new class of radially and unradially pulsating stars”), and also a possible long period of 142.3 d; the authors suggest that if the latter is real, then although the more likely explanation is a 142.3-d rotation, nevertheless gravitational pull from an unknown exoplanet is conceivable; AAVSO(VSX), as of 2022 Jul. 31, for its part notes the existence of an NSV entry, fails to find a record of AAVSO observations, suggests the V-passband range 2.75–2.77 (AAVSO(VSX) assessment: status flag = suspected variable; no variability classification symbol) (further photometric study advisable?)</p>	Cebralrai
$\mu$	Her <u>Aa</u> <sup>†</sup>	17 47.4+27 43	3.42	0.75	G5 IV	~120.3	3.8	27.1	0.804	201	-16 V	<p>BC: 9.78comb., A–BC 36”, PA 240°→249°, 1781→2015 this stable “double double” is the third-closest quadruple star system to the Sun, and is one of the best-studied double doubles (<a href="#">2016AJ...151..169R</a> consequently writes that <math>\mu</math> Her “serves as an archetype for understanding stellar system formation”): <math>\mu</math> Her Aa,Ab is in tight orbit, and <math>\mu</math> Her BC is in tight orbit, with each of these two pairs experiencing the other as essentially a point mass; <a href="#">2016AJ...151..169R</a> gives Aa,Ab a period of ~100 y, with wide uncertainties, concluding also that Ab is an M-dwarf (and thus more massive than a mere substellar object; Aa is mag. 3.5, Ab mag. 12.7, with this magnitude difference making the astrometry difficult; the Aa,Ab binary has been measured 24 times, 1998→2015 (angular separation 1.8” in 2015); BC has period 43.127 y, B-to-C distance 1.5 au min, 3.6 au max, 2.2 au average (angular separation 0.6” in 2016), abundantly observed (360 measurements, 1854→2016), with B, C of respective mags. 10.2, 10.7; the distance between the Aa,Ab centre of mass and the BC centre of mass is <math>\geq</math> 300 au, with orbital period <math>\geq</math> 3700 y; in contrast with the two pairings Aa,Ab and BC, there is no published orbital solution for the wide and slow pairing AB</p> <p>¶ <math>m_v</math>, B–V values are for <math>\mu</math> Her Aa,Ab combined light; additionally, GCPD gives <math>m_v</math>, B–V values 9.77, 1.50 for <math>\mu</math> Her AB combined light</p> <p>¶ AAVSO(VSX) situation in 2022: no status flag (so not confirmed variable, not suspected variable, not confirmed non-variable), and a fortiori no variability classification type</p> <p>¶ despite having finished core-hydrogen fusion, <math>\mu</math> Her Aa is a fast rotator, and is consequently magnetically active and an X-ray source</p>	
$\iota^1$	Sco A	17 49.2–40 08	3.02	0.51 <sup>†</sup>	F2 Ia	2 <sup>†</sup>	-5.9	2000 <sup>†</sup>	0.006	180	-28 SB	<p>since the SB that is <math>\iota^1</math> Sco A is not as yet resolved (even by interferometry), WDS is not as yet able to write “<math>\iota^1</math> Sco Aa,” <math>\iota^1</math> Sco Ab”</p>	

G	Sco A	17 51.5–37 03	3.20	1.16	K2 III	25.9	0.3	126	0.049	56	+25	<p>¶ a rare instance of a yellow supergiant (dead helium core; the star is now cooling, and is now in transition to the less exotic status of red supergiant)</p> <p>¶ radius estimates vary; CADARS (2001A&amp;A...367..521P) value is ~1.9 au</p> <p>¶ mass loss ~1e-7 M<sub>☉</sub>/y</p> <p>¶ slow rotator (≥ 0.5 y)</p> <p>¶ AAVSO(VSX) situation in 2022: no status flag (so not confirmed variable, not suspected variable, not confirmed non-variable), and a fortiori no variability classification symbol</p> <p>¶ distance and mass are rather uncertain</p> <p>¶ the modest angular distance of ι<sup>1</sup> Sco from ι<sup>2</sup> Sco is the result of a mere line-of-sight coincidence (with ι<sup>2</sup> ~2 times as distant as ι<sup>1</sup>; again by coincidence, not ι<sup>1</sup> alone, but also ι<sup>2</sup>, is a supergiant)</p> <p>HR6630, Fuyue</p> <p>although masses of K giants are in general uncertain, in this particular case the mass is known via <i>WIRE</i> salvage-mission asteroseismology (being determined in 2008ApJ...674L..53S as 1.44 M<sub>☉</sub>, with just a 15% uncertainty)</p> <p>¶ AAVSO(VSX) situation in 2022: no status flag (so not confirmed variable, not suspected variable, not confirmed non-variable), and a fortiori no variability classification flag</p>
γ	Dra A	17 57.2+51 29	2.23	1.52	K5 III <sup>†</sup>	21.1	-1.1	154	0.024	200	-28	<p><b>Eltanin</b></p> <p>in 1728, James Bradley used γ Dra to demonstrate aberration of light (“velocity aberration”); his demonstration strongly confirmed the heliocentric (and thus non-Ptolemaic) kinematics of the Solar System</p> <p>¶ AAVSO, as of 2022 Jul. 31, asserts variability, fails to find a record of AAVSO observations, tentatively categorizes the variability as of a slow irregular type, and suggests V-passband range of 0.01 mag.</p> <p>(AAVSO(VSX) assessment in 2022: status flag = confirmed variable; variability classification symbol = “<b>LB</b>”); further photometric study advisable?</p>
v	Oph +2P <sup>†</sup>	18 00.3 -9 46	3.34 <sup>†</sup>	1.00	G9.5 IIIa <sup>†</sup>	22	0.0	150	0.117	185	+13	<p>¶ Fe is slightly underabundant</p> <p>var. of unknown type? (range ~0.04 in V passband?)</p> <p>AAVSO, as of 2022 Jul. 31, notes the existence of an NSV entry and the existence of 4 AAVSO observations (AAVSO(VSX) assessment: status flag = suspected variable; no variability classification symbol); further photometric study advisable?</p> <p>¶ brown-dwarf companions, not optically resolved (so WDS cannot write “v Oph A,” “v Oph B,” “v Oph C”) with masses ≤ 24× Jupiter and ≤ 27× Jupiter (deuterium fusion begins at a lower mass, 13× Jupiter), periods 530.3 d and 3190 d (Quirrenbach et al. 2011, and additionally 2012PASJ...64..135S; the latter paper suggests formation in circumstellar disk, with subsequent migration, in a scenario reminiscent of planet and exoplanet formation): this is the third star found to be hosting two brown dwarfs</p> <p>¶ slow rotator (≤ 234 d)</p> <p>¶ far-IR variability has been suspected</p> <p>¶ CN underabundant, Fe overabundant</p> <p>¶ Astron. Alm. (epoch 2021.5) assigns MK type G9 IIIa</p>
γ <sup>2</sup>	Sgr	18 07.3–30 25	2.98 <sup>†</sup>	1.00	K0 III <sup>†</sup>	34	0.6	97	0.189	197	+22 SB	<p><b>Alnasl</b></p> <p>metals underabundant</p> <p>¶ AAVSO(VSX), as of 2022 Jul. 31, notes existence of NSV entry but denies variability (AAVSO(VSX) assessment in 2022: status flag = confirmed non-variable; no variability classification symbol (but should the classification system “<b>CST</b>” be assigned?))</p> <p>¶ Astron. Alm. (epoch 2021.5) assigns MK type K0<sup>+</sup> III</p> <p>¶ ε Sgr and the γ<sup>2</sup>-γ<sup>1</sup> Sgr pair serve as pointers to Baade’s Window</p>

$\eta$	Sgr A <sup>†</sup>	18 19.2–36 45	3.11 <sup>†</sup>	1.56 <sup>†</sup>	M3.5 IIIab	22	-0.2 ~146	0.211	218	+1 V?	<p>¶ angular proximity of <math>\gamma^1</math> Sgr (= W Sgr; variable, mag. range 4.28–5.10 in V; ~50', to ~N of <math>\gamma^2</math> Sgr) is a mere line-of-sight coincidence</p> <p>slight irreg. var.: 3.05–3.12; B: 8.00, G8: IV., 3.5" (2016) AAVSO(VSX), as of 2022 Jul. 31, notes existence of 20 AAVSO observations and tentatively assigns a slow irregular variability type (AAVSO(VSX) assessment: status flag = confirmed variable; variability classification symbol = "<b>LB:</b>"); further photometric study advisable?</p> <p>¶ PA 100°→110°, 1879→2016; orbit <math>\geq</math> 1270 y, A-to-B distance <math>\geq</math> 165 au</p> <p>¶ <math>m_v</math>, B–V values are for <math>\eta</math> Sgr AB combined light</p> <p>¶ <math>\eta</math> Sgr A is variously asserted to be on the (very highly evolved) HR diagram AGB or at the tip of the RGB</p> <p>¶ temperature of <math>\eta</math> Sgr A not yet well determined?</p>
$\delta$	Sgr A	18 22.5–29 49	2.70 <sup>†</sup>	1.38	K2.5 IIIa <sup>†</sup>	9	-2.4 350	0.041	128	-20 V?	<p>¶ Astron. Alm. (epoch 2021.5) assigns MK type K2.5 IIIa CN 0.5</p> <p>¶ possibly a weak barium (Ba) star, <math>\delta</math> Sgr A possesses (as expected for a Ba star) a WD companion</p> <p>¶ temperature of <math>\delta</math> Sgr A not yet well determined?</p> <p>¶ AAVSO(VSX), as of 2022 Jul. 31, notes the existence of NSV entry and the nonexistence of AAVSO observations, and states V-passband range simply as "2.7–?" (AAVSO(VSX) assessment: status flag = suspected variable; variability classification symbol = "<b>VAR:</b>"); further photometric study advisable?</p> <p>¶ "Kaus" is Arabic "bow," with Kaus Borealis (<math>\lambda</math> Sgr), Kaus Media (<math>\delta</math> Sgr), and Kaus Australis (<math>\epsilon</math> Sgr) the three delineating stars of the archer's bow; by coincidence, the archer turns out to be aiming rather close both to Baade's Window and (prolonging the line of firing) to the Sgr A* black hole at the galaxy's centre</p>
$\eta$	Ser A	18 22.5 -2 53	3.25 <sup>†</sup>	0.94	K0 III–IV <sup>†</sup>	54	1.9 ~60.5	0.890	218	+9 V?	<p>slight var. of unknown type?</p> <p>AAVSO(VSX), as of 2022 Jul. 31, notes the existence of NSV entry and the nonexistence of AAVSO observations, states range in the near-IR K band as 0.91–1.28 (AAVSO(VSX) assessment: status flag = suspected variable; no variability classification symbol); further photometric study advisable?</p> <p>¶ slow rotator (but <math>\leq</math> 1.9 y)</p> <p>¶ high velocity relative to Sun suggests that <math>\eta</math> Ser is an interloper (born outside the galactic thin disk? consistently with this conjecture, Fe is underabundant)</p>
$\epsilon$	Sgr A	18 25.7–34 22	1.84	-0.03	A0 II n (shell?) <sup>†</sup>	23	-1.4 ~143	0.130	198	-15	<p><b>Kaus Australis<sup>†</sup></b></p> <p>fast rotator (consistent with shell-star classification); as might be predicted for a fast rotator, a magnetic field, and also X-ray emission, have been detected</p> <p>¶ Astron. Alm. (epoch 2021.5) assigns MK type "A0 II n (shell)"</p> <p>¶ has been classified as a <math>\lambda</math> Boo star, apparently in error</p> <p>¶ IR excess indicates debris disk (possibly also detected in polarimetry), at average separation 155 au; and yet a companion (other than the WDS-documented celestial-sphere neighbours <math>\epsilon</math> Sgr B, <math>\epsilon</math> Sgr C, <math>\epsilon</math> Sgr D) is also asserted, surprisingly present within this debris-disk radius</p> <p>¶ <math>\epsilon</math> Sgr B,C,D are at mags. 14.3, 8.4, 9.0 respectively; AC astrometry is 2.2"→2.4", PA 146°→142°, 1992→1999; AD astrometry (as a pair at wide angular separation) is 858"→858", PA 36°→36°, 1980→1999</p> <p>¶ AAVSO(VSX) situation in 2022: no status flag (so not confirmed variable, not suspected variable, not confirmed non-variable), and a fortiori no variability classification symbol</p>
$\alpha$	Tel	18 28.7–45 57	3.51 <sup>†</sup>	-0.17	B3 IV <sup>†</sup>	12	-1.2 280	0.056	198	0 V?	<p><a href="http://stars.astro.illinois.edu/sow/alphatel.html">stars.astro.illinois.edu/sow/alphatel.html</a></p> <p>remarks that MK luminosity class IV notwithstanding,</p>



															<p><math>\alpha</math> Tel is still on the astrophysical (as opposed to the MK-phenomenological) Main Sequence (in other words, is still fusing core hydrogen)</p> <p>¶ said in <a href="#">2005ApJS...158..193S</a> to be among the (rare) He-rich stars; these authors list <math>\alpha</math> Tel as a candidate-and-unconfirmed <math>\beta</math> Cep variable, and say they suspect it is a variable in the slowly pulsating B-star class; although <math>\alpha</math> Tel has <i>HIPPARCOS</i> microvariability (0.909 d), it is absent from the AAVSO(VSX) database as viewed 2022 March 03, 2022 Jul. 31 (since no status flag, not confirmed variable, not suspected variable, not confirmed non-variable; a fortiori no variability classification symbol)</p> <p>[THIS STAR ONLY IN ONLINE VERSION OF TABLE]</p>
$\lambda$	Sgr A	18 29.4–25 24	2.82	1.04	K1 IIIb	~41.7	0.9	78	0.191	194	-43 V?				<p>Kaus Borealis<sup>†</sup></p> <p>modest X-ray emission indicates some magnetic activity (not usual in a duly evolved, stable core-He-fusing, HR diagram “clump star”)</p> <p>¶ <math>\lambda</math> Sgr B is mag. 9.9; AB 101”→82”, PA 188°→184°, 1911→2019</p> <p>¶ lunar occultations are possible, planetary occultations possible yet rare; most recent planetary occultation was by Venus, on 1984 Nov. 19</p> <p>¶ AAVSO(VSX) situation in 2022: no status flag (so not confirmed variable, not suspected variable, not confirmed non-variable), and a fortiori no variability classification symbol</p> <p>¶ unusual in occupying fully three roles in the Western pictorial traditions: as northernmost star of the Archer’s Bow, as westernmost (handle-tip) star of the Little Milk Dipper, and as uppermost (lid-knob) star in the Teapot</p>
$\alpha$	Lyr A	18 37.7+38 48	0.03 <sup>†</sup>	0.00	A0 Va <sup>†</sup>	130	0.6	25.0	0.350	35	-14 V				<p>pole-on rapid rotator with circumstellar disk <b>Vega</b></p> <p>¶ pole-on rotators are useful for asteroseismology, since all but the axisymmetric modes (whether radially symmetric or radially not symmetric) are helpfully rendered invisible to photometry (and in a rather analogous way, equator-on rotators are also useful through suppression of some modes);</p> <p><math>\alpha</math> Lyr pole-on orientation represents an extreme on a continuum whose other extreme is represented by the equator-on rapid rotator <math>\alpha</math> Leo A (Regulus); the <math>\alpha</math> Lyr A rapid rotation (<a href="#">2015A&amp;A...577A..64B</a>) now confidently asserts 0.68 d) yields oblate spheroid shape (here as with <math>\alpha</math> Leo A); it is this, with consequent latitude-varying photosphere (severe temperature and luminosity gradients along the arcs of photospheric longitude, with equator coolest and darkest), rather than any evolution beyond core-hydrogen-fusion stage, that explains the anomalously high luminosity (<math>\alpha</math> Lyr A is in MK luminosity class Va, rather than in the slightly dimmer V class that would be observed if its orientation was equator-on)</p> <p>¶ <math>\alpha</math> Lyr A is now known to harbour all three of the classical circumstellar-dust regimes (~1500 K, near-IR; ~120-170 K, mid-IR, as an analogue of our own zodiacal dust; and ~50 K, far-IR, as an analogue of our own Kuiper Belt: for regimes overview without specific reference to <math>\alpha</math> Lyr, cf <a href="#">2013ApJ...763..118S</a>, section 1):</p>

[2013ApJ...763..118S](#) is the paper announcing discovery of the second of these around  $\alpha$  Lyr (with sections 5.1 and 5.2, respectively, summarizing previous  $\alpha$  Lyr A work on the first and third of the three regimes): a question of recent interest is the origin of the  $\alpha$  Lyr A exozodiacal (warm-regime, mid-IR) dust (episode analogous to our own planetary system's Late Heavy Bombardment? or, rather, some steady-state replenishment mechanism?); efforts at detecting exoplanet(s) to account for the complex inferred, and indeed in some wavelengths also now directly imaged, disk structure have not yet succeeded ¶ [2007ASPC...364...305G](#), reviewing the history of  $\alpha$  Lyr A photometry, considers modest variability likely, the historical use of  $\alpha$  Lyr A as a photometric standard notwithstanding [and indeed  $\alpha$  Lyr A is described at AAVSO(VSX) as a low-amplitude  $\delta$  Sct variable, in the now-obsolete AAVSO(VSX) "DSCTC" classification bin, fluctuating between  $-0.02$  and  $+0.07$  in V, with period 5 h (AAVSO(VSX) assessment in 2022: status flag = confirmed variable; variability classification symbol = "DSCTC" )] ¶ [2010A&A...523A..41P](#), with [2014A&A...568C...2P](#) corrigendum, is a recent discussion of  $\alpha$  Lyr A magnetism (the authors note that  $\alpha$  Lyr A "may well be the first confirmed member of a much larger, as yet unexplored, class of weakly-magnetic stars now investigatable with the current generation of stellar spectropolarimeters"; for origin, they somewhat favour dynamo over fossil, and radiative dynamo over core dynamo): consistently with magnetism, [2015A&A...577A..64B](#) finds, via line-profile variations, multiple (bright, not dark) star spots, in some undetermined complex pattern (authors comment that this is "first strong evidence that standard A-type stars can show surface structure"); [2015A&A...577A..64B](#) is additionally one of several papers summarizing recent work on an interrelated complex of  $\alpha$  Lyr A themes, comprising (in addition to magnetism) rotation, spots, photovariability, and pulsation ¶ in [2010ApJ...725.2401F](#), Fig. 8 with its accompanying discussion summarizes studies on elemental abundances (important because  $\alpha$  Lyr A, as a rather "normal" star for MK temperature type A, might serve as a benchmark for appraising chemically peculiar A stars) ¶ WDS documents an exceptionally large number of celestial-sphere neighbours for  $\alpha$  Lyr A (B,C,D,E,F,G,H, Ia,Ib,J,Ka,Kb,L,M,N,O,P,Q,R,S,T); all of these are faint apart from B and E (each at mag. 9.5, and each known to be not gravitationally bound to A): AB  $37'' \rightarrow 84''$ , PA  $117^\circ \rightarrow 184^\circ$ , 1781  $\rightarrow$  2017; AE  $150'' \rightarrow 87''$ , PA  $40^\circ \rightarrow 39^\circ$ , 1831  $\rightarrow$  2017 ¶ E(B-V) = 0.00 ¶ E(B-V) = 0.00 apparent duplicity now discounted

$\beta$  Lyr Aa1<sup>†</sup> 18 50.9+33 23 3.42v<sup>†</sup> 0.01<sup>†</sup> B7 Vpe (shell)<sup>†</sup> ~3.4<sup>†</sup> -3.8 ~960<sup>†</sup> 0.004 152 -19 SB

(erroneous lunar-occultation observation)  
 ¶ [stars.astro.illinois.edu/sow/phisgr.html](http://stars.astro.illinois.edu/sow/phisgr.html)  
 discusses some difficulties in physical modelling (if pole-on rotator, then there will be troublesome temperature and luminosity gradients along the arcs of photospheric longitude)  
 ¶ AAVSO(VSX) situation in 2022: no status flag (so not confirmed variable, not suspected variable, not confirmed non-variable), and a fortiori no variability classification symbol  
 eclipsing, 3.30–4.35 in V, 13 d Sheliak  
 period is increasing at constant rate of ~19 s/y;  
 orbital solution has been published for  $\beta$  Lyr Aa1,  $\beta$  Lyr Aa2 (although not for the wider binary  $\beta$  Lyr Aa,  $\beta$  Lyr Ab, and not for the still wider binary  $\beta$  Lyr A,  $\beta$  Lyr B);  
 this orbit is seen nearly edge-on;  
 prototype of the  $\beta$  Lyr class of eclipsing systems (but has also been assigned to the new class of “W Ser stars”: [1980AJUS...88..251P](https://ui.adsabs.org/1980AJUS...88..251P));  
 AAVSO supplies information both via VSX database (showing, e.g. the high-precision recent determinations of period; assessment in 2022 is as follows:  
 status flag = confirmed variable;  
 variability type = “DPV:EB”) and via [www.aavso.org/vsots\\_betalyr](http://www.aavso.org/vsots_betalyr)  
 (a detailed astrophysics discussion, with bibliography):  
 alternating deep and shallow visible-light minima, with the object eclipsed in the deep minima (the “donor”) a Roche-lobe-filling giant, currently ~3  $M_{\odot}$  and diminishing, and the object eclipsed in the shallow minima (the “gainer”) embedded in a thick accretion disk, currently ~13  $M_{\odot}$  and increasing; mass transfer is copious (~2e–5  $M_{\odot}$ /y); this disk renders the gainer dim, and its eclipses consequently shallow, even though the (presently dim) gainer is (now, at this rather late stage in mass transfer) already ~4 times more massive than the (bright) donor (cf [1963ApJ...138..342H](https://ui.adsabs.org/1963ApJ...138..342H));  
 further, instabilities in the accretion disk, from which ~20% of the light comes, make the light curve liable to vary slightly from cycle to cycle; the presently dim gainer is destined to be first (1) brightening, and spun up by conservation of angular momentum, as its obscuring accretion disk disappears by being dumped down into photosphere, and then (2) to become a slower rotator, tidally locked with the secondary; at stage “(1),” the system will be a so-called “Rapidly Rotating Algol,” at stage “(2),” on the other hand, the system will be simply a “classical Algol”  
 ¶ [2008ApJ...684L..95Z](https://ui.adsabs.org/2008ApJ...684L..95Z) presents the first (CHARA-interferometric) binary-resolving imaging, achieving resolution ~0.5 mas or ~0.7 mas (and for the first time in astrophysics deduces a  $\beta$  Lyr Aa1,Aa2 astrometric orbit); the bright low-mass donor, and the presently dim high-mass gainer, are evident, corroborating the overall conception of

[1963ApJ...138..342H](https://ui.adsabs.org/1963ApJ...138..342H) ; [2008ApJ...684L..95Z](https://ui.adsabs.org/2008ApJ...684L..95Z)

discusses also polar outflow jets on the gainer (these do not alter the essential situation: for the gainer, equatorial gain exceeds polar loss), and deduces a distance to  $\pm 15\%$  (a distance consistent-to-within-uncertainties with the *HIPPARCOS* distance)  
 ¶ [2012ApJ...750...59L](https://ui.adsabs.org/2012ApJ...750...59L) discusses possible hot spot at edge of accretion disk, on the basis of spectropolarimetry (and [2013MNRAS.432..799M](https://ui.adsabs.org/2013MNRAS.432..799M) has modelling that provides for hot spot, and additionally for a bright spot, on the accretion disk)  
 ¶ some observations have been made in radio and (a regime especially relevant to hot-spot studies) X-ray  
 ¶ strictly speaking, this is a hierarchical system, Ab experiencing the binary that is Aa1+Aa2 as essentially a point mass; for the Aa+Ab pairing, and for

												possibility of further pairings (AB, AC, ... , Be, ...), cf WDS and (a source that reports inter alia <i>Gaia</i> ) <a href="https://en.wikipedia.org/wiki/Beta_Lyrae">en.wikipedia.org/wiki/Beta_Lyrae</a> ¶ m <sub>v</sub> , B–V values are for β Lyr Aa1,Aa2,Ab combined light ¶ although we here, following Garrison, assign a rather straightforward spectral type, this should be taken only as a starting point: cf, eg., <a href="#">2000A&amp;A...353.1009B</a> , which lists six systems of spectral lines, while repeating an old O. Struve warning that spectrum involves circumstellar matter ¶ Kaler comments in <a href="https://stars.astro.illinois.edu/sow/sheliak.html">stars.astro.illinois.edu/sow/sheliak.html</a> “one of the most confusing, heavily studied, and important stars of the nighttime sky” ¶ the rather long period, with the large magnitude swing, and the readily discoverable difference in depths of the alternating minima, make this object a suitable binoculars-or-naked-eye photometry project (using γ Lyr A as a comparison) even from locations suffering rather frequent cloud <b>[THIS STAR ONLY IN ONLINE VERSION OF TABLE]</b>
σ	Sgr Aa	18 56.7–26 16	2.08	−0.21	B3 IV	14	−2.2	230	0.056	164	−11V	<b>Nunki</b> fast rotator ¶ the σ Sgr Aa,Ab duplicity was discovered, with suggestion that mags. are roughly equal, but with the system not resolved, through the Narrabri intensity interferometer ( <a href="#">1974MNRAS.167..121H</a> ); WDS documents a single measurement of Aa,Ab (in 1991, with separation found to be ~12 mas), through aperture-masking interferometry at AAT ( <a href="#">1994A&amp;A...290.340B</a> ); σ Sgr B, at mag. 9.95, is known to be not gravitationally bound to σ Sgr Aa,Ab; AB 309″→349″, PA 244°→239°, 1837→1999 ¶ m <sub>v</sub> , B–V values are for σ Sgr Aa,Ab combined light ¶ AAVSO(VSX) situation in 2022: no status flag (so not confirmed variable, not suspected variable, not confirmed non-variable), and a fortiori no variability classification symbol ¶ lunar occultations are possible, and planetary occultation possible-yet-rare (most recently Venus, 1981 Nov. 17) ¶ E(B–V) = +0.02
ξ <sup>2</sup>	Sgr	18 59.1–21 04	3.51	1.18	K1 III	9	−1.7	400	0.034	113	−20	occultations (at any rate lunar) are possible ¶ AAVSO(VSX) situation in 2022: no status flag (so not confirmed variable, not suspected variable, not confirmed non-variable), and a fortiori no variability classification symbol ¶ the angular proximity of ξ <sup>1</sup> Sgr is a mere line-of-sight coincidence <b>[THIS STAR ONLY IN ONLINE VERSION OF TABLE]</b>
γ	Lyr A	18 59.8+32 43	3.25	−0.05	B9 II <sup>†</sup>	5	−3.1	600	0.003	290	−21 V <sup>†</sup>	possible slight var. (3.23–3.26 in V passband?) Sulafat AAVSO(VSX), as of 2022 Jul. 31, notes existence of NSV entry and nonexistence of AAVSO observations (AAVSO(VSX) assessment in 2022: status flag = suspected variable; no variability classification symbol); further photometric study advisable? ¶ has been both asserted and denied to be SB ¶ <a href="#">2001A&amp;A...371.1078A</a> reports many metals underabundant
ζ	Sgr <u>AB</u> <sup>†</sup>	19 04.1–29 51	2.60	0.08	A2 IV–V + A4:V:	37	0.4	90	n.a.	n.a.	+22 SB	A: 3.3; B: 3.5, 0.3″ (2020), 21.1 y, compos. spectrum Ascella PA is 223° in 2020; the system is astrometrically well observed, with 210 astrometry measurements 1867→2020; A-to-B distance is 10.6 au min, 16.1 au max, average 13.4 au ¶ m <sub>v</sub> , B–V values are for ζ Sgr AB combined light ¶ under IAU rules, “Ascella” designates ζ Sgr A, not ζ Sgr B ¶ <a href="https://stars.astro.illinois.edu/sow/ascella.html">stars.astro.illinois.edu/sow/ascella.html</a> discusses uncertainty in masses, remarks that temperatures are not yet directly measured ¶ AAVSO(VSX) situation in 2022: no status flag (so not confirmed variable, not suspected variable, not confirmed non-variable)

ζ	Aql A <sup>†</sup>	19 06.5+13 54	2.98 <sup>†</sup>	0.01 <sup>†</sup>	A0 Vann	~39.3	1.0	83	0.096	184	-25 SB	<p>¶ Sgr C (17.6" in 2013) is probably a mere optical companion</p> <p>among the most rapidly rotating stars known (period 16 h)</p> <p>¶ since the SB is not as yet resolved (not even interferometrically), WDS is as yet unable to write "ζ Aql Aa," "ζ Aql Ab"; in the faint angular-proximity grouping ζ Aql B,C, D, E, ζ Aql B, at mag. 12.0, has been asserted to be a gravitationally bound companion of the ζ Aql A SB pair (5"→7", PA 60°→46°, 1874→2016; A-to-B distance is ≥ 125 au, period ≥ 800 y); also, the exceedingly faint (mag. 16.2) ζ Aql E may be gravitationally bound</p> <p>¶ m<sub>v</sub>, B-V values are for ζ Aql AB combined light (where, however, the contribution of faint ζ Aql B is minuscule)</p> <p>¶ ζ Aql assemblage may harbour slight variability: AAVSO(VSX), as of 2022 Jul. 31, notes existence of NSV entry and nonexistence of AAVSO observations, suggests possible V-passband range 2.98–2.99 (AAVSO(VSX) assessment in 2022: status flag = suspected variable; no variability classification symbol); further photometric study advisable?</p> <p>¶ <a href="#">2008A&amp;A...487.1041A</a> reports near-IR excess around ζ Aql A, and suggests that an unseen close companion is a more likely source than a close-in hot debris disk</p>	Okab
λ	Aql	19 07.5 -4 51	3.43	-0.10	B9 Vnp (kB7HeA0) <sup>†</sup> 26		0.5	120	0.093	192	-12 V <sup>†</sup>	<p>possibly SB</p> <p>¶ rapid rotator (&lt; 21h)</p> <p>¶ suspected chemically anomalous (metals-weak, in λ Boo class); Astron. Alm. (epoch 2021.5) assigns MK type "A0 IVp (wk 4481)"</p> <p>¶ AAVSO(VSX) situation in 2022: no status flag (so not confirmed variable, not suspected variable, not confirmed non-variable), and a fortiori no variability classification symbol</p>	
τ	Sgr	19 08.4-27 38	3.32	1.18	K1.5 IIIb <sup>†</sup>	27	0.5	120	0.255 <sup>†</sup>	191	+45 <sup>†</sup>	<p>possibly SB</p> <p>¶ high velocity relative to Sun suggests origin outside galactic thin disk; underabundance of metals is consistent with this conjecture</p> <p>¶ slow rotator (≤ 270 d)</p> <p>¶ AAVSO(VSX) situation in 2022: no status flag (so not confirmed variable, not suspected variable, not confirmed non-variable), and a fortiori no variability classification symbol</p>	
π	Sgr AB <sup>†</sup>	19 11.2-20 59	2.89 <sup>†</sup>	0.36 <sup>†</sup>	F2 II-III + n.a.	6	-3.1	500	0.036	182	-10	<p>possible var.? (1.75-2.38 in near-IR K band?)</p> <p>Albaldah</p> <p>AAVSO(VSX), as of 2022 Jul. 31, notes existence of NSV entry and nonexistence of AAVSO observations</p> <p>(AAVSO(VSX) assessment: status flag = suspected variable; no variability classification symbol); further photometric study advisable?</p> <p>¶ in HR-diagram terms, π Sgr A lies on blue edge of IS, without being presently observed to pulsate</p> <p>¶ tight triple system, seldom successfully resolved</p> <p>π Sgr A is mag. 3.6, π Sgr B is mag. 3.6, and π Sgr C is mag. 6.0; AB 0.1"→0.1", PA 152°→179°, 1936→1989 (only 5 astrometry measurements), with A-to-B distance ≥ 13 au, AB orbit ≥ 15 y; AB,C 0.4"→0.3", PA 122°→136°, 1936→1939 (only 3 astrometry measurements), with AB-to-C distance ≥ 40 au, orbit ≥ 100 y; under IAU rules, "Albaldah" designates π Sgr A, not π Sgr B or π Sgr C</p> <p>¶ m<sub>v</sub>, B-V values are for π Sgr ABC combined light</p> <p>¶ lunar occultations of ABC are possible, planetary occultations possible yet rare (next by Venus, 2035 Feb. 17)</p>	
δ	Dra A	19 12.6+67 42	3.07	0.99	G9 III	33.5	0.7	97	0.133	46	+25 V	<p>AAVSO(VSX) situation in 2022: no status flag (so not confirmed variable, not suspected variable, not confirmed non-variable), and a fortiori</p>	Altai

$\delta$	Aql <u>Aa</u>	19 26.7 +3 10	3.36 <sup>†</sup>	0.32 <sup>†</sup>	F2 IV <sup>†</sup>	64	2.4	51	0.268	72	-30 SB <sup>†</sup>	no variability classification symbol
												fast rotator (> 0.9 d) ¶ Astron. Alm. (epoch 2021.5) assigns MK type F2 IV–V ¶ the SB that is $\delta$ Aql Aa,Ab has been observed just once in successful astrometry (speckle interferometry, 1979, with 1.9 m telescope at OHP, with angular separation found to be $132 \pm 10$ mas); the Aa,Ab duplicity was discovered astrometrically, through periodic perturbation in proper motion of $\delta$ Aql Aa, in 1929 (Alden, as reported in <a href="#">1936AJ....45..193A</a> ); an astrometric-spectroscopic orbit is offered in <a href="#">1989AJ....98..686K</a> , with period $3.426 \pm 0.006$ y, $e = 0.36 \pm 0.07$ (rotational broadening makes the spectroscopy difficult; also, this paper deduces a large magnitude difference between $\delta$ Aql Aa and $\delta$ Aql Ab, diminishing the prospects for easy interferometry) ¶ <a href="#">stars.astro.illinois.edu/sow/deltaaql.html</a> discusses points of uncertainty (incl. the just-mentioned binarity, and possible $\delta$ Sct variability; although in 2018 $\delta$ Aql was not in the AAVSO(VSX) database, as of 2022 Jul. 31 AAVSO(VSX) flags the $\delta$ Aql assemblage as a confirmed variable, while noting the nonexistence of AAVSO observations, and gives V-passband range 3.36–3.37 and period 1.04524 d (AAVSO(VSX) assessment in 2022: status flag = confirmed variable; variability classification symbol = “ <b>GDOR:</b> ”); further photometric study advisable? ¶ $m_v$ , B–V values are for $\delta$ Aql Aa,Ab combined light B: 5.13, 35”; Aa, Ab, Ac $\leq 0.3''$ (2020) Albireo B is of B–V colour $-0.09$ , MK type B9.5 Ve ¶ if AB is true binary, orbit is possibly $\geq 100\,000$ y; the competing mere-optical-companions thesis is argued by Bob King in <i>Sky &amp; Telescope</i> 2016 Sep. 21; same conclusion is reached in 2018 by P. Plait at <a href="#">www.syfy.com/syfywire/long-standing-astronomical-mystery-solved-albireo-is-not-a-binary-star</a> , on strength of fresh <i>Gaia</i> data (which yield for $\beta$ Cyg B $\pi = 8.4$ mas $\pm 2\%$ , implying distance for $\beta$ Cyg B, to two significant figures, 390 ly; however, further analysis is needed, since astrometry of $\beta$ Cyg A is potentially perturbed by the multiplicity of A <a href="#">en.wikipedia.org/wiki/Albireo</a> recaps literature, with some reference to recent interferometry); WDS takes the firm view that AB is not a true binary, citing discrepancy in its accepted parallaxes, while also noting the binarity defence arguments of R. Griffin in <a href="#">1999JRASC..93..208G</a> (Griffin’s assumed parallaxes are for their part not discrepant; in arguing for binarity, Griffin also (a) cites the statistical improbability of two such bright objects appearing on the celestial sphere at so tight an angular separation if they are not gravitationally bound, and (b) notes the possibility of shared proper motion [with, he stresses, proper motion observations rendered difficult by possible astrometric wobble in the photocentre of the Aa,Ac binary]) ¶ Aa,Ab,Ac are at mags. $\sim 3.4$ , $\sim 5.0$ , $\sim 5.2$ respectively; $m_v$ , B–V values are for $\beta$ Cyg Aa,Ab,Ac combined light ¶ the Aa,Ac binary has been measured in astrometry 62 times, 1976–2021 (typical angular separation $\sim 0.3''$ ); <a href="#">2008AN....329..54S</a> offers a preliminary Aa,Ac orbit (with $e \sim 0.3$ ; since period is $\sim 210$ y, the orbit will remain only imperfectly known over coming decades); the Aa,Ab binary has been measured in astrometry just twice (1978,1995, with the 1978 angular separation $0.1''$ ) ¶ our values, for $\beta$ Cyg A, of $\pi = 10$ mas (strictly, $9.5$ mas $\pm 6.0\%$ ), with D consequently computed to two significant figures as 330 ly, are taken uncritically from <i>Gaia</i> $\sim 2018$ , rather than (as in our previous Handbook editions) from <i>HIPPARCOS</i> ;
$\beta$	Cyg <u>Aa</u> <sup>†</sup>	19 31.7+28 01	3.08 <sup>†</sup>	1.13 <sup>†</sup>	K3 II <sup>†</sup>	10 <sup>†</sup>	-2.3	330 <sup>†</sup>	0.009	229	-24 V	

												we do not here attempt a critical investigation of uncertainties
												¶ $\beta$ Cyg B is a fast rotator ( $< 0.6$ d), and consistently with this is in emission (as “Be,” rather than plain “B”: being very evolved, this star is not, however, an instance of the “Be phenomenon” as discussed in the final subsection of our accompanying essay)
												¶ $\beta$ Cyg B is not now thought to be a binary (WDS note, and WDS “X” flag)
												¶ AAVSO(VSX) assessment (2022) of $\beta$ Cyg system: status flag = suspected variable; no variability classification symbol
												¶ the name “Albireo,” colloquially associated with the AB pairing as visible in a small telescope, applies under IAU rules only to $\beta$ Cyg Aa
$\delta$	Cyg $\Delta^\dagger$	19 45.7+45 11	2.87 <sup>†</sup> -0.03	B9.5 III	20	-0.7	160	-0.066	~42	-20	SB	B 6.3, F1 V; 1.9”→2.7”, PA 41°→213°, 1826→2020Fawaris orbit 780 y; separation 84 au min, 230 au max, 157 au average, period 780 y
												¶ $m_v$ , B-V values are for $\delta$ Cyg AB combined light
												¶ $\delta$ Cyg A is a rapid rotator
												¶ $\delta$ Cyg C is gravitationally bound to the AB pair: mag. 12, angular distance (2017) 62.5”, PA (for AC): 66°→67°, 1913→2017
												¶ variability has been suspected both in A and in B; nevertheless, AAVSO(VSX), while noting the existence of NSV entry, asserts constant light (AAVSO(VSX) assessment in 2022: status flag = confirmed non-variable; variability classification symbol = “CST”)
												¶ E(B-V)=+0.05
$\gamma$	Aql A	19 47.4+10 40	2.72 <sup>†</sup> 1.51	K3 II <sup>†</sup>	~8.3	-2.7	390	0.017	100	-2	V	possible variability (unknown type) Tarazed
												AAVSO(VSX), as of 2022 Jul. 31, notes the existence of NSV entry and of 2 AAVSO observations, writes for V-passband range “2.72-?” (AAVSO(VSX) assessment: status flag = suspected variable; variability classification symbol = “VAR:”); further photometric study advisable?
												¶ radius ~0.5 au
												¶ a rare instance of a “hybrid” star (possessing a (hot) corona, like our Sun’s, and yet also emitting the cool high-mass wind typical in an evolved star)
$\chi$	Cyg A	19 51.5?32 58	to 3.3v <sup>†</sup> 1.92	S6,2e-S10,4e(MSe)	6	—	~500	3	298	+1.6		Mira-type variable (in 2022, was mag. 4.6 to 13.0)
												¶ we take MK type range from AAVSO(VSX), which follows <a href="#">1974ApJS...28..271</a> ; the unusual type “S” is in temperature terms similar to MK type “M,” but signifies anomalous composition, with carbon and oxygen both abundant, and their abundances roughly equal; the comma notation, as explained at <a href="https://en.wikipedia.org/wiki/S-type_star">en.wikipedia.org/wiki/S-type_star</a> , seeks to document both temperature and composition; Astron. Alm. and <a href="#">1980ApJS...43..379K</a> for their part do not attempt to give an MK range, assigning simply the one S-type “S6+/1e” (where “1e” is not a luminosity class, but is a carbon-versus-oxygen abundance index)
												¶ AAVSO(VSX) gives the V-band range as 3.3–14.2 and the period as 408.05 d; the two brightest visually estimated maxima since 2000 Jan. 01 were at some point in 2006 August (with AAVSO recording a visual estimate of ~3.1 on 2006 Aug. 13; there is no V-band measurement from this time in the AAVSO archive) and in 2013 May (with AAVSO recording a visual estimate of ~3.3; there is additionally a V-band measurement from 2008 May 13, of mag. 3.77 with uncertainty 0.04); the most recent maximum was in 2022 April (visual estimate ~4.4, 2022 April 21, in reasonable agreement with V-band measurement 4.69 from 2022 April 22), with the next

$\alpha$	Aql A	19 51.9 +8 56	0.76 <sup>†</sup>	0.22	A7 Vnn	195	2.2	16.7	0.660	54	-26	<p>maximum consequently being due in 2023 June (AAVSO(VSX) assessment 2023 Feb. 1: status flag = confirmed variable; variability classification symbol = “M”)</p> <p>¶ we take D, and by inference <math>\pi</math>, from <a href="#">2009ApJ...707..632L</a>, (which, modifying the Baade-Wesselink method, compares interferometrically measured angular size fluctuations against spectroscopically measured radial velocities); this method agrees to within uncertainties with the Hipparcos 2007 determination (the uncertainty in both cases being large, on the order of 20% or 25%).</p> <p>¶ we compute the <math>M_V</math> range from the <math>m_V</math> range, using the <a href="#">2009ApJ...707..632L</a> value of D.</p> <p>¶ AAVSO(VSX) assessment ~2022: status flag = confirmed variable; classification symbol = M.</p> <p>[THIS STAR ONLY IN ONLINE VERSION OF TABLE]</p> <p>rapid rotator (~7 h or ~8 h, latitude-dependent) <b>Altair</b> the first MS star, other than the Sun, to yield a measurement of photospheric oblateness (<a href="#">2001ApJ...559.1155V</a>); <a href="#">2007Sci...317..342M</a> announces CHARA imaging with angular resolution ~0.65 mas (the first direct imaging of any MS star other than the Sun; <a href="#">news.bbc.co.uk/2/hi/science/nature/6709345.stm</a> is a news writeup; <a href="#">2007Sci...317..342M</a> shows oblate rotation-flattened photosphere, brighter at poles than at equator)</p> <p>¶ found in <a href="#">2005ApJ...619.1072B</a>, via <i>WIRE</i> salvage mission, to be a slight variable of the <math>\delta</math> Sct type, a classification now followed by AAVSO(VSX) (which indicates a fluctuation of 0.004 mag in V, period 91.32 minutes (AAVSO(VSX) assessment in 2022: status flag = confirmed variable; variability classification symbol = “DSCT”); <i>WIRE</i> makes this the brightest known <math>\delta</math> Sct variable; second-brightest is <math>\beta</math> Cas); the <a href="#">2005ApJ...619.1072B</a> authors suggest that many <math>\delta</math> Sct variables, as residents of the IS, may be oscillating at such low amplitudes as to evade detection except by such sensitive facilities as <i>WIRE</i> (their suggestion helps relieve a longstanding astrophysical puzzlement over IS residents that appear, inexplicably, not to be pulsating)</p> <p>¶ drawing on interferometry, spectroscopy, and the <a href="#">2005ApJ...619.1072B</a> <math>\delta</math> Sct asteroseismology, <a href="#">2020A&amp;A...633A..78B</a>, while conceding a failure of uniqueness, and consequently conceding the need for further spectroscopy, offers a physical model that takes account of the rapid rotation (by assuming mere cylindrical symmetry, and not the outright spherical symmetry that would be appropriate in the modelling of a slow rotator); Table 5 of the paper summarizes its results, comparing them against earlier modelling; the paper finds a typical core rotation period ~0.6 of the rotation period of the photosphere, and with only modest latitude variation (shearing) in the rotation period in the photosphere (with middle altitudes ~7.7 h, equator ~7.8 h, immediate vicinity of poles ~8.1 h); the paper deduces a value for core metallicity that makes <math>\alpha</math> Aql A young, aged only ~100 My (but some other recent literature proposes instead ~1.2 Gy; both suggested ages are consistent with the failure of <math>\alpha</math> Aql A to have progressed significantly off the MS); the paper ascribes to <math>\alpha</math> Aql A a remarkable variation in envelope temperature, with the envelope convective (because cooler) at low latitudes and radiative (because hotter) at high latitudes (a similar latitude-governed bifurcation in envelope characteristics is believed present in the rapid rotator <math>\alpha</math> Cep A [Alderamin]); consistently with this latitude-dependent temperature</p>
----------	-------	---------------	-------------------	------	--------	-----	-----	------	-------	----	-----	--



$\eta$	Aql A <sup>†</sup>	19 53.7 +1 04	3.65v <sup>†</sup> 0.68	F6–G1 Ib	2	–4.3 1000	0.011 140	–15 SB	<p>variation, <a href="#">2009A&amp;A...497..511R</a> finds modest coronal X-ray emission, attributed to modest dynamo activity at the low or intermediate latitudes (the authors note that of the stars not in a tight binary system, <math>\alpha</math> Aql A is among the hottest known to have coronal X-ray emission)</p> <p>¶ <a href="#">2017A&amp;A...608A.113N</a> reports time-varying IR (K-band) excess, suggestive of tenuous circumstellar material (possibly debris disk: the “Be phenomenon,” present in many hot, young rapid rotators, is believed to involve a gas disk rather than a debris disk)</p> <p>¶ since proper motions of <math>\alpha</math> Aql A and <math>\alpha</math> Aql B are discrepant, AB is not a true binary (AB astrometry: 143″→196″, PA 335°→286°, 1781→2015); under IAU rules, the name “Altair” designates just <math>\alpha</math> Aql A</p> <p>Cepheid variable, 3.49–4.30 in V, 7.2 d (AAVSO(VSX) assessment in 2022: status flag = confirmed variable; variability classification symbol = “DCEP”); AAVSO(VSX) as viewed 2022 Jul. 15 gives 7.17679 d (same value as given 2021 Jan. 28 and 2022 March 03, and in 2019 January); BSC5 asserts 7.176641 d with period changes; <a href="#">2002ApJ...140..465B</a> (in centre panel of the author’s Fig 1) gives (1990s?) photometry (to tighter than <math>\pm 10</math> millimag), colour, and radial-velocity curves</p> <p>¶ <math>\eta</math> Aql AB has been split with HST WFC3 (cf <a href="#">2020ApJ...905...81E</a>, A-to-B distance is <math>\sim 200</math> au, AB period <math>\sim 900</math> y); the AB astrometry so far comprises just 2 observations (0.7″→0.7″, 2011→2012); A is for its part an SB not as yet resolved, for which <a href="#">2020ApJ...905...81E</a> gives period 4 y, suggesting that the orbit is eccentric and is seen nearly face-on (i.e. is seen in the orientation least favourable for the radial-velocity investigation required in orbit spectroscopy)</p> <p>¶ in the case of novice Northern Hemisphere observers troubled by frequent cloud, its rather long period makes <math>\eta</math> Aql A a better high-amplitude Cepheid demonstration than the more celebrated <math>\delta</math> Cep A</p> <p><b>[THIS STAR ONLY IN ONLINE VERSION OF TABLE]</b></p>
$\gamma$	Sge	19 59.8+19 33	3.48 <sup>†</sup> 1.57	M0 III <sup>†</sup>	13	–1.0 260	0.070 71	–33 V?	<p>radius 0.26 au (from interferometry; the disk subtends an angle of 6.18 mas)</p> <p>¶ Astron. Alm. (epoch 2021.5) assigns MK type M0<sup>+</sup> III</p> <p>¶ AAVSO(VSX), as of 2022 Jul. 31, notes the existence of NSV entry and the nonexistence of AAVSO observations, gives a V-passband range of 0.004 with period 6.37836 d (AAVSO(VSX) assessment: status flag = confirmed variable; variability classification symbol = “VAR”); further photometric study advisable?)</p> <p>¶ already has a dead carbon core, is not yet a Mira</p> <p><b>[THIS STAR ONLY IN ONLINE VERSION OF TABLE]</b></p>
$\theta$	Aql Aa	20 12.5 –0 45	3.24 <sup>†</sup> –0.07 <sup>†</sup>	B9.5 III <sup>†</sup>	11	–1.5 290	0.036 81–27	SB2 <sup>†</sup>	<p>a good marker of celestial equator</p> <p>¶ there may be some unclarity in the literature regarding “Aa,” “Ab” nomenclature (cf WDS Note): it is at any rate clear that the SB2 that is <math>\theta</math> Aql A has now been interferometrically resolved, and that two quite similar orbital solutions have been published (period 17.124 d, <math>e \approx 0.6</math>); recent discussions include <a href="#">1995AJ...110..376H</a>, <a href="#">2000A&amp;AS..145..215P</a></p> <p>¶ <math>m_v</math>, B–V values are for <math>\theta</math> Aql Aa,Ab combined light</p> <p>¶ <math>\theta</math> Aql A pairing is metal-rich</p> <p>¶ AAVSO(VSX), while noting the existence of an NSV entry, has the following assessment in 2022: status flag = confirmed non-variable; variability classification symbol = “CST”</p> <p>¶ Astron. Alm. (epoch 2021.5) assigns MK type B9.5 III<sup>+</sup></p>
$\beta$	Cap Aa <sup>†</sup>	20 22.3–14 42	3.08 <sup>†</sup> 0.79 <sup>†</sup>	K0: II <sup>†</sup>	10	–2.0 300	0.046 81	–19 SB	<p>hierarchical quintuplet (or greater) Dabih</p> <p><a href="https://en.wikipedia.org/wiki/Beta_Capricorni">en.wikipedia.org/wiki/Beta_Capricorni</a> has a diagram summarizing the known gravitationally bound hierarchy:</p>



												<p>finds no record of AAVSO observations, offers possible V-passband range 1.93–1.96 (AAVSO(VSX) assessment: status flag = suspected variable; no variability classification symbol); further photometric study advisable? ¶ <a href="#">1988A&amp;A...20L.273V</a> discusses galactic-astronomy implications of puzzling deuterium paucity in <math>\alpha</math> Pav A</p> <p>¶ <math>E(B-V)=+0.02</math></p> <p>¶ the name, although anomalously English, is nevertheless IAU-official: its origins lie in 1930s RAF Air Almanac project, which directed HM Nautical Almanac Office that no air-navigation star was to be left nameless</p>
$\alpha$	Ind A	20 39.2–47 12	3.11	1.00	K0 III CN–1 <sup>†</sup>	33	0.7	98	0.083	37	–1	<p>Fe overabundant (<math>\alpha</math> Ind AB born in metal-rich ISM cloud?)</p> <p>¶ AAVSO(VSX) situation in 2022: no status flag (so not confirmed variable, not suspected variable, not confirmed non-variable), and a fortiori no variability classification symbol</p>
$\alpha$	Cyg A	20 42.2+45 22	1.25 <sup>†</sup>	0.09 <sup>†</sup>	A2 Ia	2 <sup>†</sup>	–6.9~1400 <sup>†</sup>	0.003	47	–5	<p><b>Deneb</b></p> <p>blue supergiant, of radius ~0.5 au or ~1 au for context pertaining to this particular BSG in the general population of hypergiants and supergiants, cf <a href="https://en.wikipedia.org/wiki/List_of_largest_stars">en.wikipedia.org/wiki/List_of_largest_stars</a> (which adopts “~1 au”); for current state of theoretical investigations into BSG populations (crossing Hertzsprung-Russell diagram for the first time, redward? or, rather, after episode of mass loss, crossing for the second time, blueward?) cf, e.g. <a href="#">2014MNRAS.439L...6G</a></p> <p>¶ slightly variable, and the prototype of the <math>\alpha</math> Cyg variables: AAVSO(VSX) gives range 1.21–1.29 in V (AAVSO(VSX) assessment in 2022: status flag = confirmed variable; variability classification symbol = “<b>ACYG</b>”); seemingly irregular (in the <math>\alpha</math> Cyg variables, many short-period oscillations are superimposed); <a href="#">2011AJ...14L...17R</a> discusses <math>\alpha</math> Cyg A, reporting a 1977-through-2001 campaign in both photometric and spectroscopic variability</p> <p>¶ <math>\alpha</math> Cyg A core hydrogen-fusion career started in MK spectral type B, or possibly even in the rare MK spectral type O</p> <p>¶ present mass-loss rate is ~8e-7 <math>M_{\odot}/y</math></p> <p>¶ slow rotator (period possibly as long as 0.5 y, consistently with its large radius and its ongoing mass loss)</p> <p>¶ public-outreach astro audiences enjoy comparing and contrasting distance, and therefore intrinsic luminosity, of <math>\alpha</math> Cyg A with distance, and therefore intrinsic luminosity, of the other two Summer Triangle stars (nearby <math>\alpha</math> Lyr A, nearby <math>\alpha</math> Aql A; all three are similar not only in their apparent magnitudes, but also in falling within MK type A, and consequently in lacking tint, even through binoculars); it is perhaps worth stressing in such lectures that the <math>\alpha</math> Cyg A distance, although large (1500 ly? more?), is nevertheless not yet well known; Kaler in <a href="https://stars.astro.illinois.edu/sow/deneb.html">stars.astro.illinois.edu/sow/deneb.html</a>, accepting ~1500 ly, writes that if placed at distance of <math>\alpha</math> Lyr A, <math>\alpha</math> Cyg A “would /.../ be as bright as a well-developed crescent Moon, cast shadows on the ground, and easily be visible in broad daylight”</p>	
$\eta$	Cep A	20 45.8+61 56	3.42 <sup>†</sup>	0.92 <sup>†</sup>	K0 IV <sup>†</sup>	70.1	2.6	46.5	0.823 <sup>†</sup>	6	–87 <sup>†</sup>	<p>high velocity relative to Sun indicates interloper status in galactic thin disk (and observed underabundance of Fe is consistent with interloper status)</p> <p>¶ <math>m_v</math>, B–V values are for <math>\eta</math> Cep AB combined light (where, however, the contribution of faint <math>\eta</math> Cep B, at mag. ~11, is minuscule)</p> <p>¶ AAVSO(VSX) situation in 2022: no status flag (so not confirmed variable, not suspected variable, not confirmed non-variable), and a fortiori</p>

$\beta$	Pav	20 47.0–66 07	3.42	0.16	A6 IV <sup>†</sup>	–24.1	0.3	135	0.044	283	+10	no variability classification symbol
$\varepsilon$	Cyg Aa <sup>†</sup>	20 47.2+34 04	2.47 <sup>†</sup>	1.03 <sup>†</sup>	K0 III	44.9	0.7	73	0.486 <sup>†</sup>	47–11 <sup>†</sup>	SB <sup>†</sup>	still a fast rotator ( $\leq 2.3$ d), although core hydrogen fusion is ended or is close to ending $\P$ AAVSO(VSX) situation in 2022: no status flag (so not confirmed variable, not suspected variable, not confirmed non-variable), and a fortiori no variability classification symbol $\P$ Astron. Alm. (epoch 2021.5) assigns MK type A6 IV–Aljanah the SB that is $\varepsilon$ Cyg Aa,Ab (period $\geq 15$ y, only one set of lines visible) has been interferometrically measured-and-resolved just twice, in the period 1983→1991 $\P$ $m_v$ , B–V values are for $\varepsilon$ Cyg Aa,Ab combined light $\P$ AAVSO(VSX) situation in 2022: no status flag (so not confirmed variable, not suspected variable, not confirmed non-variable), and a fortiori no variability classification symbol $\P$ velocity of Aa,Ab (and of faint, outlying, gravitationally bound red dwarf C, mag. 13.4) relative to Sun is high
$\zeta$	Cyg Aa <sup>†</sup>	21 13.9+30 19	3.21 <sup>†</sup>	0.99 <sup>†</sup>	G8 IIIa Ba <sup>†</sup> 0.5	23	0.0	140	0.069	175	+17 SB	in evolutionary terms, possibly a Red Clump resident (stable helium fusion in core); but it might also be the case that core-helium fusion has yet to begin $\P$ SB is resolved, as $\zeta$ Cyg Aa, $\zeta$ Cyg Ab: <a href="#">1992Obs...112..168G</a> discusses spectroscopy, reviewing the history at a level of detail so instructive as to make this a case study for spectroscopy technique generally, even outside the particular domain of the $\zeta$ Cyg system; an orbital solution for Aa,Ab has been published, asserting period 6489 d ( $\sim 18$ y), with $e=0.22$ ; on this solution inter-component distance is 8 au min, 13 au max, 11 au average, and angular length of semimajor axis of the rectified orbit is $\sim 190$ mas; Ab is a WD, of mag. 13.2; the Aa,Ab binary was first split with far-UV imaging from <i>IUE</i> (in general, UV is a desirable regime for observing binaries with an elusive WD secondary, since WDs, although faint in the V band, are UV-bright); additionally, <a href="#">2001MNRAS.322..891B</a> announces direct imaging with HST WFPC2 (elongated smear, WD partly resolved, possibly 36 mas; but this observation was made under unfavourable conditions, near periastron); as viewed 2022 Jan. 14, WDS documents just one successful astrometric data point for Aa,Ab, from the year 2000; the $\zeta$ Cyg Aa,Ab SB binary, long observed spectroscopically, and now open to space-based UV astrometry, is of interest for WD studies, since determination of orbit yields determination of WD mass (admittedly, a very small number of WD orbits, notably including the respective WD companions of two stars in this Handbook table, Sirius (in $\alpha$ CMa) and Procyon (in $\alpha$ CMi), have been determined even from terrestrial astrometry; with the advent of space-based UV astrometry, the overall WD mass-determination situation, historically something of a bottleneck, may now be expected to improve) $\P$ $m_v$ , B–V values are for $\zeta$ Cyg Aa,Ab combined light $\P$ AAVSO(VSX) situation in 2022: no status flag (so not confirmed variable, not suspected variable, not confirmed non-variable), and a fortiori no variability classification symbol $\P$ $\zeta$ Cyg Aa is chemically a mild barium star (Astron. Alm. (epoch 2021.5) assigns MK type “G8 <sup>+</sup> III–IIIa Ba 0.5 <sup>+</sup> ”); before becoming a WD, $\zeta$ Cyg Ab, as a mass-shedding AGB star, deposited barium onto $\zeta$ Cyg Aa
$\alpha$	Cep A	21 19.1+62 41	2.46 <sup>†</sup>	0.22	A7 Van <sup>†</sup>	66.5	1.6	49.1	0.158	72	–10 V	Alderamin fast rotator ( $< 12$ h); the rotational shape distortion, into an oblate spheroid, gives $\alpha$ Cep A a remarkable variation in envelope temperature, with the envelope convective (photosphere $\sim 6600$ K) at equator and

$\beta$	Cep <u>Aa</u> <sup>†</sup>	21 29.0+70 40	3.23 <sup>v†</sup> -0.22 <sup>†</sup>	B1 III <sup>†</sup>	5	-3.4 700	0.015	56	-8 SB	<p>radiative (photosphere ~8600K) at poles (the transition temperature is ~8300 K): a similar latitude-governed bifurcation in envelope characteristics is present (cf <a href="#">2011ApJ...732...68C</a> Fig. 9) in the rapid rotator <math>\alpha</math> Aql A (Altair)</p> <p>¶ Astron. Alm. (epoch 2021.5) assigns MK type A7 V+n</p> <p>¶ listed by AAVSO(VSX) as <math>\delta</math> Sct variable, nominal <math>m_v</math> 2.46 with range 0.002 in V (<i>TESS</i> mission), salient period 71.7 minutes (AAVSO(VSX) assessment in 2022: status flag = confirmed variable; variability classification symbol = “<b>DSCT</b>”)</p> <p>¶ several factors, including X-ray emission (consistent with corona, as might be expected for convection-harboring latitudes of the envelope) indicate magnetic activity variable: 3.16–3.27 in V, 0.19 d; B: 7.8; 13.5” (2016) Alfirk PA 255°→251°, 1779 →2016</p> <p>¶ variability range 3.16–3.27 in V; the prototype of the <math>\beta</math> Cep variables (although stars of this same type are sometimes called the “<math>\beta</math> CMA variables”), and (as is typical for the type) known to be multiperiodic: AAVSO supplies a 2010 Apr. 13 backgrounder at <a href="http://www.aavso.org/vsots_betacep">www.aavso.org/vsots_betacep</a>; AAVSO(VSX) as viewed 2021 Jan. 16 and 2022 Jul. 16 asserts salient period 4.57171 h; AAVSO archives a notice for an August 2009 <math>\beta</math> Cep campaign (coordinated photometry, spectroscopy, CHARA) at <a href="http://www.aavso.org/aavso-special-notice-162">www.aavso.org/aavso-special-notice-162</a> (AAVSO(VSX) assessment in 2022: status flag = confirmed variable; variability classification symbol = “<b>BCEP</b>”)</p> <p>¶ <math>\beta</math> Cep Aa is a magnetic star</p> <p>¶ system comprises at least (the much-studied variable) Aa and Ab (mag. 6.6, probably a Be-phenomenon star, and the origin of the Be-phenomenon behaviour observed in AaAb); Aa,Ab period is variously suggested as 50 y, 85 y; astrometry is now quite good, with 62 measurements over the period 1971→2007 (angular separation 0.3”→0.2”); if <math>\beta</math> Cep B is gravitationally bound to the Aa,Ab SB (no AB orbital solution has been published), then period is <math>\geq 40,000</math> y, with (Aa,Ab)-to-B distance 3,000 au</p> <p>¶ <math>m_v</math>, B–V values are for <math>\beta</math> Cep Aa,Ab,B combined light; GCPD gives directly measured <math>m_v</math>, B–V values for <math>\beta</math> Cep Aa,Ab combined light as 3.19, –0.23; GCPD gives directly measured <math>m_v</math>, B–V values for <math>\beta</math> Cep B as 7.83, 0.12</p> <p>¶ MK luminosity class III (“giant”) notwithstanding, <math>\beta</math> Cep Aa is still fusing hydrogen in its core</p>
$\beta$	Aqr A	21 32.8 –5 28	2.90 <sup>†</sup> 0.83 <sup>†</sup>	G0 Ib <sup>†</sup>	6	-3.2 500	0.020	114	+7 V?	<p>a rare instance of a yellow supergiant Sadalsuud</p> <p><math>\beta</math> Aqr A is possibly now evolving blueward in a second crossing of the HR diagram</p> <p>¶ spectroscopically a “hybrid” star, combining signature of hot corona with signature of cool massive wind; <a href="#">2005ApJ...627L..53A</a>, in a study jointly covering <math>\beta</math> Aqr A and the astrophysically similar hypergiant (likewise a hybrid) <math>\alpha</math> Aqr A, reports <i>Chandra</i> observation of coronal X-rays (first X-ray detection from a hybrid G supergiant; such supergiants are X-ray deficient, their coronae notwithstanding)</p> <p>¶ <math>\beta</math> Aqr lies in the IS on the HR diagram, and yet is not known to be a pulsator</p> <p>¶ their ~10° separation on the celestial sphere notwithstanding, <math>\beta</math> Aqr A and <math>\alpha</math> Aqr A have shared proper motion and similar parallaxes (and WDS <math>\beta</math> Aqr A is the same object as WDS <math>\alpha</math> Aqr C; this pairing of <math>\beta</math> Aqr A a.k.a. <math>\alpha</math> Aqr C with <math>\alpha</math> Aqr A is further discussed in <a href="http://bestdoubles.wordpress.com/2014/12/15/the-alpha-beta-gamma-of-aquarius-%CE%B1-%CE%B2-and-%CE%B3-aquarii/">bestdoubles.wordpress.com/2014/12/15/the-alpha-beta-gamma-of-aquarius-%CE%B1-%CE%B2-and-%CE%B3-aquarii/</a>)</p> <p>¶ AAVSO(VSX) situation in 2022: no status flag</p>

$\mu$	Cep A	21 44.2	58 53	4.0v <sup>†</sup>	2.35 <sup>†</sup>	M2- Ia	1? <sup>†</sup>	-8?	3000? <sup>†</sup>	0.007	155	+21	<p>(so not confirmed variable, not suspected variable, not confirmed non-variable), and a fortiori no variability classification symbol</p> <p>red supergiant, of dramatic colour in binoculars</p> <p>¶ semiregular var., 3.4–5.1 in V; <a href="#">1997JBA...107..135B</a>, using 1959-1993 data, finds periods of ~840 d and ~4400 d, and notes (“Table 3”) the broad consistency of this result with 7 other multi-decade photometric studies, all starting from various points in the 19th century; typical recent V-passband measurements from the AAVSO main database (not VSX) are, as expressed by us at the Handbook to just three significant figures, 4.00 (2022 Dec. 07), 3.84 (2022 Nov. 03), and 3.79 (2022 Oct. 16); the most recent brightness plateau, in the ~30-day period centred on 2022 Jul. 01, yielded V-band measurements of mag. ~3.4;</p> <p>for wider <math>\mu</math> Cep A photometric context, cf. <a href="#">en.wikipedia.org/wiki/List_of_semiregular_variable_stars</a></p> <p>¶ we take MK type from <a href="#">1989ApJS...71..245K</a>; however, <a href="#">2011ARep...55...31S</a> asserts emission (at temperature type M2, luminosity class Ia)</p> <p>¶ the determination of D is difficult, with the HIPPARCOS parallax very uncertain, and with <math>\mu</math> Cep A parallaxes in general troubled by the non-negligible angular width (~20 mas) of this extraordinarily distended star; following Table 2 in <a href="#">2020MNRAS.493..468D</a>, we compute D as 940 pc * (3.26 ly/pc) = 3064 ly, stating this to one significant figure (the cited “Table 2” gives a large uncertainty in the direction of increase, almost 15%, while giving a smaller uncertainty in the direction of decrease) as 3000 ly; the “Table 2” determination of D is based on reliable <i>Gaia</i> DR2 (2018) parallaxes for hot (and to the <i>Gaia</i> telescope adequately punctiform) celestial-sphere neighbours of <math>\mu</math> Cep A, assumed by the “Table 2” compilers to belong to the stellar association into which <math>\mu</math> Cep A itself was born; <a href="#">en.wikipedia.org/wiki/Mu_Cephei</a> refers to this “Table 2” determination, but additionally to two other very discrepant determinations; upon conversion to our own favoured units of ly, followed by reduction to a single significant figure, these Wikipedia-cited determinations become 1000 ly and 6000 ly, thereby yielding a discouragingly wide distance bracket</p> <p>¶ we compute pi directly from the “Table 2” D determination, as 1 mas</p> <p>¶ although <math>\mu</math> Cep A has no IAU-official name, it is often informally referred to as “Herschel’s garnet star”: in 1783, W.Herschel noted to its “very fine deep garnet colour,” and indeed many garnets are deep red</p> <p>¶ AAVSO(VSX) assessment ~2022: status flag = confirmed variable; classification symbol = <b>SRC</b></p> <p><b>[THIS STAR ONLY IN ONLINE VERSION OF TABLE]</b></p>
$\varepsilon$	Peg A	21 45.3	+9 59	2.39 <sup>†</sup>	1.52 <sup>†</sup>	K2 Ib <sup>†</sup>	5	-4.2	700	0.027	89	+5 V	<p>slight irregular var.: 2.37–2.45 in V (flare in 1972) <b>Enif</b> (AAVSO(VSX) assessment in 2022: status flag = confirmed variable; variability classification symbol = “LC”); <a href="#">1972IAUC.2392....1W</a> reports extreme flare-like brightening, ~10 minutes, to V mag. 0.7</p> <p>¶ orange-class supergiant</p> <p>¶ Astron. Alm. (epoch 2021.5) assigns MK type K2 Ib–II</p> <p>¶ <a href="#">1987MNRAS.226..563S</a> discusses abundances, finding that, earlier literature notwithstanding, <math>\varepsilon</math> Peg A is unremarkable in its barium (and unremarkable in its strontium), and therefore</p>

$\delta$	Cap A	21 48.3–16 01	2.86 <sup>v†</sup> 0.30	A3mF2 IV:†	84	2.5 38.7	0.396 139	–6 SB†	discounting an earlier suggestion that $\epsilon$ Peg A outer layers have hosted nucleosynthesis in slow-neutron capture $\P$ $m_v$ , B–V values are for $\epsilon$ Peg A alone (i.e. are not combined-light values) $\P$ BSC5 suggests “cooler shell surrounding” $\P$ WDS documents as celestial-sphere neighbours the faint $\epsilon$ Peg B (mag. 12.7, known to be not gravitationally bound to $\epsilon$ Peg A (angular separation 83” in 2013), and additionally the less faint $\epsilon$ Peg C (mag. 8.6; AC astrometry is 138”→144”, PA 323°→318°, 1825→2018) ecl. binary: V 2.81–3.05, 1.0 d, 3.2 + 5.2 Deneb Algedi $\P$ since SB has not been measured as a visual binary (not even interferometrically; the binarity has, admittedly, been demonstrated in at least one occultation: lunar occultations are possible, planetary occultations possible-yet-rare), WDS is not as yet able to write “ $\delta$ Cap Aa,” “ $\delta$ Cap Ab”; the $\delta$ Cap A pair is classified at AAVSO(VSX) as an Algol-type eclipsing binary, 1.0227688 d in AAVSO(VSX) as viewed 2021 Jan. 16, but 1.0227672 d as viewed 2022 Feb. 23 and 2022 Jul. 16; AAVSO(VSX) as viewed 2022 Jul. 16 asserts also $\delta$ Sct variability (denied, however, by <a href="#">1994MNRAS.266L..13L</a> ) in the $\delta$ Cap assemblage; further, AAVSO(VSX) has O-C, i.e. period-monitoring, plotting from 2016; secondary in the SB is ~3 mag. fainter than primary and is judged in <a href="#">1992MNRAS.259.251W</a> to be mildly active, possibly tidally locked, with large spot; A is known to be SB since 1906 (Slipher), and yet is known to be eclipsing only as of <a href="#">1956PASP...68..541E</a> ; <a href="#">1994MNRAS.266L..13L</a> remarks that “given the brightness of the system, $\delta$ Cap is poorly observed,” with period awkward for any one solitary observatory (an implication of this remark is that coordinated intercontinental photometry would now be helpful); AAVSO(VSX) as on 2022 Jul. 16 indicates that the AAVSO database has just 23 observations (further photometric study desirable?); AAVSO(VSX) assessment in 2022 is as follows: status flag = confirmed variable; variability classification symbol = “EA+GDOR+DSCT”) $\P$ Astron. Alm. (epoch 2021.5) assigns MK temperature type F2m and does not assign an MK luminosity class
$\gamma$	Gru	21 55.3–37 15	3.00 –0.12	B8 IV–Vs	15	–1.1 210	0.099 98	–2 V? Aldhanab	$\P$ AAVSO(VSX) situation in 2022: no status flag (so not confirmed variable, not suspected variable, not confirmed non-variable), and a fortiori no variability classification symbol
$\alpha$	Aqr A	22 07.0 –0 12	2.94 <sup>†</sup> 0.97 <sup>†</sup>	G2 Ib <sup>†</sup>	6	–3.1 –520	0.021 117	+8 V? Sadalmelik	a good marker of celestial equator $\P$ a rare instance of a yellow supergiant; possibly now evolving blueward in a second crossing of the HR plane; resides in the IS (under at least one definition of IS) and yet is nonpulsating (cf further <a href="#">2017AstL...43..265U</a> ) $\P$ spectroscopically a “hybrid star,” combining signature of hot corona with signature of cool, massive wind; <a href="#">2005ApJ...627L..53A</a> , in a study jointly covering $\alpha$ Aqr A and the astrophysically similar supergiant (likewise a hybrid star) $\beta$ Aqr A reports <i>Chandra</i> observation of coronal X-rays (first X-ray detection from a hybrid G supergiant; such supergiants are X-ray deficient, their coronae notwithstanding) $\P$ AAVSO(VSX) situation in 2022: no status flag (so not confirmed variable, not suspected variable, not confirmed non-variable), and a fortiori no variability-classification symbol $\P$ despite their ~10° separation on the celestial sphere, $\alpha$ Aqr A and $\beta$ Aqr A have shared proper motion and similar parallaxes (and WDS $\beta$ Aqr A is the same object as WDS $\alpha$ Aqr C; this pairing of $\beta$ Aqr A a.k.a.

$\alpha$	Gru A	22 09.7–46 51	1.74 <sup>†</sup>	–0.14	B7 Vn	32	–0.7	101	0.194	139	+12	<p><math>\alpha</math> Aqr C with <math>\alpha</math> Aqr A is further discussed in <a href="https://bestdoubles.wordpress.com/2014/12/15/the-alpha-beta-gamma-of-aquarius-%CE%B1-%CE%B2-and-%CE%B3-aquarii/">bestdoubles.wordpress.com/2014/12/15/the-alpha-beta-gamma-of-aquarius-%CE%B1-%CE%B2-and-%CE%B3-aquarii/</a></p> <p>possible slight variability (1.70–1.76 in V?) AAVSO(VSX), as of 2022 Jul. 31, notes the existence of NSV entry, finds no record of AAVSO observations (AAVSO(VSX) assessment in 2022: status flag = suspected variable; no variability classification symbol); further photometric study advisable? ¶ rapid rotator (&lt; 1d) ¶ E(B–V)=–0.02</p>	Alnair
$\theta$	Peg	22 11.4 +6 19	3.52 <sup>†</sup>	0.09	A2mA1 IV–V <sup>†</sup>	35	1.3	90	0.284	84	–6 SB2	<p>possible slight variability (3.48–3.56 in V?) AAVSO(VSX), as of 2022 August 01, notes existence of NSV entry, finds no record of AAVSO observations (AAVSO(VSX) assessment in 2022: status flag = suspected variable; no variability classification symbol) (further photometric study advisable?); assertion of <math>\delta</math> Sct variability, from somewhere in the earlier literature, seems to be now discounted ¶ rapid rotator (&lt; 20 h); consistently with rapid rotation, and therefore with a stirred atmosphere, elemental abundances are unremarkable [THIS STAR ONLY IN ONLINE VERSION OF TABLE]</p>	Biham
$\zeta$	Cep	22 11.7+58 19	3.35 <sup>†</sup>	1.57 <sup>†</sup>	K1.5 Ib <sup>†</sup>	3.9	–3.7	800	0.014	69	–18 SB	<p>slight irregular variability (3.31–3.40 in V) (AAVSO(VSX) assessment in 2022: status flag = confirmed variable; variability classification symbol = “LC”) ¶ orange supergiant either approaching core-helium fusion or already in core-helium fusion ¶ an eclipsing companion has been suggested, with suggestion later questioned ¶ metals somewhat overabundant</p>	
$\alpha$	Tuc	22 20.1–60 08	2.85	1.39	K3 III <sup>†</sup>	16	–1.1	200	0.081	241	+42 SB <sup>†</sup>	<p>SB 11.5 y, separation possibly 11.5 au ¶ primary in the SB is a giant, with carbon underabundant, nitrogen overabundant ¶ <a href="https://stars.astro.illinois.edu/sow/alphatuc.html">stars.astro.illinois.edu/sow/alphatuc.html</a> discusses uncertainties in the evolutionary stage of this giant, offering three scenarios ¶ AAVSO situation in 2022: no status flag (so not confirmed variable, not suspected variable, not confirmed non-variable), and a fortiori no variability classification symbol</p>	
$\delta$	Cep A <sup>†</sup>	22 30.0+58 32	3.60v <sup>†</sup>	0.48	F5–G2 Ib	4 <sup>†</sup>	–3.0	900 <sup>†</sup>	0.016	77	–15 SB <sup>†</sup>	<p>the prototype Cepheid variable: 3.49–4.36 in V, 5.4 d second-nearest Cepheid (<math>\alpha</math> UMi is still nearer) ¶ AAVSO offers a tutorial at <a href="https://www.eso.org/public/outreach/eduoff/aol/market/collaboration/varstar/pg2.html">www.eso.org/public/outreach/eduoff/aol/market/collaboration/varstar/pg2.html</a> and an initial backgrounder at <a href="https://www.aavso.org/vsots_delcep">www.aavso.org/vsots_delcep</a>; the first three sections of a paper directed inter alia to AAVSO observers, 2016JAVSO..44..179N, constitute a deeper backgrounder on the Cepheids ¶ AAVSO(VSX) has, as viewed 2021 Jan. 28 and again 2022 Feb. 24, 2022 Jul. 16, period 5.366266 d (is this value possibly now stale?); although Cepheids experience both period jitter and (monotonic) period slide, with a slide of even 200 s/y possible, 2014ApJ...794...80E finds <math>\delta</math> Cep period sliding slowly, at just –0.1 s/y (period decrease-increase is a signature of evolution, specifically of density increase-decrease, as a Cepheid passes across the HR diagram [<math>\delta</math> Cep is now making its second such passage, moving blueward]) (AAVSO(VSX) assessment in 2022: status flag = confirmed variable; variability classification symbol = “DCEP”) ¶ 2015ApJ...804..144A announces that <math>\delta</math> Cep A is SB, with period 2201 d ¶ accurate distances to Cepheids are foundational</p>	



												in cosmology, which needs independently known (galactic) Cepheid distances before embarking on its external-galaxy distance deductions through applications of the Cepheid Period-Luminosity (PL) Law; it is reassuring that the 2007 <i>HIPPARCOS</i> distance and the distance implied by the usual PL calculation agree to within uncertainties; although we have here stated the 2007 <i>HIPPARCOS</i> parallax, on which distance of $\delta$ Cep depends, as 4 mas, our cited 2007 <i>HIPPARCOS</i> determination is more formally, with decimal fractions and the uncertainty made explicit, $3.77 \pm 0.16$ mas; <a href="#">2015ApJ...804..144A</a> proposes instead $4.09 \pm 0.16$ mas, with the remark that impending <i>Gaia</i> may be expected, in part in the light of these authors' SB announcement, to secure an authoritative parallax; an already reassuring state of affairs may thus be expected to improve further
$\zeta$	Peg A	22 42.6+10 57	3.41 <sup>†</sup> -0.09	B8.5 III <sup>†</sup>	16	-0.6	210	0.078	98	+7 V?	¶ mass loss $\sim 1e-6 M_{\odot}/y$ ; bow shock in ISM has now been detected ¶ $\delta$ Cep C, at mag. 6.1, is in slow and wide orbit with $\delta$ Cep A (no orbital solution published; period 345,000 y; AC astrometry is $42'' \rightarrow 41''$ , PA $195^{\circ} \rightarrow 191^{\circ}$ , 1800 $\rightarrow$ 2018) [THIS STAR ONLY IN ONLINE VERSION OF TABLE] slight variability (3.40–3.41 in V) Homam AAVSO (VSX) as viewed 2022 Jul. 16 gives period 22.952 h, assigns the variability to the “slowly pulsating B stars” type (AAVSO(VSX) assessment: status flag = confirmed variable; variability classification symbol = “SPB”); additionally, <a href="#">2007PASP...119.483G</a> discusses satellite detection of amplitude $\sim 0.5$ millimag ¶ our (Garrison) MK spectral type notwithstanding, B8 V has been suggested	
$\beta$	Gru	22 44.1–46 46	2.11v <sup>†</sup> 1.62 <sup>†</sup>	M5 III <sup>†</sup>	18	-1.6	180	0.135	92	+2	¶ fast rotator (< 1.4 d) semiregular variable, 1.90–2.3 in V, 37 d Tiaki classified at AAVSO(VSX) as semiregular late-type giant, perhaps on the basis of <a href="#">2006JAVSO...34..156O</a> (this paper might serve as a case study for effective amateur-budget intercontinental photometry collaboration) (AAVSO(VSX) assessment in 2022: status flag = confirmed variable; variability classification symbol = “SRB”) ¶ among the rather uncommon cool red giants, with radius slightly > 0.8 au ¶ Astron. Alm. (epoch 2021.5) assigns MK type M4.5 III compos. spectrum, SB period 813 d Matar the $\eta$ Peg Aa,Ab SB is resolved in speckle interferometry, with orbital solution published (66 measurements: $0.1'' \rightarrow 0.1''$ , 1975 $\rightarrow$ 2005); BC is itself a tight pairing of equally bright stars, probably a true binary ( $0.3'' \rightarrow 0.2''$ , 1889 $\rightarrow$ 2011), probably in orbit with the Aa,Ab binary, making this a hierarchically organized quadruple system; A,BC astrometry is $90'' \rightarrow 92''$ , PA $339^{\circ} \rightarrow 338^{\circ}$ , 1824 $\rightarrow$ 2012 ¶ we take the MK type from WDS Note ¶ $m_v$ , B–V values are for $\eta$ Peg Aa,Ab combined light; it is known that individually, in the R passband as distinct from the V passband, $\eta$ Peg Aa and $\eta$ Peg Ab are respectively mag. 4.1, mag. 6.9 ¶ AAVSO(VSX), as of 2022 Jul. 16, indicates possible slight variability for the $\eta$ Peg assemblage (range 2.92–2.96 in V), finds no record of AAVSO observations, notes existence of NSV entry (AAVSO(VSX) assessment: status flag = suspected variable; no variability classification symbol); further photometric study advisable? ¶ $\eta$ Peg Aa is a slow rotator (818 d?) possible slight variability (3.47–3.53 in V?) AAVSO(VSX), as of 2022 Jul. 16,	
$\eta$	Peg Aa <sup>†</sup>	22 44.1+30 21	2.95 <sup>†</sup> 0.85 <sup>†</sup>	G8 II–III + F0 IV	15	-1.2	210	0.029	153	+4 SB		
$\varepsilon$	Gru	22 50.0–51 12	3.48 <sup>†</sup> 0.08	A2 Va	25	0.5	130	0.126	121	0 V		

												finds no record of AAVSO observations, notes existence of NSV entry (AAVSO(VSX) assessment: status flag = suspected variable; no variability classification symbol) ¶ rapid rotator (< 0.65 d) [THIS STAR ONLY IN ONLINE VERSION OF TABLE]
ι	Cep	22 50.5 +66 19	3.52	1.06	K0 III <sup>†</sup>	28.3	0.8	115	0.141	208	-12	Astron. Alm. (epoch 2021.5) assigns MK type K0 III ¶ AAVSO(VSX) situation in 2022: no status flag (so not confirmed variable, not suspected variable, not confirmed non-variable), and a fortiori no variability classification symbol [THIS STAR ONLY IN ONLINE VERSION OF TABLE]
μ	Peg	22 51.1 +24 44	3.49	0.94	G8 III <sup>†</sup>	31	0.9	106	0.151	106	+1	Astron. Alm. (epoch 2021.5) assigns MK type G8 III ¶ AAVSO(VSX) situation in 2022: no status flag (so not confirmed variable, not suspected variable, not confirmed non-variable), and a fortiori no variability classification symbol [THIS STAR ONLY IN ONLINE VERSION OF TABLE] Sadabari
δ	Aqr	22 55.9 -15 42	3.27 <sup>†</sup>	0.06	A3 IV-V	20	-0.2	160	0.051	237	+18 V	possible slight variability (3.25-3.29 in V?) AAVSO(VSX), as of 2022 Jul. 16, finds no record of AAVSO observations, notes existence of NSV entry (AAVSO(VSX) assessment: status flag = suspected variable; no variability classification symbol); further photometric study advisable? ¶ weak λ4481 ¶ rapid rotator (< 3.0 d) 2008 (HST) image was debris cloud, not exoplanet <b>Fomalhaut</b> HST putative 2008 “exoplanet” α PsA Ab was IAU-named Dagon, after a Semitic deity; at ~125 au, in the outermost of the debris rings; Dagon was in always-wide (albeit eccentric) orbit, making direct imaging, as opposed both to spectroscopy (for star Doppler wobble) and astrometry (for star transverse wobble) the tool of choice: 32 au min, 320 au max; period ~1700 y; in more recent years, it was suggested that Dagon could be a mere dust cloud, or an aggregation of rubble, or a single rocky body; an explanation was needed for the fact that Dagon proved so readily HST-visible (e.g. visibility enhanced by circumplanetary dust sphere, or by circumplanetary ring system?); Dagon mass was uncertain (< 2× Jupiter, perhaps even ~Earth); but with Dagon now no longer HST-visible, it would appear that the 2008 HST image was of an expanding debris cloud, now become too tenuous for detection ¶ the nested circumstellar dust rings extend as far as radius ~150 au (a distance recalling the Solar System Kuiper Belt); <a href="#">2017ApJ...842...8M</a> reports complete outer debris-ring mapping, via ALMA (223 GHz radio), finding ring mass of 0.015 Earths, eccentric, with α PsA Aa offset from the ring centroid ¶ α PsA Aa is a fast rotator (< 1d) ¶ in evolutionary terms, α PsA Aa is sufficiently young to be undergoing an analogue of the Solar System’s Late Heavy Bombardment (and consistently with this, <a href="#">2017ApJ...842...9M</a> writes that exocometary gas is detected, in ALMA 230 GHz radio) ¶ <a href="#">2017ApJ...842...8M</a> comments that “given its unique characteristics and architecture, the Fomalhaut system is a Rosetta stone for understanding the interaction between planetary systems and debris disks” ¶ α PsA Aa has low metallicity ¶ <a href="#">2013AJ...146..154M</a> , working both from proper motion (across the celestial sphere) and from velocities along the line of sight, concludes that α PsA Aa, α PsA B, and α PsA C belong to the same system: B (a flare star) is V mag. 7, at angular separation almost 2° from Aa (period ≥ 7.6 My), while C is V mag. 13, at enormous angular separation 5.7° from Aa Skat
α	PsA Aa <sup>†</sup>	22 58.9 -29 30 <sup>†</sup>	1.16	0.09	A3 Va <sup>†</sup>	130	1.7	25.1	0.368	1 17	+7	

											(and yet at a sufficiently low distance from Aa-with-B to have the Aa-with-B gravitational field dominate the general external gravitational field at its location; period is $\geq 35$ My) ¶ AAVSO(VSX), as of 2022 Jul. 16, finds no record of AAVSO observations, notes existence of NSV entry, writes “1.15-?” for V-passband range (AAVSO(VSX) assessment: status flag = suspected variable; no variability classification symbol = “LC”); further photometric study advisable? ¶ $\beta$ Peg, $\alpha$ Peg serve as pointers: since $\alpha$ PsA Aa lies a couple of arcminutes N of DEC $\approx$ -30°, $\alpha$ PsA Aa rises (if briefly) above the horizon even for such Canadian subarctic communities as Churchill, and for such Scandinavian communities as Stavanger semiregular variable, mag. 2.31–2.74 in V, 43.3 d Scheat (AAVSO(VSX) assessment in 2022: status flag = confirmed variable; variability classification symbol = “SR”) ¶ Astron. Alm. (epoch 2021.5) assigns MK type M2.5 II–III ¶ an intermediary between straightforward red giant and red bright giant (radius $\sim 0.5$ au); mass-loss rate is notably low for such a star ( $\leq 1e-8 M_{\odot}/y$ ; i.e. $\sim 100\times$ lower than mass loss rate of $\alpha$ Ori Aa (Betelgeuse); IRAS detected no IR excess)
$\beta$	Peg A	23 04.9+28 13	2.46v <sup>†</sup> 1.67	M2 II–III <sup>†</sup>	16.6	-1.5 ~196	0.232	54	+9 V		
$\alpha$	Peg	23 05.9+15 20	2.48 -0.04	A0 III–IV	24	-0.6 133	0.073	124	-4 SB	<b>Markab</b>	
										the existence of an NSV entry notwithstanding, AAVSO(VSX), as of 2022 Jul. 16, asserts constant light, and notes existence of 143 AAVSO observations (AAVSO(VSX) assessment in 2022: status flag = confirmed non-variable; variability classification symbol = “CST”) ¶ rapid rotator (1.5 d)	
$\gamma$	Cep $\underline{\Delta}^{\dagger}+1P^{\dagger}$	23 40.3+77 46	3.21 1.03	K1 III–IV <sup>†</sup>	71	2.5 46	0.135	339	-42 V?	<b>Errai</b>	
										possible slight variability (3.18–3.24 in V?) AAVSO(VSX), as viewed 2022 Jul. 16, finds no record of AAVSO observations, notes existence of NSV entry (AAVSO(VSX) assessment: status flag = suspected variable; no variability classification symbol); further photometric study advisable? ¶ the binary $\gamma$ Cep A, $\gamma$ Cep B has been split with adaptive optics at Subaru, with direct imaging reported in <a href="#">2007A&amp;A...462..777N</a> (AB astrometry: 3 measurements, 0.9 $\rightarrow$ 0.9", PA 257 $\rightarrow$ 256°, 2006 $\rightarrow$ 2006); the IAU-official name “Errai” applies to $\gamma$ Cep A rather than to the two-star system $\gamma$ Cep AB; Errai hosts an exoplanet, IAU-named Tadmor, which is not as yet astrometrically observed by any optical technique (so WDS is not as yet able to write “ $\gamma$ Cep Aa,” “ $\gamma$ Cep Ab”); Tadmor, circumstellar without being circumbinary, is among the few exoplanets discovered in a two-star system; Tadmor orbital period is 2.47 y, with average distance from Errai 2.05 au, and with mass between 3x Jupiter and 16x Jupiter; A-to-B distance is 12 au min, 25 au max, AB orbital period (orbital solution has been published) is 66 y or 67 y ¶ Errai rotation period is possibly 781 d (making this star a slow rotator) ¶ Astron. Alm. (epoch 2021.5) assigns MK type K1 III–IV CN 1	