

PREFACE: Scope and purpose of this essay

This online essay is an extended version of the essay in the printed-edition Handbook, containing all the material of its printed-edition accompaniment, but adding material of its own. The accompanying online table is likewise an extended version of the printed-edition table, (a) with extra stars (after providing for multiplicity, as we explain below, the brightest MK-classified 322, allowing for variability, where the printed edition has almost 30 fewer, allowing for variability: our cutoff is mag. ~ 3.55), and (b) with additional remarks for most of the duplicated stars. We use a dagger superscript (\dagger) to mark data cells for which the online table supplies some additional information, some context, or a caveat.

The online essay and table try to address the needs of three kinds of serious amateur: amateurs who are also astrophysics students (whether or not enrolled formally at some campus); amateurs who, like many in RASC, assist in public outreach, through some form of lecturing; and amateurs who are planning their own private citizen-science observing runs, in the spirit of such “pro-am” organizations as AAVSO. Additionally, we would hope that the online project will help serve a constituency of sky-lovers, whether professional or amateur, who work with the heavens in an unambitious and contemplative spirit, seeking to understand at the eyepiece, or even with the naked eye, the realities behind the little that their limited circumstances may allow them to see. (This is the same contemplative exercise as is proposed for the Cyg X-1 black hole, with its gas-dumping supergiant companion HD226868, in the Handbook printed-editions “Expired Stars” essay: with a small telescope, or even with binoculars, we first find HD226868, and then take a moment to ponder in awe the accompanying unobserved realities of gas-fed hot accretion disk, event horizon, and spacetime singularity.)

Our online project, started as a supplement to the 2017 Handbook, must be considered still in its rather early stages. We cannot claim to have fully satisfied the needs of our various constituencies. Above all, we cannot claim to have covered all the appropriate points from stellar-astronomy news in our “Remarks” column, important though news is to amateurs of all three types. We would hope in coming years to remedy our deficiencies in several ways, most notably by relying more in our writing on recent primary-literature journal articles, with appropriate explicit citations.

In our citations, we favour the now-preferred astrophysics “bibcode” formalism. The formalism is documented in <http://simbad.u-strasbg.fr/guide/refcode/refcode-paper.html>, and again in section 1.2.3 (headed “Bibliographic Identifiers”) in http://adsabs.harvard.edu/abs_doc/help_pages/data.html.

A bibcode can be transformed into the display of a more human-readable bibliography entry, often with clickable hyperlink to an underlying online full-text, all-illustrations PDF publication, in various ways. We illustrate some possibilities by taking an extreme case, namely our bibcode reference to the classic 1910 Joel Stibbins *Astrophysical Journal* paper that reports the electric-photometry discovery of a secondary minimum in the Algol light curve. Old though the paper is, it is nevertheless available online. The bibcode (as we state again in our “Remarks” for the Algol entry in our table) is [1910ApJ...32..185S](http://simbad.u-strasbg.fr/simbad/sim-ref?bibcode=1910ApJ...32..185S). A browser display with hyperlink to the desired full-text, all-illustrations PDF is available from the Centre de Données Stellaires (CDS) server (probably in Strasbourg) as <http://simbad.u-strasbg.fr/simbad/sim-ref?bibcode=1910ApJ...32..185S>. If something has gone wrong—and experience suggests that things can go wrong, even when a bibcode appears to casual inspection to be correctly typed, at any rate in some such autonomous-agent computing environment as Microsoft Office—then one can recover through CDS as <http://simbad.u-strasbg.fr/simbad/sim-fid> if the star of interest and year of publication are known. In this particular case, recovery involves giving <http://simbad.u-strasbg.fr/simbad/sim-fid> some convenient identifier, for instance the IAU-promulgated name “Algol” or the Bayer identifier “beta Per.” In the Algol-specific input form generated, one next asks, in the “References” section of the form, for all references from 1910 to 1910. The duly displayed bibcode, [1910ApJ...32..185S](http://simbad.u-strasbg.fr/simbad/sim-fid), for the sole 1910-through-1910 reference, is shown as a clickable hyperlink. Upon further clicking, the hyperlink eventually yields the PDF. A similar browser display is available from a (probably North American) ADS-NASA server as <https://ui.adsabs.harvard.edu/abs/1910ApJ...32..185S/abstract>. As a fourth possibility, the PDF is retrievable through a self-evident set of steps that starts by copying and pasting the bibcode into the “Bibliographic Code Query” box at the paper-workflow, as distinct from the more obviously accessible paperless-workflow, online form <https://ui.adsabs.harvard.edu/paper-form>. This fourth method has the advantage that multiple bibcodes can be entered within a single query. As a fifth possibility, which in our view cannot be guaranteed to work (but there seem to be intermittent problems with the fourth possibility as well; and in general, servers should not be presumed fully reliable, in any discipline) is simply to put [1910ApJ...32..185S](http://simbad.u-strasbg.fr/simbad/sim-fid) naively into a general Google search, and to explore the ensuing chain of hyperlinks: in the case of at least a heavily cited paper, one is likely soon enough to reach an abstract at ADS-NASA or some similar authority, with accompanying PDF.

The bibliographic support of <http://simbad.u-strasbg.fr/simbad/> and <https://ui.adsabs.harvard.edu/>, as the principal tools for our primary-literature searching, is herewith gratefully acknowledged, as are Wikipedia (in exact-science topics, generally careful and up to date); *Sky & Telescope*, the web materials of Prof. James Kaler, and at a more technical level, two key sources of data, the Washington Double Star Catalog (WDS) and AAVSO. Helpful at AAVSO are not only the general graphing facility and the general AAVSO record of observations, but also a more recent offering, the VSX online database.

SECTION 1: Selection bases for our 315 nominal “bright stars,” strictly 322 MK-classified bright stars

Of our selected 315 nominal stars, three call for extra comment pertinent to this mag. ~ 3.55 naked-eye selection criterion. (1) κ (kappa) CMa (at RA $\sim 6^{\text{h}}50$) brightened in the 1960s or 1970s, just managing to meet the cutoff, and has remained bright. This change

was unfortunately not noted in the RASC Handbook until 2019. Should κ CMa now once again fade, we propose to keep listing it for at least a few years, since it is a variable of the γ Cas type (and may therefore be liable to yet further episodes of brightening during the 21st century; in general, the γ Cas variables, whether temporarily bright or temporarily faint, are desirable targets for ongoing, regular, citizen-science spectroscopy, and even naked-eye, monitoring, being closely associated with the amateur-relevant “Be phenomenon,” which we discuss in the final subsection of this essay). (2) We discontinued listing L₂ Pup (at RA ~07h14) in the 2017, 2018, and 2019 Handbooks. Now, however, we revert to our pre-2017 policy, since L₂ Pup is a semi-regular pulsator, occasionally bright. (3) T CrB (at RA ~16h00) has shown nova behaviour, brightening from its current very faint state (mag. ~10) to mag. 2.0 in 1866 and to mag. 3.0 in 1946. We have for years listed this star in our table and propose to continue listing it (since the history suggests the possibility of a 21st-century outburst).

An omission from our selection of 315 nominal stars also calls for comment: while mindful of the fact that η Car brightened greatly, attaining even mag. 0 for a few years from 1837 onward, we omit it from our table since there is no firm prognosis of a 21st-century repetition of that outburst. On 2021 January 7, η Car was being variously reported visually as at mag. 4.3 and 4.5, in either case being decidedly fainter than our naked-eye cutoff. The star was not reported by AAVO visual observers as any brighter than mag. 3.9 in the second half of 2020.

We may now explain in what sense the set of 315 is nominal. In a strict accounting, the selection is a set of 315 objects that are in naked-eye terms “bright stars,” i.e. are bright unresolved point sources of starlight. Three kinds of situation need to be distinguished here, as we move from naked-eye impressions to underlying physical realities:

(1) In some cases (to cite an example at random, β Tau (Elnath)) what is to the naked eye a point source actually is, so far as is known, a solitary star.

(2) Very common is a situation in which a bright star is a component in a multi-star system, with the other member(s) making either a very small or a negligible contribution to the naked-eye retinal signal. An instance of the former type of pairing is γ And A, from which at a distance of just over 9" lie two fainter stars, γ And B and γ And C, themselves separated by a mere 0.2", and so faint that the BC pairing shines at around mag. 5. This has the consequence that BC makes just a modest contribution to the overall γ And ABC naked-eye neuron response. An instance of the latter type of pairing is α CMa A (Sirius), with α CMa B a white dwarf shining at mag. 8.5, in other words shining so feebly as to play essentially no role in the signal generated by the naked-eye retina. This binary constitutes a not trivial, and yet also at the present favourable time a not hopeless, project for the small telescope. (At <http://www.rasc.ca/sirius-observing-challenge>, RASC notes that with apastron due in 2025, “an extremely difficult feat has become merely a very demanding one.”) Since our table is officially a table of bright stars, we take care, at any rate in our various table revisions from early 2021 onward, to write in our first table column “ γ And A” (not “ γ And AB” or “ γ And”), and “ α CMa A” (not “ α CMa AB” or “ α CMa”). Helpfully, the naming rules promulgated since around 2016 at IAU, and reflected in our concluding “Remarks” column, stipulate, in parallel with our first-column decision, that a name such as “Sirius” applies to a star such as α CMa A, rather than to the binary system α CMa AB.

(3) In ten other cases, the naked-eye point, shining at mag. ~3.55 or brighter, is the combined light of two binary-system components, each individually bright enough to count as a “bright star”—perhaps with each component exceeding our mag. ~3.55 cutoff, but also perhaps with one or both components just a little fainter than our mag. ~3.55 cutoff, and yielding a “star” brighter than mag. ~3.55 upon combining the light.

These ten, so-to-speak awkward, cases (awkwardly forcing us to write the binary designations “AB” or “Aa,Ab” in the first column) are the following:

- β Phe AB (with each of A, B individually around mag. 4, yielding an aggregate naked-eye impression of mag. 3.2)
- γ Per Aa,Ab (with each of Aa, Ab a little brighter than mag. 4, yielding an aggregated naked-eye impression of mag. 2.91)
- α Aur Aa (Capella), Ab (with each of Aa, Ab very close to mag. 0)
- β Aur Aa (Menkalinam), Ab (with magnitudes nearly equal, yielding an aggregated naked-eye impression a little brighter than mag. 2)
- γ Vir A (Porrina), B (magnitudes nearly equal, and with each individually very close to our mag. ~3.55 cutoff, yielding an aggregated naked-eye impression a little brighter than mag. 3)
- β Cen Aa (Hadar), Ab (magnitudes nearly equal, with each individual star much brighter than our mag. ~3.55 cutoff)
- η Oph A (Sabik), B (with B at mag. 3.5)
- λ Sco Aa (Shaula), Ab (with even Ab well above our cutoff, at mag. ~2.8)
- ζ Sgr A (Ascella), B (with B at mag. 3.5)
- π Sgr A (Albaldah), B (a poorly documented pairing, with the faint outlier C also poorly documented: WDS implies that B is of nearly the same magnitude as A, with each of these two stars very close to our mag. ~3.55 cutoff)

It is tempting to consider the η Peg system to be an eleventh case, requiring entry as “ η Peg Aa (Matar), Ab”. But since Ab is decidedly fainter than our mag. ~3.55 cutoff, and Aa only slightly fainter than our mag. ~3.55 cutoff, we are obliged instead to enter this case simply as “ η Peg Aa (Matar)”, drawing attention in the table “Remarks” column to the fact that our stated magnitude of 2.93 is for the combined light.

Somewhat like the η Peg system is the o (omicron) Leo system, where o Leo Aa (Subra) is very close to mag. 3.5, and where o Leo Ab is, while fainter than mag. 3.5, nevertheless bright enough to make a non-trivial contribution to the overall visual impression.

Before 2021, our table unfortunately had the erroneous information that α Leo Aa,Ab is a binary system in which the components are of equal magnitude.

We thus have a table of nominally 315 stars, comprising a more refined, i.e. less nominal, analysis $315 + 10 = 325$ bright stars. With 3 exceptions, each of the 325 has a known (at worst, an uncertainty-flagged) MK temperature type and MK luminosity class (with the Sun, of course, better observed than any of the others). The final result is accordingly a set of $325 - 3 = 322$ bright stars of known MK classification.

SECTION 2: General characteristics of our 322 MK-classified bright stars

Our 322-element sample is found to lie in a region, around 3000 ly in radius, essentially confined to the sandwich-filler, or “thin disk,” part of the overall galactic disk. Of the few Sample-S interlopers born outside the sandwich filling, and now temporarily passing through it on orbits oblique to the thin disk, the best known is α Boo (Arcturus). It is convenient here to use the term “Population P” for the ensemble of non-brown-dwarf, non-white-dwarf stars in the much larger, 3000-ly radius, subdisk-of-the-thin-disk from which our (tiny) Sample S is drawn. This P-region is itself only a (tiny) fraction of the overall galactic thin-disk region of stars, $\sim 50,000$ ly in radius.

Sample S, being formally defined by an apparent-magnitude cutoff as opposed to a distance cutoff, is itself far from statistically representative of Population P. (a) In P, the O stars are vanishingly rare. A tabulation by Glenn Ledrew, in *JRASC* **95** (2001), pp. 32ff (bibcode [2001JRASC..95...32L](https://ui.adsabs.org/2001JRASC..95...32L)) suggests an O-star frequency within P of just 0.00003%. By contrast, O stars comprise a hefty $\sim 2\%$ of S. A similar overrepresentation occurs for the B, A, F, G, and K stars, with Ledrew’s tabulation suggesting that these MK temperature types might have a respective frequency within P of 0.1%, 0.6%, 3.2%, 8.0%, and 12.9%. By contrast, the first three of these five rare types comprise $\sim 30\%$, $\sim 20\%$, and $\sim 10\%$, respectively, of S, and the last $\sim 20\%$ of S. (b) In P, something on the order of 76% or 78%—different authorities are perhaps mildly discrepant—must be M stars. (Ledrew’s tabulation, in particular, suggests an M-star frequency of 78.2%.) Only a few of these (the Ledrew tabulation suggests 0.04%) have evolved to beyond the main-sequence stage of stable-core hydrogen fusion. By contrast, the M stars comprise just $\sim 5\%$ or $\sim 10\%$ of S. All of them have evolved beyond the main sequence, having started their lives as types hotter than M or K.

The statistically anomalous character of S is further illustrated by the fact that in S, in each of the Big Six MK temperature types hotter than M, the numerical majority comprises the stars that have ended stable-core hydrogen fusion (and so have, as a generally reliable rule—we return below to a necessary caveat regarding reliability—evolved out of MK luminosity class V into one of the brighter MK luminosity classes IV, III, II, or I). In Ledrew’s tabulation, the percentages of evolved stars in F, G, and K, as a percentage of the overall respective F, G, and K populations, are just 2.0%, 2.5%, and 3.8%. Consistently with this, the 1991 Gliese-Jahreiss catalogue of the nearest 1000 stars (containing, admittedly, not only the local OBAFGKM VI, V, IV, III, II, and I stars, but also at least many of the local white dwarfs) assigns less than 1% of its population to MK luminosity classes IV, III, II, or I.

Sample S—so rich in varieties of star statistically infrequent within Population P—harbours physical extremes. Although the extremes are for the most part not written into our table, they can be studied easily, from such sources as Prof. James Kaler’s <http://stars.astro.illinois.edu/sow/sowlist.html>.

At least 58 of our 322-star set each radiate, across the full spectrum from X-ray through UV and optical to IR and radio, at least as much power as is radiated by 10,000 Suns. Possibly the most dramatic is ζ Ori, with a bolometric luminosity of 375,000 Suns—making ζ Ori notable not within S alone, but even within the overall galaxy. Several others are not far behind, among them ζ Pup (360,000 Suns, suggests Kaler, as of July 2008 revising his earlier, circa-1999, suggestion of $\sim 750,000$ Suns). We believe that just two stars in Sample S, nearby τ Cet and nearby α Cen B, radiate more feebly than our Sun, each at about half of the Sun’s bolometric luminosity.

The principal determinant of stellar luminosity, for any given phase in stellar evolution, is mass, with even small variations in mass translating into large variations in energy output. The exceptional luminosities of ζ Ori and ζ Pup, in particular, are a consequence of their exceptionally high respective masses, $20 M_{\odot}$ and $40 M_{\odot}$. (Kaler now suggests $40 M_{\odot}$ for ζ Pup, while having previously suggested $60 M_{\odot}$. He additionally notes from the literature the lower suggested value of $22.5 M_{\odot}$.)

Theory does predict, although our small Sample S does not succeed in illustrating, the possibility of masses up to the Eddington stellar-mass limit, somewhere above $100 M_{\odot}$, and even of some “super-Eddington” stars. (Eddington’s limit is by definition attained when luminosity rises so high as to make the outward radiation push, tending to tear a star apart, exceed the inward gravitational pull.)

Rotation periods in Sample S vary from far in excess of our Sun’s to far short of our Sun’s (which we may here take as a nominal 27 d; refined treatments of solar rotation provide for rotation-period variations both with solar latitude and with solar depth). Spectroscopy yields for γ Cep a period of 781 d, i.e. of 2.14 y. Kaler suggests that the respective rotation periods of α Hya and ϵ Crv could be as long as 2.4 y and 3.9 y. Perhaps our slowest rotator, however, is α Ori, now (cf [2009A&A...504..115K](https://ui.adsabs.org/2009A&A...504..115K)) assigned the period of 8.4 y. At the other extreme, Kaler suggests for ζ Aql A, α Aql, and ζ Lep, respectively, 16 h, at most 10 h, and around 6 h.

Radii (as distance from centre to outermost opaque layer, perpendicular to the axis of stellar rotation) are typically greater than the solar radius. Two notable instances of stellar expansion—in other words, of notably tenuous stellar atmosphere—are α Sco (with a radius of 3.4 AU, not far short of the Sun-Jupiter distance) and α Ori (with a radius of 4.1 AU or 4.6 AU from interferometry, or alternatively 3.1 AU or 3.4 AU from luminosity-temperature deductions). Results in these extreme cases depend strongly on the wavelength selected for evaluating opacity. Observations within Population P do indicate, although our sample S does not succeed in illustrating, the possibility of still more-extreme stellar radii, to values approaching ~ 10 AU. (Among these extreme-radius cases is a vividly red star well known to binocular-equipped observers, though a bit too faint for our table, μ Cep.)

The broad range of temperatures (a topic whose MK conceptual subtleties we examine in subsection 4.1, below) is reflected in the

fact that all of the Big Seven temperature-type bins in the traditional MK temperature sequence are well occupied, however statistically skewed (as we have argued above) is the distribution in the MK Big Five luminosity-class bins. At the MK temperature extremes are the hot ζ Pup (O5; 42,000 K) and the cool \omicron (omicron) Cet (M5–M10; a typical temperature for this variable is variously suggested as ~2000 K or ~3000 K).

Interesting spectral anomalies in Sample S include the “Be phenomenon” and “shell spectrum” stars, as discussed at length in our final subsection.

SECTION 3: Initial user guide to the columns in our 315-entry table

In our first column, we use the flags “+nP” (n = 1, 2, ...) for companions of sub-stellar mass, such as have been found outside our Solar System, in an accelerating sequence of discoveries, from the 1990s onward, that has now reached even the tiny Sample S. Such companions are typically planets but could in principle also be brown dwarfs. We do not attempt here to define formally the difference between a planet and a brown-dwarf companion.

In this same column, we apply the WDS naming scheme for multiplicity, both in the case of true binarity and in the case of mere optical doubles (in all but ten awkward cases, as noted in section 1 above, putting into the first column just the name of the brightest WDS-catalogued component; but we additionally try to supply particulars, at any rate in the online table, for binary and mere-optical companions brighter than mag. 10, in the “Remarks” column).

An overview of the WDS naming rules is now in order.

Suppose, as a hypothetical case, that a certain bright naked-eye point source has been familiar from Johann Bayer’s 1603 atlas onward as “omega FooBaris,” or ω FBr. Suppose ω FBr to have been discovered by some 1830s filar micrometrists to be a tight double, with components separated on the celestial sphere by an angular distance of 0.7”. It does not matter whether the pair is a true binary or a mere line-of-sight coincidence: in either case at the 1960s launch of WDS the pairing is catalogued as ω FBr A and ω FBr B.

Now suppose, as a refinement of this basic scenario, that around 1910, ω FBr A was found by some spectroscopist to be a spectroscopic binary (in our penultimate-column notation, to be an “SB”), and that nothing further was known about ω FBr A until 1974. What are the 1973 WDS implications of the 1910 discovery? Under WDS rules, ω FBr A had at that early stage in the development of WDS to be ω FBr A (not ω FBr Aab), since as of 1973 its components had not been individually resolved.

Stellar interferometry was launched in a modest way in the 1920s. It is perhaps reasonable to say that a “Second Generation” of optical interferometers was ushered in by the team of Robert Hanbury Brown, operating the Narrabri Stellar Intensity Interferometer from 1963 to 1974. Suppose, then, that in 1974 some interferometer, such as Narrabri, succeeded in resolving ω FBr A into two components, say at a separation of 0.1”. At this stage, the WDS multiplicity catalogue was at last able (and under its self-imposed rules was required) to refer not to “ ω FBr A” but to ω FBr Aa and ω FBr Ab.

Finally, suppose that in the current, arguably “Third,” generation of optical interferometry, some such instrument as CHARA or NPOI, perhaps working in the year 2020 or 2030, discovers ω FBr Ab to be itself a (very tight, very rapid) binary, with the separation even at apastron amounting to just a few tens of milliarcseconds. At this stage, WDS is able (and under its self-imposed rules is required) to refer not to ω FBr Aa and ω FBr Ab but, rather, to ω FBr Aa, ω FBr Ab1, and ω FBr Ab2.

Apparent Visual Magnitude ($m_v = V$): Apparent magnitudes, with “v” appended for large-amplitude variables, are from *HIPPARCOS*. In the case of variable, we take as authoritative the ranges (where possible, in V), and also the periods, published in the online AAVSO(VSX) database. Our reasoning here is that AAVSO has critically appraised and filtered data originally presented in more upstream sources, such as the primary (journal-article) literature. Our “V” is the usual “V” of UBV photometry, as introduced by H.L. Johnson and W.W. Morgan in [1953ApJ...117..313J](#). The (yellow) V filter corresponds roughly to the response of the eye. We retain, without having attempted our own independent error analysis, the assertion of our Handbook predecessor R.F. Garrison (working essentially before *HIPPARCOS*) that the “probable error” of each of our cited V values is at most 0.03 mag. (in other words, that of the actually and potentially available V measurements from the world’s duly competent photometry facilities, at least half will lie within 0.03 mag. of our own cited V values). Some small inaccuracies in magnitudes may be present in cases of combined light: readers needing confirmation may check our values against WDS, or where possible against the magnitude-specifying atlas pages of AAVSO. (By the nature of its mission, AAVSO is constrained to supply in its cartography not only details of variables, but also magnitudes of stars that are constant, and which can be used by amateur photometrists as comparison stars and check stars.) We hope to rectify these possible small inaccuracies in the next major revision of this document, in the “6.x.x” series, perhaps around 2021 Dec. 31.

Spectral Classification (MK Type): The “MK temperature type” (O, B, A, F, G, K, M) is given first, followed by a finer subtype (0-9) and an “MK luminosity class” (Roman numerals I-V, with “a” or “b” added occasionally to indicate slightly brighter or fainter stars within the class). As we discuss in detail in subsection 4.1 below, O stars are the hottest, M stars coolest; Ia stars are termed the most luminous “supergiants”; III stars are termed “giants”; and V stars are termed “dwarfs.” V stars form the largest class in the cosmos, comprising the observational Main Sequence (MS) (as a region in two-dimensional MK-luminosity-class-versus-MK-temperature-type classification space). Other MK symbols include “e” for hydrogen emission; “f” for broad, non-hydrogen emission in hot stars; “m” for strong metallic absorption; “n” or “nn” for unusually broad absorption; “p” for peculiarities; “s” for a mixture of broad and sharp lines; and “:” for a minor uncertainty. (The flags “n” and “nn” are a signature of rotation. It seems that historically “n” and “nn” signified “nebulous”, as references to the photographic-plate appearance of a rotationally broadened absorption line.) Where a single star (e.g. α CMa A) is given two types, with the second flagged “m”, the first is the type that best characterizes the hydrogen lines, the second the type that best characterizes the metal lines.

MK classifications are in some cases controverted. We have inherited our own types for the most part from the judgements of our

predecessor R.F. Garrison, who, as a principal historical authority in MK classification, drew both on what he judged to be the best of the literature and on some of his own unpublished classifications. As of 2021 Jan. 13, we have made a modest beginning at flagging the cases of controverted MK phenomenology (in our online, but not in our printed-edition, “Remarks” column), in two ways: (a) Where the literature suggests a real difficulty in MK classification, we draw attention to the difficulty, discussing it in a few words. (b) Where we have not found reason in the literature to suspect an MK-classification uncertainty, but nevertheless find our assigned MK type diverging (even in a small way) from the type assigned as of epoch 2021.5 in the official United States Naval Observatory and HM Nautical Almanac Office publication *Astronomical Almanac*, Section H (bright stars), we document the divergence, without further discussion.

Parallax (π), Proper Motion (μ), and Position Angle (PA): Parallaxes, in milliarcseconds (mas), proper-motion vector norms ($''/y$), and vector position-angles (degrees, from N through E) are derived from the *HIPPARCOS* 2007 data reduction, with a few exceptions. It may be hoped that in future years more precise parallaxes will be forthcoming from the *Gaia* mission, which has now found an engineering solution significantly easing its initial restriction to the fainter stars. (Detector overload had been feared.) Like *HIPPARCOS*, *Gaia* has to cope with the special challenges posed in measuring to high precision (i) the parallax of a (orbitally wobbling) star possessing a gravitationally bound, and not necessarily well documented, companion, and (ii) the parallax of a star with perturbed photosphere, and consequently with displaced photocentre (as when a tight binary system contains a bright mass-transfer stream).

Absolute Visual Magnitude (M_V) and Distance in Light-Years (D): Absolute magnitudes and distances are determined from parallaxes, except where a colon follows the absolute magnitude; in these cases, both quantities are determined from a calibration of the spectral classification. The absolute magnitude is left uncorrected for interstellar absorption. The appropriate correction is typically $\sim +0.06$ mag. per 100 ly outside the Local Bubble, i.e. beyond ~ 100 ly. A special difficulty, not fully grasped by us, arises in the case of the controverted ϵ Aur system distance (for which we now use *Gaia* DR2, additionally supplying references to the recent literature).

We take account of uncertainties in parallaxes by stating the derived distances, in ly, to no more than the appropriate number of significant figures (rounding where necessary). In cases where rounding would itself be misleading, we use a tilde as an indicator of imprecision.

Radial Velocity (V_{rad}): Radial velocities are from BSC5. “SB” indicates a spectroscopic binary, an unresolved system whose duplicity is revealed by periodic Doppler oscillations in its spectrum and for which an orbit is possibly known. If the lines of both stars are detectable, “SB2” is used; “+” and “-” indicate, respectively, motion away from and toward the observer. “V” indicates a variable velocity in a star not observable as a spectroscopic binary. (In most “V” cases, the orbit is unknown.)

Remarks: Remarks include data on variability and spectra, particulars of any companions, and (for the most part, only in our online table) prominent bits of observational-astronomy news. In a departure from our practice prior to 2017, we now give star names in all and only those cases in which star names are formally promulgated in the International Astronomical Union (IAU) star-naming project, as launched in 2016 at http://www.iau.org/public/themes/naming_stars. Readers requiring further information on names could start with the individual star descriptions in <http://stars.astro.illinois.edu/sow/sowlist.html>. Richard Hinckley Allen’s 1899 book *Star Names: Their Lore and Meaning* has been much cited over the decades. More recent scholarship, with due professional attention to Arabic philology, is, however, presented in Paul Kunitzsch and Tim Smart, *Short Guide to Modern Star Names and their Derivations* (Wiesbaden, 1986), and (by the same pair of authors) *Dictionary of Modern Star Names: A Short Guide to 254 Star Names and their Derivations* (Cambridge, MA, circa 2006). In the **Remarks** column, a **boldface** star name indicates a navigation star.

SECTION 4: Supplementary user guide, concerning the more detailed interpretation of our MK-classification column

SUBSECTION 4.1: Conceptual underpinnings of the MK classification system

In strict conceptual accuracy, the MK temperature types are a purely phenomenological record of which elements are present (a) in which stages of ionization, and (b) at what densities (in other words, under what local strength of the local downward-directed gravitational field), in the photosphere of the given star.

Decades before the 1943 Morgan-Keenan-Kellman publication of the full two-dimensional MK scheme, it had already been found possible to set up the phenomenological spectral types under our heading “(a)” in a single orderly OBAFGKM sequence, in which individual types gave way smoothly to their neighbouring types. (This process was itself not quite straightforward. First came a simple Harvard “A, B, C, D, ...” scheme. This was followed by the realization that “A,” for example, linked smoothly in its phenomenology with “B” and “F,” with some of the old alphabet having to be altogether dropped or repurposed. In working out this ordering, it was found necessary by the Harvard pioneers to subdivide the OBAFGKM categories, for instance in the sense of “G rather similar to F” and “G rather similar to K” and “G about equidistant between F and K.” Hearnshaw’s *Analysis of Starlight*, now in its second edition as [2014anst.book.....H](#), is the definitive history both of the MK scheme and of its predecessors.)

It was then not a matter of definition, but of astrophysical discovery (cf, e.g. the already-cited [2014anst.book.....H](#), or again [1994AJ....107..742G](#), or again the detailed MK reference-work exposition [2009ssc.book.....G](#)), that the OBAFGKM sequence corresponded to a temperature-ordered sequence of stellar groupings, running from the hottest photospheres to the coolest, with each of the various subdivisions within each of the O, B, A, F, G, K, and M types corresponding to a particular temperature range.

With the 1943 introduction of the two-dimensional MK scheme, the luminosity classes I, II, III, IV, V likewise had strictly a phenomenological, not an astrophysical, definition (proceeding now from our heading “(b)”, as opposed to the “(a)” that yielded O, B, A, F, G, K, and M). It was then once again conceptually speaking not a matter of definition, but of astrophysical discovery, that the I-through-V sequence corresponded to a luminosity-ordered sequence of stellar groupings, running from the most luminous to the least

luminous.

Admittedly, this conceptual picture, for the history of work under our heading “(b)”, is idealized. It was evident on the theoretical front even some decades before 1943 that the “(b)”-heading phenomenological features highlighted in 1943 by the developers of the MK taxonomic system, and signalling differences in photospheric gas densities (i.e. to differences in the strength of the local downward-pointing gravitational field) in fact correspond to differences in stellar luminosities. The developers of the MK taxonomy thus had a theoretical motivation for their definitions of classes I, II, III, IV, and V, resolutely phenomenological though their definitions were required to be, under observational-astrophysics methodology. — The MK system now serves as a paradigm of successful taxonomy, even for fields outside astronomy. A classification system is defined in terms of mere phenomenological fieldwork, and yet in the expectation (successfully realized in the case of MK) that the phenomenological classification bins will in due course be discovered by the theoreticians to correspond to relevant, important, physical differences in the materials observed. (Parallels might be suggested with, e.g. 18th- or 19th-century medicine: whereas (i) the old clinical-phenomenology definition of “tertian fever” and “quartan fever,” in terms of the observed duration of body-temperature anomalies, have been found in physiology theory not to correspond to useful fundamental realities at the level of microbiology, (ii) the gross empirical observation, as with the pre-Victorian stethoscope, of heartbeat anomalies has been found to correspond to useful fundamental realities at the level of cardiac neuroanatomy.)

When the MK system was introduced, it was already evident that if the classes I through V signalled a progressive decrease in stellar luminosities, then they had to signal a corresponding progressive decrease in stellar radii. The temperature of a given photosphere determines the amount of energy that photosphere radiates per unit time per unit of photosphere area. Consequently, if two stars in the same temperature type are found to differ in luminosity class, the one in the brighter luminosity class must have a larger total photosphere area, and so must be of greater radius.

It was therefore natural to adopt theory-informed, but nevertheless in official terms purely mnemonic, labels for the phenomenologically conceived luminosity classes, with I called for convenience the “supergiants,” II the “bright giants,” III the “giants,” and IV the “subgiants.” V had to be given some mnemonic label opposed to “giant,” with “dwarf” consequently pressed into service, and “subdwarf” used for the underluminous class VI (important in studies of congenital metallicity, but irrelevant to our own Sample S). (It is admittedly troublesome that the terms “white dwarf”—and nowadays also “brown dwarf”—prove necessary in other contexts, with the “white dwarfs” and the now-celebrated “brown dwarfs” radiating at luminosities far below even classes V and VI.)

SUBSECTION 4.2: MK classification and stellar evolution: preliminary remarks

In 1943, when the MK system was introduced, stellar-evolution theory was not yet on a sound footing. Only the broad outline, that a star may be expected to increase in photospheric radius after completing the fusion of hydrogen in its innermost portion, was at that point known. With the theoretical nuclear-physics advances of the 1950s and 1960s, and with the advent of increasingly detailed computer modelling from the 1960s onward, it became possible to map the elaborate excursions (we outline these in subsections 4.7 and 4.8 below) that evolving stars perform in the two-dimensional luminosity-class-versus-temperature-type phenomenologically defined MK plane (the “observational HR diagram”). In particular, it is now known that every star in the phenomenological class V in our 322-star set from our 315-entry table is still performing stable fusion of hydrogen in its innermost portion. (We repeat that this class V is best termed, with correct deference to the MK classification conceptual underpinning, not simply the “Main Sequence” (MS), but the “observational MS”—as at p. 342 of the authoritative [2006ima.book....C.](#)) Further, membership in the phenomenological class IV is a good (though even in our small 322-star set not an infallible) indicator that stable hydrogen fusion in the innermost portion is over, with the subject star now having performed at least some part of its (in general, elaborate) later-life excursions over the MK phenomenological plane.

The distribution of the set of 322 stars across MK luminosity classes I through V accordingly turns out to be a reasonable indication of the evolutionary spread of the set.

It follows that the naked-eye bright-star night sky is a different place from the daytime sky, with its lone proximate class-V star. Something on the order of a mere fifth of our 321 MK-classified bright nighttime stars (for the most part stars in luminosity class V) resemble the Sun (the sole daytime object in our set of 322 MK-classified bright stars) in stably burning hydrogen at their centre. Even most of these are far hotter than the Sun and are consequently destined to spend less time than the Sun in in this process of stable burning. All the rest have in one way or another moved beyond that stage, as shown by their luminosity classes—with the nocturnal 321 falling overwhelmingly into classes III and IV, but with classes I and II also rather well populated.

SUBSECTION 4.3: MK classification and stellar evolution: starbirth and MS

A star has at birth (i.e. has upon condensing sufficiently from its local ISM cloud to begin hydrogen fusion) four key characteristics. If the star happens not to be in the disturbing environment of some proximate star (most notably, in the disturbing environment of a binary companion so close as to transfer matter) then these four characteristics jointly entail its various other characteristics, for each point in its entire subsequent career. Prominent among those other characteristics are the duration of overall life, and at each point in the overall life additionally those time-varying key characteristics, which are radius, luminosity, and its photosphere effective temperature. Here, then, are the “Governing Four”: (a) birth-epoch mass (the more massive stars are also the hotter, the more luminous, and the shorter-lived); (b) birth-epoch elemental composition (the most important aspect of composition is simply the birth-epoch “metallicity”—i.e. the extent to which, thanks to the specific properties of the local gestating ISM cloud, the subject star contains at the time of birth any elements, in whatever detailed proportions, heavier than hydrogen and helium); (c) absence or

(possible) presence of inherited fossil magnetism, from (possible) magnetism in the gestating ISM cloud; and (d) birth-epoch speed of rotation.

Of the four listed properties, the first is the most important, accounting, along with the accidental circumstances of distance-from-Earth and time-elapsed-since-gestation, for essentially all the stellar variety that the naked eye can discern.

Regarding the accidental circumstance of time-elapsed-since-gestation, a parenthetical caveat, relevant even to interpreting the casual naked-eye experience, is needed: stars condensed from the same ISM cloud are of the same age. This is the case not only with binaries but also, more dramatically, with associations (such as the dramatic naked-eye association in the northern sky whose most familiar members comprise β UMa (Merak), γ UMa A (Phecda), δ UMa A (Megrez), ϵ UMa A (Alioth), and ζ UMa Aa (Mizar), in other words comprise all but the first and last of the seven Big Dipper stars).

In contrast with mass and present age, congenital elemental composition does not vary greatly across our set of 322 MK-classified bright stars. The pronounced chemical differences across the set of 322 (evident from the notations for chemical peculiarities in many of the 322 bright-star MK types in our 315-entry table) are due, rather, to processes of stellar aging, notably (i) gravitational settling and radiational lofting of selected elemental species in cases in which the outer layers are quiet (in particular, not rotationally disturbed), and (ii) processes known as “Dredge-Up” (discussed again in subsection 4.8, below), when convection in an evolving star raises such elements as carbon or nitrogen into the photosphere from the buried thermonuclear furnaces.

We will not attempt to discuss congenital magnetism. But we do remark that like chemical peculiarities, magnetism can develop and change as a star ages (with, for instance, convection in outer layers, under rotation, producing a dynamo, and with the dynamo in turn generating the kind of looping-field locally dipolar magnetic structures present in the Sun, and hinted at in the small telescope by the Sun’s appearance through a hydrogen Balmer- α filter).

The fourth property in our list, congenital rotation, is a consequence of the vagaries of possible motions in the gestating ISM. The local part of the condensing gas was likely to have some kind of coordinated spin, and this spin tended to increase, under conservation of angular momentum, as the gas became more and more condensed—even though some angular momentum also was possibly shed via gas outflows, as the condensation proceeded toward starbirth.

We will not discuss congenital rotation further. We do, however, remark that the rotation speed of a solitary, undisturbed star is once again a property that can evolve as the given star ages, under the combined influence of its evolving mass distribution (although the mass of all but the hottest stars remains rather constant until late in life, after cessation of core hydrogen fusion the mass gets distributed over larger radii, forcing (under conservation of angular momentum) an increase in rotation period) and its (possibly, as already noted, evolving) magnetism.

The process of change has two aspects. On the one hand, as an aging star evolves out of luminosity class V into IV, III, and in the case of congenitally massive stars even into II or I, increases in its radius cause (because angular momentum is conserved) a slowing of rotation.

On the other hand, and quite apart from this general slowing-through-bloating, a spin-braking mechanism exists within class V for those stars that succeed in generating the right kind of local, looping, dipole magnetic-field structures. The mass shed by such a star in winds, although modest, is nevertheless constrained by magnetic fields not to orbit the star freely, but to rotate at the about the same angular velocity as the star itself. Under conservation of angular momentum, this so-called “magnetic braking” then slows the rotation. In the overall galactic population of V stars, those cooler than MK temperature type F5 are capable of achieving magnetic braking, and those hotter than F5 are not. The F5 type thus constitutes a so-called “rotation break” within class V.

In our set of 322 MK-classified bright stars, all but six of the class V stars lie on the hot side of the break. The brightest V-class stars in Earth’s sky have to be either the most luminous, and therefore the hottest, or those nearest to Earth. The scarcity of V-class bright stars on the slow side of the rotation break therefore indicates that it is the first of these two brightness-promoting characteristics that predominates, in our overall set of 322.

Although we here largely neglect stars in the disturbing environment of other proximate stars, we do have to remark parenthetically that in the case of a close binary, rotation (like also chemical composition) can be affected by processes of mass transfer. This is very notably the case with one of the more heavily studied stars in the 322-member set, α Leo A (Regulus). Here the rapid rotation is the result of a now-completed spinning-up process, involving a copious mass transfer, from the now diminutive, and therefore now observationally elusive, pre-white dwarf. In the “Remarks” for α Leo A in the table, we point out that this elusive companion, having for decades escaped observation, is at last reported in [2020ApJ...902...25G](#) as detected spectroscopically.

The F5 “rotation break” within MK luminosity class V is ultimately due to, and is nearly coincident with, a transition (as one proceeds along the observational MS from the hottest stars to the coolest, i.e. as one advances in the sense OBAFGKM) from stars in which the hydrogen fusion is predominantly the work of the carbon-nitrogen-oxygen (CNO) cycle to stars in which the hydrogen fusion is predominantly the work of the proton-proton chain. The point at which the two processes deliver, per unit of fusion-depth mass, roughly equal energy-per-unit time is at or near a total stellar mass of $1.2 M_{\odot}$.

To what extent are the four key properties reflected in the MK type of a young star (in observational terms, a star found to lie in MK luminosity class V)?

(a) Mass is well correlated with MK temperature type, in the sense that the OBAFGKM progression within class V proves to be a progression from the most massive stars to the least massive. This fact is itself far from obvious. It was, however, established in the early decades of the 20th century by spectrally classifying the elements of binary systems, of known distance, in which the orbit is not so tight as to allow the disturbing feature of mass transfer, and yet in which the orbit is tight enough, and consequently fast enough, to permit determination of orbital geometry and orbital period. For such binaries, individual masses can be determined from Newtonian mechanics.

(b) Birth-epoch elemental composition is not really reflected in the observationally assigned MK class. We have already remarked that the elemental-composition flags present in many of the 322 bright MK types are due, if not to “Dredge-Up” in the case of an aging star, then to segregation of elements through gravitational settling and radiative lofting (processes that can occur even for a

young star, provided its atmosphere is quiet, as in cases where rapid rotation is absent).

(c) The MK scheme does not attempt to flag magnetism, even though magnetism is observed spectroscopically, through the Zeeman splitting of emission and absorption lines when a magnetic field is strong.

(d) Rotation can be inferred in favourable cases, but not in all cases, from the presence of the MK-type flags “n” and “nn.” In a favourable case, a rapidly rotating star is seen more or less equator-on, causing its emission and absorption lines to be Doppler-broadened (since half of the photosphere is rapidly receding from the spectrograph, and the other half rapidly approaching it). In, however, the unfavourable case in which the star is seen more or less pole-on, there is no rotational broadening. A particularly well-known example of a rapid pole-on rotator (with “n” and “nn” therefore absent from the observed MK type) is α Lyr A (Vega).

We might add by way of background that it is only in recent decades that the detection of pole-on rotators has become feasible at all. If the star is close and bright enough, interferometry, while powerless to detect the shape deformation of a pole-on rapid rotator, may nevertheless succeed in picking up the equatorial darkening that accompanies rotational flattening (in the pole-on case, as an anomalous darkening, over and above the normal “limb darkening,” toward the edges of the interferometrically discerned stellar disk, at whose centre is the Earth-facing stellar pole).

SUBSECTION 4.4: MK classification and stellar evolution: rotation largely neglected here

It is now helpful to outline the various possibilities for stellar evolution, as experienced by that majority of stars in the 322 MK-classified set that are already in MK luminosity classes IV, III, II, or I, as opposed to the “observational MS” which is class V. But an initial caveat is needed: we here largely neglect the disturbing influence of stellar rotation, important though that influence is.

Regarding rotation, we do remark at this point that rotation can produce flows of matter along lines of stellar longitude (“meridional flows”), and that where such flows extend some significant distance into the stellar interior, they help replenish the supply of hydrogen, as a thermonuclear fuel, in the stellar depths. The effect of rotation is in general to somewhat shift the evolutionary track of a star on the phenomenological MK plane (by promoting mixing of stellar layers that would otherwise be more sharply separated) without radically changing the shape of the track.

Difficulties in constructing an evolutionary model for the interior of a rapid rotator are among the themes of Section 1 in [2011ApJ...732...68C](#). This same paper discusses difficulties involved in deducing the mass and age of a rapid rotator, and the problem of deviations from the von Zeipel 1925 gravity-darkening law for oblate-spheroid stars. The law would give the correct result for gravity darkening if the flattened star had a purely radiative envelope. With rotation, however, gravity darkening can lower the photosphere effective temperature at the equator, causing convection to set in there even when the envelope is radiative at the poles. In our 322-star set, this pathology is present in at least α Aql A (Altair) and α Cep A (Alderamin).

Even where the convective regime is uniform, it is not possible to assign a single photospheric effective temperature to a rapid rotator: its observed MK temperature type is now a mongrel, the result of light entering the spectrograph from the differing temperature regimes of (hot) poles and (cool) equator.

SUBSECTION 4.5: MK classification and stellar evolution: structure, energy flows

As a further preface to details of evolution, it is now necessary to introduce discussion-guiding concepts of stellar structure and stellar energy flows.

A star still stably fusing hydrogen in its innermost portion (whether predominantly via the CNO cycle or predominantly via the proton-proton chain) is said to have a hydrogen-fusing “core.” The layers outside the energy-producing “core” of such a star are said to comprise its “envelope.” Under this definition of “envelope,” the envelope is not a place of energy generation, but merely a place of energy transport. This transport involves a cascade, in which a single core-produced photon is absorbed by some envelope atom, causing the envelope atom to re-radiate multiple photons, each individually less energetic, and with the same aggregate energy as the now-vanished input photon. Each of these less energetic photons is in turn absorbed by some envelope atom in a still higher layer, which in its turn re-radiates a plurality of correspondingly less energetic photons. Eventually, as that outer-skin part of the envelope that is the photosphere is reached, photons begin travelling freely, without processes of absorption and re-radiation.

Those young stars with cores hot enough to have the CNO cycle as their principal mode of hydrogen fusion have convective cores. In the case of the very hottest O stars (perhaps hotter than any of the 35 or 40 or so O stars in our set of 322 MK-classified bright stars), not only the core but even the envelope is convective. The more usual case, however, for a CNO-dominated star, and perhaps the only case appearing for the CNO-dominated subset of our 322-star set, involves a convective core overlain by a radiative envelope.

Where the temperatures at the core are low enough for the proton-proton chain to predominate, the core of a young star is radiative. High envelope opacities in this low-temperature case make radiation an inefficient mode of energy transport, causing envelopes to be convective. As one advances along the temperature sequence in the sense OBAFGKM, stars at first present just a thin convective layer (setting in at a photosphere effective temperature of ~ 8300 K), with convection then running deeper and deeper (and in particular, in the case of our own Sun, as a G2V star, pervading the entire envelope).

Here (once again) a caveat is necessary regarding rotation. A rapid rotator can straddle the ~ 8300 K boundary, with convection absent at its (hot) poles, and at least a thin convective layer present at its (cooler) equator.

As an irony of nature, an extreme case exists at the cool end of the OBAFGKM progression, just as for its already-discussed hot end. In the coolest young M stars, convection extends all the way down to the core. As for the extreme O stars, so also, however, the extreme-M case is irrelevant for us: our set of 322 MK-classified bright stars contains no young M stars at all.

SUBSECTION 4.6: MK phenomenology of early evolution within the theoretically defined MS

Having so far mentioned just the “observational MS,” we may now proceed to the theoretical definition of the MS, or more strictly of departure-from-MS (and soon we shall also be relating this bit of theory to the already-presented observational MS concept). The theoretical MS will turn out (subsection 4.8, below) to be defined in such a way that departure perhaps can occur already within class V, but can also be delayed until an aging star has brightened enough to take it into class IV.

It is a sufficient, although not a necessary, condition for a star lying within the theoretical MS that it be still fusing hydrogen within its core.

Even within this early, seemingly placid, stage of a star’s life, large changes can occur. While our own Sun has another four or five gigayears of life before its core-hydrogen fusion is over, the placid process of early-MS evolution will after just a single gigayear already drive its luminosity high enough to destroy Earth’s biosphere.

At the heart of this early-MS process is a gradual change in core composition, as helium ash accumulates. With the core becoming progressively helium-richer even while core hydrogen nuclei continue to fuse, the number of particles constituting the aggregate of gas that is the core progressively falls. Given this rise in the mean mass of the core-gas particles (the free electrons, and a diminishing number of hydrogen nuclei, and a rising number of helium nuclei: but the increased helium comes at the expense of the hydrogen, with two hydrogens yielding one helium) the core, while maintaining the pressure needed to support the overlying envelope, is under the Ideal Gas Law forced to contract. Under a physical principle known as the Virial Theorem, half of the gravitational potential energy liberated by the contraction is translated into thermal energy, i.e. into a rise in the temperature of the core. With this rise in temperature, core hydrogen fusion (a process already decidedly dependent on temperature in the case of the proton-proton chain, and very strongly dependent on temperature in the case of the CNO cycle) becomes more vigorous. As a result, the star overall becomes more luminous, and also experiences a modest increase in radius.

It is now convenient to distinguish in our set of 322 MK-classified bright stars between (A) the very massive ones (possessing at birth a mass greater than around $8 M_{\odot}$ or $10 M_{\odot}$) and (B) all the others. The very massive stars are destined to die as supernovae (leaving behind perhaps a black hole, perhaps a neutron “star”). The others are destined to die as white dwarfs.

SUBSECTION 4.7: MK phenomenology of evolving high-mass stars (eventual supernovae)

In observational terms, the very massive MS stars are of MK temperature class O, or else of the hot B subdivisions B0, B1, or B2. In our set of 322 MK-classified bright stars, at least the following ten (in order of increasing RA) can be said with confidence to meet this condition: η Ori Aa (B0.5 V), θ Car (B0.5 V), α Cru B (B1 V), β Mus Aa (B2 V), π Sco A (Fang; B1 V), β Sco Aa (Acreb; B0.5 V), τ Sco (Paikauhale; B0 V), ζ Oph (O9.5 V), α Ara A (B2 V). Additionally, 31 are observed to be on the borderline for meeting this condition (being in IV, or being classified “IV–V,” or being of MK temperature class B2.5).

In the process leading up to the supernova climax, these massive stars will eventually rise in observational terms into the MK “supergiant” luminosity class I. In the set of 322, 35 are clearly now at that late stage in their development.

We will not discuss at any length the details of massive-star evolution once core hydrogen is exhausted, instead contenting ourselves with just five brief points:

(i) The very concept of MS is a little misleading for the most extreme of the massive stars, since in the most extreme cases scarcely has starbirth (the commencing of core hydrogen fusion) been achieved before gross observable evolutionary changes have set in. We will not here attempt to chart this territory (and in particular will not attempt to define for this group of stars the tricky theoretical concept of “departure from MS”). We remark only that a safe early-life theoretical concept for the most massive stars is the concept of a mere instant, as opposed to an interval—namely arrival on the “Zero Age [Theoretical] MS,” as the instant at which core hydrogen fusion starts.

(ii) In their so-short lives, these very massive stars fuse progressively heavier elements, in a central aggregation and in shells overlying the aggregation. The fusion after helium is finished is fuelled first by carbon, then by oxygen and neon and magnesium, and finally by sulphur and silicon, yielding the eventual dumping of iron ash, from sulphur-silicon burning in a shell, onto a growing inert central aggregate of iron.

(iii) A “core-collapse” supernova eventuates after the iron aggregate exceeds the “Chandrasekhar limit” of $\sim 1.4 M_{\odot}$

(iv) The complexities of core and shell burning, with burning at various levels switching itself on and off in the process leading up to the supernova, translates in observational terms into movements across the MK luminosity-class-vs-temperature-type surface, with luminosity not changing much, but with temperature type changing dramatically (and with changes possible both in the redward, or OBAFGKM, sense and in the blueward, or MKGFABO, sense). Each of the MK types OBAFGKM is represented in our group of 35 supergiants, with at the hot (blue) extreme ζ Pup (Naos; O5 Ia) and ζ Ori Aa (Alnitak; O9.5 Ib), and at the cool (red) extreme α Sco A (Antares; M1.5) and α Ori Aa (Betelgeuse; M2 Iab).

(v) In its redward or blueward progressions, an evolving supergiant can pass, possibly more than once, through the “Instability Strip” (IS) in the luminosity class-vs-temperature type MK plane, thereby temporarily becoming a Cepheid variable. This possibility is presently actualized in our set of 35 class-I stars by (in order of increasing RA) α Umi Aa (Polaris), β Dor, I (ell) Car, η Aql A, and δ Cep A.

SUBSECTION 4.8: MK phenomenology of evolving lower-mass stars (eventual white dwarfs)

(B) We may now proceed to explain the sense in which, extreme cases of lower-mass cases of rotation aside (where rotation yields gas flows so violent as to leave no gas unmixed), all stars in the 322-star set with masses below $\sim 8 M_{\odot}$ or $\sim 10 M_{\odot}$, and not disturbed by mass transfer from some companion star, proceed from a readily definable theoretical-MS interval of life to the theoretical Sub-Giant Branch (SGB), then to the theoretical Red Giant Branch (RGB), then to either the theoretical Horizontal Branch (HB) or the theoretical Red Clump, then to the theoretical Asymptotic Giant Branch (AGB), and finally (as almost-corpses or corpses) to a post-theoretical-AGB phase, which, in the fulness of time, yields a white dwarf.

It might seem natural to set up a definition of “theoretical MS” for our eventual-white-dwarf stars on which such a star is deemed to leave the theoretical MS upon finishing core hydrogen fusion. The definition actually employed is, however, different (Carroll-and-Ostlie 2006ima.book....C, pp. 452, 453). The definition actually employed has (surely?) been motivated, over the past few decades of theory construction, by a desire to make the theoretical-astrophysics demarcations correspond as closely as possible to the actual spectrograph-observable changes of direction (i.e. to the actual observed bends) as a star traces its path, over a span of megayears or tens or hundreds or thousands of megayears, on the phenomenological I-through-V vs O-through-M surface. Under the standardly employed definition, a star is said to remain on the theoretical MS not only through the process of luminosity increase attributed in Subsection 4.6 to the Ideal Gas Law, but somewhat later, even a little after the depletion of core hydrogen has brought core fusion to a halt.

The matchup of theory and phenomenology is, despite efforts at fine-tuning the theoretical definitions, imperfect. Awkwardly enough, not only can a star be on the theoretical MS even after finishing core-hydrogen fusion: conversely, a star can even have left the observational MS, in other words can have left the MK luminosity class V, while residing so far within the theoretical MS as to be still burning its core hydrogen. In terms of our table, this awkward converse possibility is illustrated by at least the following (in order of increasing RA): χ Car (B3 IV (p?)), λ UMa (Tania Borealis; A1 IV), β Cru A (Mimosa; B0.5 III), ν Cen (B2 IV), ζ Cen (B2.5 IV), ι Lup (B2.5 IVn), α Tel (B3 IV), and the celebrated variable β Cep Aa (Alfirk; B1 III). Additionally, α Lyr A (Vega) is still far within the theoretical MS, and yet might erroneously be thought to have evolved to the edge of the observational MS, since its MK class is A0 Va. Here the cause of the “Va,” as distinct from “V,” is rotation (with Vega presenting itself to the spectrograph pole-on while rotationally flattened, in other words presenting a misleadingly increased radius).

At the moment at which the depletion of core hydrogen has brought core fusion to a halt, the luminosity of the star derives from fusion in a core-surrounding hydrogen shell, now raised to a fusion-capable temperature by the increased temperature of the inactive helium-ash core. For some modest time after core-hydrogen fusion has ceased, nothing dramatic happens from an observational MK standpoint. Departure from the MS is defined as occurring when the central deposit of non-fusing helium ash becomes so massive as to trigger a rapid internal reorganization of the star, with one or the other of two possible types of rapid contraction, to be distinguished below as “(B.a)” and “(B.b).” This is the point at which something MK-noteworthy, i.e. something that registers strongly in the spectrograph, finally happens.

(B.a) For stars in the 322-star set of mass below $\sim 1.25 M_{\odot}$, the growing central deposit of still-inert helium ash becomes so massive as to trigger a further, this time rapid, contraction of the core. Some of the gravitational potential energy present before the abrupt contraction, and now liberated by infall, is under the Virial Theorem translated into an increase in the thermal energy of the shell (in which fusion of hydrogen is therefore in turn speeded up). Paradoxically, although the core has decreased in radius, the rise in temperature of the shell causes the shell to expand, increasing the radius of the star overall.

Two contending factors are now at work. On the one hand, the star has become more luminous. On the other hand, it is now larger. The latter factor outweighs the former, entailing a fall in the photosphere effective temperature. (Total luminous output from the photosphere is determined both by the attained photosphere effective temperature and by the attained photosphere radius, i.e. by the extent of stellar bloat. If the overall radius increase is large, then a reasonable modest increase in total luminous output has to be accompanied by a temperature decrease.)

In MK observational terms, the star, now defined to have departed the theoretical MS and simultaneously arrived on the theoretical SGB, has on the one hand moved some modest distance upward out of luminosity class V, and has on the other hand advanced redward, i.e. has evolved in the sense OBAFGKM.

(B.b) For stars of mass above $\sim 1.25 M_{\odot}$ (and nevertheless not, we repeat, attaining the $\sim 8 M_{\odot}$ or $\sim 10 M_{\odot}$ threshold that makes an eventual supernova possible), the star is found under computer modelling to undergo a more radical internal reorganization. On this more radical scenario, not just the inactive helium-rich core, but the entire star, suffers a rapid contraction. It is this spectrograph-detectable event that is in the “(B.b)” case taken to define the end of the theoretical MS phase.

As in the less radical “(B.a)” scenario, the star increases in luminosity, with some of the liberated pre-contraction gravitational potential energy once again translated into an increase in temperature (with, once again, a consequent speeding up of hydrogen fusion in the shell). In contrast with the “(B.a)” scenario, however, the star is of a reduced radius overall. Under the unavoidable correlation of overall luminous output with both attained photosphere effective temperature and attained photosphere radius, the now shrunken, and yet now brightened, photosphere must now be of a higher temperature. In MK observational terms, the star therefore now quite abruptly not only advances upward in the V–IV–III–II–I sense, but also advances blueward, i.e. evolves in the sense MKGFABO.

Whereas in scenario “(B.a),” the star is said to arrive on the theoretical SGB simultaneously with its departing the theoretical MS, in the “(B.b)” scenario now under consideration arrival on the theoretical SGB is defined as occurring just a little later than departure from the theoretical MS, with a further episode of core contraction following the overall contraction that under “(B.b)” defines departure from the theoretical MS. This further episode of core contraction yields a cooling of the photosphere, and consequently a spectrograph-observable change in the sense OBAFGKM.

In scenario “(B.a),” i.e. for stars exceeding $\sim 1.25 M_{\odot}$, movement through the SGB is rapid, making the detection of such stars statistically improbable, and generating the so-called “Hertzsprung Gap” in HR-diagram plots of same-age stars when the subject

population is so selected as to be duly rich in masses exceeding $\sim 1.25 M_{\odot}$, and duly rich both in observational-MS stars and in observational-RGB stars. (Many open clusters meet this sampling requirement.) The statistical improbability notwithstanding, our 322-star set does succeed in capturing several fleeting residents of the Hertzsprung Gap, at any rate (in order of increasing RA) α Aur Ab (the close Capella companion), ε Leo, ζ Leo A (Adhafera), α UMa A (Muscida), and ζ Her A.

From this point onward, it is no longer necessary to distinguish scenarios “(B.a)” and “(B.b).” Under both scenarios, residency on the SGB (admittedly started, as we have just said, in one way in the “(B.a)” scenario, in a different way in “(B.b),” with residency in the former case brief) in due course yields a cooling of the photosphere. With this cooling, the photosphere opacity rises, causing not only the photosphere-proximate layers but even much of the deeper interior to convect. Since, however, convection is a markedly efficient mode of energy transport, the star becomes progressively more luminous and larger, while keeping its photosphere effective temperature roughly constant. As this observationally dramatic increase in luminosity starts, the star is defined as leaving the theoretical SGB and (simultaneously) arriving on the theoretical RGB.

As in the late phases of theoretical MS life, and as in the theoretical SGB, so also here on the theoretical RGB, the star is fusing hydrogen in a shell overlying an increasingly massive, although still inactive, central ball of helium. Now, however, luminosity is much higher than in the MS and SGB phases. As the still-inactive central helium ball increases in mass, it gradually contracts under its own weight. Some of the gravitational potential energy thus liberated once again becomes thermal energy in the ball, as dictated by the Virial Theorem. With the helium ball now getting gradually hotter, the overlying hydrogen-fusing shell becomes gradually hotter also, producing in turn a gradual speeding-up of its hydrogen fusion, and therefore a gradual increase in the star’s (already high) luminosity.

RGB life comes to an end with one of two possible kinds of transition to core helium burning, both entailing a decrease in overall luminosity and yet without much change in photosphere temperature. The transition is violent in the case of the less massive stars in our set, less violent in the case of the more massive stars in our set: we again omit details. The core-helium-fusion phase is analogous to, and yet is briefer than, the core-hydrogen fusion that characterizes the earlier part of the theoretical MS. The exact destination of this transition depends on whether the star was at the time of its birth (its arrival on the theoretical MS) metal-poor or metal-rich.

For a star born as metal-poor, exit from the RGB takes it rapidly to the “theoretical HB.” This region of the theoretical luminosity-vs-photosphere-effective-temperature plot corresponds to a long, roughly horizontal, roughly straight locus of points, which we might term the “observational HB,” on the MK surface. Since globular clusters are metal-poor, the observational HB becomes prominent when a globular is (at least partly) resolved into its constituent stars, for which spectroscopy then yields individual MK types. Different metal-poor stars switching on their core helium fusion are found to arrive at different points on the observational HB, i.e. to attain different photosphere effective temperatures. The particular attained photosphere effective temperature is found in computer modelling to depend chiefly not on the mass of the newly ignited helium core (this proves on modelling to be rather constant across the metal-poor population), but on the mass of the outer, non-helium, layers.

However, with just two or three or so known exceptions—the most celebrated of these being α Boo (Arcturus)—our 322 MK-classified bright stars are metal-rich. Moreover, the exceptions in our set of 322 are perhaps all at phases of evolution either preceding or following residency on the theoretical HB and observational HB. We will therefore not discuss the HB further.

For a star born as metal-rich, exit from the RGB, i.e. the switching on of core helium fusion, involves a rapid transition to the theoretical and observational “Red Clump” (in effect the redmost rump of the grander theoretical and observational HB), as further discussed at, e.g. https://en.wikipedia.org/wiki/Red_clump. Since the Red Clump is the helium-fusion analogue of theoretical-MS core-hydrogen fusion, it is unsurprising that it is followed by an evolutionary phase analogous to the observational RGB and theoretical RGB, namely the observational AGB and theoretical AGB.

On the AGB, helium core fusion has come to an end, with the star at this late stage in its life harbouring an inert core rich in carbon and oxygen. Fusion now proceeds, simultaneously or alternately, in an inner shell of helium and an overlying (and in terms of overall luminous output, for most of the AGB lifetime dominant) shell of hydrogen. With more than one shell in play, evolution becomes rather elaborate. In particular, it is possible for the helium shell to be temporarily inactive, simply accreting mass from the helium ash being dumped on it by the overlying hydrogen shell. Once the helium shell becomes sufficiently massive, it turns on helium fusion, causing the overlying hydrogen shell to expand and briefly switch off. The net result of this is a temporary drop in the luminosity of the star, until the helium burning in turn subsides and the hydrogen burning resumes. In its overall evolution along the AGB, and in its post-AGB transition to the quiet, dead state of a white dwarf, a star can undergo even many tens of such “helium shell flash” episodes. Additionally characteristic of evolution on the AGB are pulsation and mass loss. The possibility is dramatically illustrated in our 322-star set by \omicron (omicron) Cet Aa (Mira).

We will skip over the further details of stellar evolution toward the white-dwarf corpse phase, remarking here only that in the case of a star nearly, but not quite, massive enough to die as a supernova, even carbon may be fused before all thermonuclear activity finally ceases.

Two concluding remarks are now in order.

(1) Mention has already been made of “Dredge-Up” as a process affecting the elemental composition of the spectroscopically observed photosphere. In terms of the concepts now laid out, it can be added that “Dredge-Up” may occur in the violent and deep convection of the RGB, as “First Dredge-Up” (FDU), or after the RGB, as “Second Dredge-Up” (SDU) and “Third Dredge-Up” (TDU). A highly evolved star may experience more than one episode of TDU, and it is also possible for FDU and TDU to occur without SDU. Our table cites α Tau A (Aldebaran) as a star that has undergone FDU. On the other hand, our table in its present state of development does not cite instances of SDU or TDU.

(2) The deducing of a star’s evolutionary stage from its observed MK type, as it makes its way off the MS toward, eventually, the AGB, is not always straightforward. In the case of the most massive stars (with masses greater than $\sim 8 M_{\odot}$ or $\sim 10 M_{\odot}$, and with death-by-supernova therefore impending, and with temperature evolution late in life at one or more stages proceeding in the sense OBAFGKM and at one or more stages proceeding in the contrary sense MKGFABO), temporary observed residence, as a Cepheid

variable, on the Instability Strip (IS) raises the question (not always easy to answer) “Is this star making a first, a second, or a third crossing of the IS?” As pointed out in the table “Remarks,” this problem complicates, in particular, the analysis of that rather untidy Cepheid variable that is α UMi Aa (Polaris). For those stars massive enough to achieve core helium fusion at some point in their lives, and not so massive as to die a supernova death (a condition met in our 322-star set by all the stars below $\sim 8 M_{\odot}$ or $\sim 10 M_{\odot}$), it sometimes proves difficult to distinguish residency on the theoretical RGB, residency in the theoretical Red Clump, and residency in the theoretical AGB from the available spectroscopy. Indeed the theoretical “Asymptotic Giant Branch” is so named because it corresponds in observational terms to a locus of MK-surface points running perilously close to, so-to-speak asymptotically approaching, that just slightly redder locus of points that is the observational RGB.

SUBSECTION 4.9: Supplementary remarks on rotation with “Be phenomenon” and “shell” (in MK types O, B, A)

Some of our 322 bright MK-classified B stars have an “e” flag, for emission lines in spectroscopy. Some, and yet not all, such cases involve the important, and not yet well understood, “Be phenomenon.” Strictly speaking, the presently known “Be-phenomenon” stars in our set of 322 are at least the following 19 (in order of increasing RA): γ Cas A, α Eri (Achernar), ϵ Cas, η Tau Aa (Alcyone), η Ori Aa, ζ Tau (Tianguan), α Col A (Phact), κ CMa, β CMi A (Gomeisa), ω Car, ρ Car (HR4140), γ UMa A (Phecda), δ Cen Aa, μ Cen Aa, η Cen, δ Sco A (Dschubba), α Ara A, ζ Oph, and β Cep Aa (Alfirk). As we discuss again below, β Lyr Aa1 (Sheliak) may or may not constitute a twentieth case, and some doubt hangs additionally over γ Ara A (in our treatment, not a “Be phenomenon” star, because too evolved; but perhaps we are wrong). Further, closely related to the Be phenomenon is the spectroscopic (predominantly B-star) “shell” phenomenon. The amateur-spectroscopy essay in the Handbook current printed editions notes that the spectroscopic-shell phenomenon, and by implication the Be phenomenon, is a potentially fertile field for amateur spectroscopy. We accordingly supply here a general briefing on the Be phenomenon and its “shell” associate, highlighting the connection of both Be and shell with the often-troubling topic of rotation.

Although many of the most tempting amateur targets in the Be-phenomenon and “shell” fields are members of our 322-star set, we nevertheless discuss the Be and shell phenomena for the most part in general terms, without restriction to the set of 322. We hope thereby to maximize the value of our discussion, and in particular to stimulate an interest in Pleione, as a Be and sometimes-“shell spectrum” star not much fainter than our mag. ~ 3.55 cutoff.

Of all the non-cluster B stars in the galaxy, about 17% at some point in their lives present the “Be phenomenon,” with the phenomenon more prevalent at the hotter (near-O) than at the cooler (near-A) end of the B range. Within the overall set of galactic stars, the exceedingly rare O stars are known to sometimes present the same phenomenon (with the term “Oe” star therefore used occasionally in the literature). In our set of 322, ζ Oph, as an O star with a photosphere almost, and yet not quite, cool enough to entail classification as a hot B, is an instance. Also within the overall set of galaxy stars, some A stars are known to present the Be phenomenon. Again, our 322-star set furnishes an instance, namely γ UMa A (Phecda): this star is of MK temperature type A0, and so is just barely cool enough not to fall into the B classification bin. Nevertheless, since the phenomenon (which we will soon describe in proper detail) occurs predominantly in the B stars, the term “Be phenomenon” is standardly applied to stars in all three of the O, B, and A observational MK temperature types.

Several qualifying comments are now necessary.

The Be-phenomenon stars are not to be confused with the “Herbig Ae/Be ‘stars.’” The latter are not stars in the strict sense, but instead are contracting starlike bodies that have not yet achieved starbirth, i.e. have not yet started core hydrogen fusion. In their present stage of development, they are continuing to heat up under gravitational contraction, and are (unsurprisingly for objects condensing out of ISM clouds) embedded in circumstellar dust.

A true “Be phenomenon star” need not currently have emission lines in its spectrum. It must, on the other hand, be known to have at some point in its past presented emission. In observational practice, the emission is always found to occur in at least one or more lines of the hydrogen Balmer series.

The condition of past-or-present emission, while necessary, is not in its turn sufficient. A supergiant in MK type B, with Balmer emission, is not a Be-phenomenon star. For a star to be Be-phenomenon, it must either lie on the theoretical MS or (as in the case of Be-phenomenon ζ Tau (Tianguan) in our table, observationally in MK luminosity class III) be evolved only modestly beyond the theoretical MS.

Also not harbouring a Be-phenomenon star is a theoretical-MS or near-theoretical-MS member of a binary system with mass transfer, in which the observed hydrogen Balmer emission comes from an incandescent mass-transfer stream. In the table, this is perhaps the case for β Lyr Aa1 (Sheliak), which certainly has such a mass-transfer stream. Confusingly, however, a “shell” spectrum is observed for Sheliak, and “shell” in the case of a young B star (as we explain below) is generally, or even inevitably, associated with the Be phenomenon. Perhaps all that can be said here is that Sheliak is a confusing case. (It has certainly been notorious over the decades, in one way or another, as a challenge to modelling.) The conceptual point remains that if, hypothetically speaking, emission in a young B star were to come from no source other than a mass-transfer stream, thanks to that star’s membership in a tight binary system, then that star, while being obliged to show the observation-driven “e” flag in its MK type, would not count as an instance of the Be phenomenon

This, then, concludes the qualifying comments. To recapitulate: the true Be-phenomenon stars are theoretical-MS or near-theoretical-MS stars with presently observed or historically observed emission lines, where the emission is not due to a mere mass-transfer process attributable to membership in a tight binary system.

The astrophysical task is now to determine what produces the emission. Emission must mean that the star has somehow managed to shed significant quantities of incandescent gas. Copious shedding cannot be attributed to stellar winds, since winds play only a minor role in mass-shedding for stars within or near the MS (except, perhaps, for the case of stars at the hottest end of the O range,

where even the concept of time-spent-on-theoretical-MS is, as noted above, problematic). Our own Sun, for instance, as an MS star, sheds a mere tenth-of-a-trillionth of its mass per year.

The cause of the copious shedding has not yet been determined with confidence. It is possible that all Be stars are rapid rotators (although, as we remarked in Subsection 4.4, spectroscopy, with its incorporation of “n” or “nn,” as occasionally appropriate, in an MK type, cannot by itself detect rotation when the star is oriented pole-on to Earth). On the other hand, there are many rapidly rotating theoretical-MS or near-theoretical-MS O, B, and A stars that do not present the Be phenomenon.

The following picture therefore suggests itself: if the star is a rapid rotator, and in addition possesses some mechanism “X” for launching photosphere gas from near its equator into its equatorial plane, then an incandescent disk forms, girdling the star, and registering as emission at the spectrograph. With the star a rapid rotator, it will not be a sphere but a rotationally somewhat flattened object, with local gravity in the photosphere somewhat lower at the equator than at the poles, and with launching into an equator-plane orbit consequently favoured. The observed hydrogen Balmer emission is on this picture a signature of hydrogen ionization in the disk, under a violent barrage of UV from the (hot, as O-or-B-or-A) photosphere: Balmer-lines hydrogen light is emitted as part of the process in which free electrons and hydrogen nuclei recombine, where a captured electron falls to the penultimate energy level from some higher level.

The equatorial-disk picture was first proposed in 1931. Now quite widely accepted is a “Viscous Decretion Disk” elaboration of this idea, introduced in [1991MNRAS.250.432L](#). “Decretion” proves a useful contrived astronomical term, created as an antonym for “accretion.” Accretion disks figure in various astrophysics contexts, for instance in such black-hole binaries as Cyg X-1 (material shed by the readily amateur-visible member of this binary falls first onto an accretion disk around the black-hole event horizon), and again in the case of starbirth, where material from the gestating ISM cloud forms an accretion disk around the protostar, in a process that might see the disk eventually transform itself into a bevy of exoplanets, with perhaps also a belt of small rocky asteroid-like bodies, and with some analogue of our Solar System’s zodiacal dust, all orbiting an infant star. Correspondingly, a “decretion disk” forms when an astronomical object (in our case the Be-phenomenon star) for one reason or another releases matter into orbit in its neighbourhood.

Although the dimensions of the hypothesized disk are not easily investigated, emission in the Balmer hydrogen- α line in the cases so far studied has been found to come from a disk on the order of 0.3 AU to 0.6 AU in radius. We seem to have here, in other words, one of the grandest of all theoretical-MS or near-theoretical-MS stellar spectacles.

Unfortunately, it is a spectacle that at best can be imaged only fuzzily, even with the most capable current optical interferometers. Let the Jupiter disk, of diameter $\sim 50''$, familiar from the small telescope, become a circular tea-tray 50 cm in diameter. The binaries resolvable in good seeing by the small telescope, at a separation of $\sim 1''$, thereby become a pair of points on that tray lying 1 cm apart. The most celebrated of the Be-phenomenon stars, γ Cas A, already noted as spectroscopically peculiar by the first stellar spectroscopist, Fr Angelo Secchi, in or shortly before 1866, lies at a distance of 600 ly from Earth. A disk of incandescent gas on the order of 0.5 AU in radius, or 1 AU in diameter, is seen at this distance as an object a mere 5 mas across. In terms of the tea-tray, this corresponds to an object around 50 microns wide, in other words to an object having the approximate width of a human hair. Consistent with the picture of gases launched by “Mechanism X” into circumstellar orbit is the [2007A&A...464...59M](#) discovery that the gas in Be-phenomenon star α Ara A is in a normal central-gravitational-field (i.e. “Keplerian”) orbit, moving unconstrained by any such nongravitational forces as magnetism, and not possessing the kinetics of a mere stellar wind.

What, then, can “Mechanism X” be? It is possible that different Be-phenomenon stars have different gas-launching mechanisms. Outflows from the poles are not currently considered relevant. Nonradial pulsation, on the other hand, may play a role in at least some cases, as may also local magnetic phenomena at the low latitudes. (There is perhaps no known case of a Be-phenomenon star with a strong global magnetic field.) Helpfully, all hitherto scrutinized Be-phenomenon stars have been found to be pulsating variables, although in some cases the pulsation-produced luminosity variation is at the millimagnitude level or below, eluding detection by ground-based photometry. (In addition to facing possible very low-amplitude variations, photometric monitoring of the stellar pulsation is confronted by the complication that the disk itself may vary photometrically (possibly with high amplitude).)

Nonradial pulsation aside, it is possible that in some cases, where the Be star is a member of a binary with tight orbit, or at any rate with an orbit possessing a tight periastron, the “X” role is played by the perturbing gravitational field of the companion.

Some Be-phenomenon stars have emission (from, on the currently accepted modelling, equatorial disks) which is, so far as the existing multidecade observational record goes, stable. Other Be-phenomenon stars, however, present emission lines only intermittently, in their years or decades of “outburst.” Two prominent instances of outburst-and-quiescence in our 322-star set are the already-cited γ Cas A and the recently active δ Sco A (Dschubba). Another well-known instance, although a little too dim for inclusion in the 322-star set, and sharing the notoriety of bright γ Cas A, is Pleione. This star, easy in binoculars as the northern neighbour of Atlas at the eastern extremity of the Pleiades, presented an emission-line outburst of uncertain commencement extending to 1903, and presented additional emission-line outbursts in the periods 1955–1972 and 1989–2005.

Where the disk is permanent, the “X” mechanism works steadily to launch fresh consignments of photospheric gas into orbit, i.e. to perpetuate the decretion. The ongoing launch compensates for the ongoing accretion of matter from at least the inner part of the disk back onto the photosphere. If the mechanism should for some reason cease to operate, decretion ceases, and yet accretion continues. This has the consequence that the disk vanishes (with, however, some of the outlying parts of the disk lost not to accretion onto the photosphere but to outflow, into the embedding ISM).

On some current modelling, a typical Be disk increases in thickness rather gently as one progresses outward (with radially directed tangents to the disk, as taken at the points where disk meets photosphere, yielding a tight “full-opening angle” of $\sim 10^\circ$). A further geometrical detail from some current modelling may also be noted: if, as is often the case, the Be-phenomenon star is a member of a binary not tight enough to produce mass transfer, and yet tight enough to produce a gravitational perturbation from the companion star, and if the Be-phenomenon star equatorial plane diverges somewhat from the orbital plane of the binary system, then the disk is warped.

We may now turn from Be to the related “shell spectrum” phenomenon. The term is somewhat unfortunate, being perhaps a relic from discussions in the early 20th century, when it was perhaps thought that an O or B or A star in or near luminosity class V could under the right circumstances surround itself not with an equatorial disk of gaseous ejecta (as on the currently accepted modelling) but with a literal “shell” of gaseous ejecta, in other words with an enclosing blanket. For better or worse, the term has stuck, surviving the acceptance of the disk morphology (and has nothing to do with thermonuclear-fusion shells in stellar interiors, as discussed in subsections 4.6, 4.7, and 4.8 of this essay). A “shell” spectrum in a rapid rotator, oriented equator-on to Earth, occurs when some lines are seen not in the expected broadened absorption typical of an equator-on rapid photosphere, but in, or also in, narrow absorption. Typically, though not inevitably, the unexpected narrow absorption lines occur as narrow absorption cores within Balmer emission.

On the current understanding, “shell” in this sense typically results when a Be star not only generates its (perhaps temporary) disk of equatorial ejecta, but happens to be oriented more or less equator-on in relation to the spectrograph. Under these circumstances, part of the disk lies between photosphere and spectrograph, yielding the absorption. Since this part of the disk is moving more or less orthogonally to the line of sight, i.e. is neither approaching the spectrograph nor receding, its absorption lines escape the rotational broadening characteristic of absorption lines from the photosphere.

Although a Be-phenomenon star with equator-on orientation can, as just noted, be simultaneously in emission-line outburst and in “shell,” it sometimes happens that shell absorption is present in a Be-phenomenon star even after its emission has for the time being subsided. The Be-phenomenon star Pleione, in particular, had a shell spectrum without emission in the period 1938–1954, and then again for some years after 1973.

Rapid rotators fitting the definition of “shell spectrum” occur even somewhat outside our present domain of interest, the Be-phenomenon stars, with instances known even in type F, right down to the F5 “rotation break.” It remains the case, however, that “shell” is most prominently connected with the Be, as a phenomenon contemporaneous with a Be outburst or present in a star that at some earlier or later time is observed to be in Be outburst.

What, in this general Be-cum-“shell” field, are the possible lines of activity for the amateur spectroscopist?

On the humblest level (even with a visual spectroscope and no camera, as in the case of 1860s Fr Angelo Secchi) it is possible to monitor theoretical-MS or near-theoretical-MS rapid rotators, to see whether emission is currently present or currently absent. The sudden onset of emission would be newsworthy of communication to AAVSO, to the LESIA laboratory at Paris-Meudon (as mentioned again below), or to other appropriate pro-am authorities.

On a less humble level, where spectrograms are taken, and are converted into intensity-against-wavelength plots, or “extracted one-dimensional spectra,” with such professional astrophysical tools as IRAF, the evolution of emission-line and shell-absorption-line profiles could be tracked. In particular, where shell absorption is present simultaneously with emission, as in the (conveniently strong) hydrogen Balmer lines, duly equipped amateurs could examine from month to month whether emission is currently stronger on the violet, or on the contrary on the red, side of the partitioning absorption.

Finally, we suggest in a speculative spirit that it might prove possible to keep a month-upon-month polarimetry log (although we do not ourselves know whether any amateurs in any country have attempted polarimetry, whether in a Be-phenomenon context or in other contexts): if the Be-phenomenon star is not seen pole-on, then some light from its photosphere will be scattered toward the polarimeter by free electrons in the disk and will therefore be linearly polarized.

The recent literature includes a long review article, [2013A&ARv..21...69R](#), on the Be phenomenon. The IAU Working Group on Active B Stars (a group whose domain of interest includes, and yet is not confined to, the Be and shell phenomena) has a homepage at <http://activebestars.iag.usp.br/bstars/>, with a link to its newsletter materials, including a newsletter archive. The LESIA laboratory at the Observatoire de Paris-Meudon maintains the “BeSS Database” comprising Be-phenomenon stars, the Herbig Ae/Be “stars” briefly mentioned near the beginning of this subsection, and a “B[e]” category of supergiants, at <http://basebe.obspm.fr/basebe/>.

APPENDIX: Glossary of acronyms and similar designation

The following is a glossary of the acronyms and similar designations used in our essay and table. We omit, as sufficiently obvious, a small handful of universally known acronyms (e.g. NASA), designations of chemical elements and chemical compounds (e.g. CO, for carbon monoxide), and designations of particular satellites or similar space missions (e.g. *Gaia*, *HIPPARCOS*, *MOST*, *ROSAT*).

- AAVSO: American Association of Variable Star Observers
- AAVSO(VSX): AAVSO International Variable Star Index (<http://www.aavso.org/vsx/>)
- ALMA: internationally funded Chile-based radio interferometer (“Atacama Large Millimetre/submillimetre Array”)
- AMBER: spectro-interferometer at VLT (“Astronomical Multi-BEam combineR”)
- AGB: asymptotic giant branch (as a region in the two-dimensional MK luminosity-versus-temperature stellar classification space)
- Astron. Alm.: *The Astronomical Almanac*, as the joint annual publication, in print and to a reduced extent online, of the United States Naval Observatory and HM Nautical Almanac Office; “Section H” (not necessarily always up to date in the online version) provides V magnitudes, B–V and V–I colours, and MK types for several hundred bright stars; Astron. Alm. particulars can be had from <http://asa.hmnao.com/> and https://en.wikipedia.org/wiki/Astronomical_Almanac
- AU: astronomical unit (the formal 1976 IAU definition is in effect a precisification of the earlier epoch-of-Kepler AU)

concept, under which the AU was the average of the Earth-Sun distances at aphelion and at perihelion)

- BSC5: Yale Bright Star Catalog, Version 5
- BSG: blue supergiant
- CADARS: Catalogue of Absolute Diameters and Apparent Radii of Stars (<https://doi.org/10.1051/0004-6361:20000451>)
- CHARA: the Mount Wilson optical interferometer (Center for High Angular Resolution Astronomy)
- CME: coronal mass ejection
- CNO cycle: the carbon-nitrogen-oxygen-catalyzed cycle under which the hotter stars fuse hydrogen into helium
- COAST: the Cambridge optical interferometer (Cambridge Optical Aperture Synthesis Telescope)
- CODEX: a series of computer codes for the numerical simulation of stellar atmospheres (Cool Opacity-sampling Dynamic EXtended)
- DR2: Data Release 2 (at *Gaia*)
- FDU: First Dredge-Up (as a stage in stellar evolution, soon after a star evolves out of the MS)
- FUV: far ultraviolet
- GCVS: General Catalogue of Variable Stars (Sternberg Astronomical Institute, Moscow)
- GTR: general theory of relativity
- Hp: a visible-light passband used for photometry at *HIPPARCOS*
- HM Nautical: “Her/His Majesty’s Nautical” (for UK publications and UK agencies)
- HR diagram, HR plot: two-dimensional luminosity-versus-temperature plot for the members of some given population of stars; it is useful to distinguish the “observational” (phenomenological, MK-classification) and the “theoretical” HR diagrams
- IAU: International Astronomical Union (Paris)
- IR: infrared
- IRAF: Image Reduction and Analysis Facility: a suite of software tools, for astronomical tasks including aperture photometry and the “extraction of one-dimensional spectra” from raw spectrograms, available free of charge from the National Optical Astronomy Observatory (USA); very widely used at North American professional observatories, and quite widely also, but in competition with MIDAS, at professional observatories outside North America: <http://ast.noao.edu/data/software>
- IS: Instability Strip (as a region in the two-dimensional luminosity-versus-temperature stellar-classification space)
- ISM: interstellar medium
- LESIA: Laboratoire d’Études Spatiales et d’Instrumentation en Astrophysique (physically at Paris-Meudon): <https://lesia.obspm.fr/>
- LPV: long-period variable
- LSR: Local Standard of Rest (as reference frame for kinematics of bodies in our own galaxy)
- M_{\odot} : solar mass
- mas: milliarcsecond
- MK: Morgan-Keenan (two-dimensional phenomenological, non-theoretical, stellar classification scheme, with “MK luminosity classes” and “MK temperature types”)
- MS: main sequence (as a region in the two-dimensional luminosity-versus-temperature stellar classification space; it is useful to distinguish the “observational MS,” in other words the empirical MK luminosity class V, from the “theoretical MS”)
- My: megayears
- NCP: North Celestial Pole
- NPOI: a US Naval Observatory facility (Navy Precision Optical Interferometer)
- NSV: New Catalogue of Suspected Variable Stars (Sternberg Astronomical Institute, Moscow)
- OBAFGKMLTY: the temperature-ordered sequence of MK types, with O the hottest and Y the coolest; until the discovery of brown dwarfs, in types L, T, and (very recently) Y, the sequence was simply OBAFGKM, recalled by 20th-century students with the unfortunate mnemonic “Oh Be A Fine Girl Kiss Me” (implementing gender-neutrality, and allowing for the three progressively cooler brown-dwarf types, one might instead propose “Oh Be A Fine Gymnast, Kiss Me Like This, Yowee”); outside this sequence are the special MK labels (marking gross chemical anomalies) W (for the Wolf-Rayet

stars; these turn out to be hot, like O stars), C (for stars whose photospheres are rich in carbon; these turn out to be cool, like K or M) and S (for stars with chemically anomalous photospheres, these are in terms of spectral phenomenology intermediate between M and C, and turn out to be cool); C is the current label for a group that was in earlier decades divided into R and N: additionally, the special “D” and “P” flags are used, in a more colloquial MK spirit, for planetary nebulae hosts and white dwarf “stars”

- PA: position angle
- PTI: Palomar Testbed Interferometer
- R_{\odot} : solar radius
- R^* : stellar radius (with reference to some given, reasonably spherical, star)
- R_{eq} : equatorial radius (with reference to some given rotationally flattened star)
- RGB: red-giant branch (as a region in the two-dimensional luminosity-versus-temperature stellar classification space)
- R_{pol} : polar radius (with reference to some given rotationally flattened star)
- RSG: red supergiant
- SAAO: South African Astronomical Observatory
- SB: spectral binary
- SETI: search for extraterrestrial intelligence
- SGB: sub-giant branch (as a region in the two-dimensional luminosity-versus-temperature stellar classification space)
- SMEI: Solar Mass Ejection Imager, as an instrument on the *CORIOLIS* satellite
- SN: supernova
- SNR: supernova remnant
- SWB: stellar-wind bubble
- UV: ultraviolet
- V: the visible-light passband in the UBVRI photometric passband system that best approximates the response of the human eye, as lying between the blue (“B”) and red (“R”) visible-light passbands
- VLT: a Chile-based facility of the European Southern Observatory (Very Large Telescope)
- VLTI: the interferometer at VLT
- VSX: AAVSO International Variable Star Index (www.aavso.org/vsx)
- WFC3: Wide Field Camera 3 (as an instrument on the *Hubble Space Telescope*)
- WFPC2: Wide Field and Planetary Camera 2 (as an instrument on the *Hubble Space Telescope*)
- WD: white dwarf
- WR: Wolf-Rayet (as a type of star)
- WDS: Washington Double Star Catalog: www.usno.navy.mil/USNO/astrometry/optical-IR-prod/wds/WDS
- ZAMS: zero-age main sequence (the subregion of the MS comprising stars that have just begun stable core-hydrogen fusion)

History of recent revisions to both essay and table

Recent revisions are tracked with UTC YYYYMMDDThhmmssZ timestamping, in the “major.minor.patch” version-numbering scheme common in software development.

- [20210914T035200Z/5.2.1](#): bolded “Rigel Kentaurus”
- [20210807T203107Z/5.2.0](#): Made various copy-edit corrections (such as insertion of missing punctuation, correction of a few spelling errors), and additionally on the side of scientific substance made a few corrections or amplifications (chiefly as follows: amplified the essay elucidation of “n”, “nn” in MK types; improved an essay remark on rotation in stellar evolution; made essay correction regarding protracted-versus-brief membership of SGB; corrected essay list of Be-phenomenon stars (the phenomenon is not observed in Adhara); added “SGB” to glossary of acronyms; improved table discussion of exoplanet status for α Tau A (Aldebaran); corrected table magnitude range for α Ori Aa (Betelgeuse); corrected table typo for angular distance in α Cru AB (Acrux and companion; correct value is 3.5”, not 35”); corrected table typo for magnitude of η Oph B (correct value is 3.3, not 7.3); updated α PsA Aa (Fomalhaut) table entry to reflect the fact that *HST*-imaged “exoplanet” Dagon (2008) has now faded below the imaging threshold, and is therefore now

believed to be an expanding, and therefore an increasingly tenuous, debris cloud rather than a true exoplanet); this version is a supplement to the 2021 Handbook, with the upcoming 6.x.x series intended to support instead the 2022 Handbook

- [20210217T042710Z/5.1.1](#): Made minor adjustments to tabs and spacing for paragraphs before creating online PDF.
- [20210216T161213Z/5.1.0](#): Made minor adjustments (small points of syntax, spelling, punctuation, or similar, with much bibcode error correction). Added a long paragraph with five methods for retrieving a full-text, all-illustrations PDF from a typical astronomical bibcode citation. Corrected a mistake of astrophysical substance, in the subsection 4.8 discussion of onset-of-helium-core-fusion (violence in the onset of core helium fusion is characteristic of the less massive, not of the more massive, incipient fusers-of-core-helium) This yielded a work sufficiently updated to support uploading to the online-version server.
- [20210128T145046Z/5.0.0](#): Made major revisions of the 4.0.0 version series, by adding several thousand words to the introductory online essay, with stellar-evolution background and a detailed briefing on the amateur-relevant “Be phenomenon” and “shell spectra” (and to a lesser extent by expanding “Remarks,” most notably for α Eri (Achernar), ζ Tau (Tianguan), and α Aql A (Altair); other work on “Remarks” included routine updates for such things as binary position angles and celestial-sphere distances, and also comparison of our MK types against MK types as assigned by Astron. Alm. for epoch 2021.5, with the MK discrepancies logged). The work was not yet sufficiently polished to support uploading to the online-version server.
- [20200815T190800Z/4.0.0](#): Performed sufficient updating of the 3.x.x version series to support the print edition of the 2021 Handbook, but without sufficient updating to support uploading to the online-version server.
- [20191231T235959Z~/3.x.x series](#): Supplemented previous editions of this online publication in various ways, most notably by adding the (rather prolix) results of (rather detailed) primary-literature inspections for α Cet Aa (Mira), α Umi Aa (Polaris), β Per Aa1 (Algol), α Tau A (Aldebaran), ε Aur A (Almaaz), α Ori Aa (Betelgeuse), γ Vel Aa, α Leo A (Regulus), α Vir Aa (Spica), ζ Oph, and α Lyr A (Vega).
- [20181231T235959Z~/2.x.x series](#): Supplemented the 1.x.x version series with some (rather detailed) primary-literature inspections for selected familiar bright stars, thereby expanding “Remarks.”

Star Name	RA (2021.5) Dec h m ° '	m_V	$B-V$	MK Type	π mas	M_V	D ly	μ "/y	PA °	V_{rad} km/s	Remarks
Sun		-26.75	0.63	G2 V		4.8	8 lm				
α And Aa	0 09.5 +29 13	2.07	-0.04	B9p IV: (HgMn)	34	-0.3	97	0.214	140	-12 SB	Aa,Ab < 0.001" Alpheratz
β Cas A	0 10.3 +59 16	2.28v	0.38	F2 III	60	1.2	55	0.554	109	+12	var.: 2.25–2.29 in V, 0.1010 d Caph second-brightest of the δ Set variables (the brightest is α Aql A (Altair)) ¶ rapid rotator: 2011ApJ...732...68C finds the rotation to be > 90% of breakup rate, and radius at poles to be ~24% less than radius at equator, with β Cas A of mass ~2 M_\odot , seen nearly pole-on; β Cas A is notable for being cooler than typical rapid rotators, lying just barely on the rapid side of the F5 “rotation break,” and additionally is notable for being old enough to have evolved off the MS, having in its MS career been an A star rather than an F star (in generally, rotation slows as an aging star increases in radius: but our table of bright stars does harbour at least one other such evolved rapid rotator in type F, namely θ Sco A); an envelope at this modest photospheric temperature is dominated by convection not only at the equator but even at the (~1000 K hotter) poles; consistently with this picture of an envelope everywhere convective, interferometry of β Cas A is found to yield results for low-latitudes gravity darkening inconsistent with 1920s von Zeipel law (the law is accurate only if an envelope is radiative); 2011ApJ...732...68C suggests that in its process of evolution off the MS (in which a core contracts, an envelope expands) β Cas A has been efficient in transferring angular momentum from core to envelope ¶ 2011ApJ...732...68C Fig. 4 presents imaging of β Cas A, from CHARA interferometry var. in β Cep class: 2.82–2.86 in V, 0.1518 d Algenib ¶ $E(B-V) = +0.01$ possible exoplanet ¶ high space velocity (interloper from remoter galactic region?)
γ Peg A	0 14.3 +15 18	2.83v	-0.19	B2 IV	8	-2.6	400	0.009	168	+4 SB	
β Hyi†	0 26.9 -77 08	2.82	0.62	G1 IV	134.1	3.5	24.3	2.243†	82	+23†	
α Phe	0 27.3 -42 11	2.40	1.08	K0 IIIb	38.5	0.3	~85	0.426	147	+75 SB	
δ And Aa	0 40.5 +30 59	3.27	1.27	K3 III	~30.9	0.7	106	0.142	126	-7 SB	Aa,Ab 0.40" Ankaa

α	Cas A	0 41.7	+56 39	2.24	1.17	K0 IIIa	~14.3	-2.0	230	0.060	122	-4 V?	<p>¶ possible debris disk AAVSO(VSX) millimag. variable (from starspots) Schedar ¶ limb darkening observed interferometrically (disk diameter 5.25 mas)</p>
β	Cet	0 44.7	-17 52	2.04	1.02	K0 III [†]	~33.9	-0.3	96	0.235	82	+13 V?	<p>anomalous in being X-ray-bright and yet a slow rotator ¶ evolutionary status uncertain (helium core ignited already, or still contracting?) ¶ Astron. Alm. (epoch 2021.5) assigns MK type G9 III CH-1 CN 0.5 Ca 1</p>
η	Cas A [†]	0 50.4	+57 56	3.46	0.59	G0 V [†]	168	4.6	19.4	1.222	117	+9 SB	<p>B:7.51, K4 Ve, 13.4", PA:62°→326°, 1779→2019 Achird orbit 480 y ¶ Astron. Alm. (epoch 2021.5) assigns MK type F9 V var.:1.6-3.0 (V); B:10.9, 2.1", PA:255°→259°, 1888→2002 orbit >1500 y</p>
γ	Cas A [†]	0 58.0	+60 50	2.15v [†] -0.05		B0 IVnpe (shell) [†]	5	-4.2	600	0.026	98	-7 SB	<p>¶ first "Be phenomenon" discovery (Secchi, 1866); additionally the prototype for the γ Cas type of eruptive irregular variables; background on Be phenomena and γ Cas-type variability is given in www.aavso.org/vsots_gammacas; 2002ASPC..279..22IH summarizes the observational history, including major shell-spectrum phases in 1935-1936 and 1939-1940; despite its historical importance, however, Cas A cannot safely be taken as a typical "Be phenomenon" star, since it presents the peculiarity of hard thermal X-ray emission (cf 2013A&ARv..21...69R p. 42, and also e.g. 2012A&A..537A..59N), derived from magnetic heating (perhaps from magnetic star-disk interaction, perhaps from disk intrinsic magnetic field); rotationally flattened (period = 1.21 d, axial tilt=45°); one of only three Be-phenomenon stars so far observed (via polarimetry, not via interferometry) to produce ejecta disks with differing position angles at different outbursts (2013A&ARv..21...69R p. 42; the other two known instances of this geometrical variation are Pleione and 59 Cyg, both too faint to be in this Handbook table of bright stars) ¶ as of at least 2007, AAVSO has called for amateur assistance with photometry: γ Cas A has been as bright as V mag. 1.6, as faint as V mag. 3; AAVSO reports visual 2021 Jan. 14 as mag. 2.1 or 2.2 ¶ dimming through ISM dust, ~0.35 mag. AB similar, 0.7", PA: 26°→76°, 1891→2018 orbit 168 y, highly eccentric; masses nearly equal; A is mag. 4.1 and B is mag. 4.2</p>
β	Phe AB [†]	1 07.0	-46 36	3.32	0.88	G8 III + G8 III	16	0.3:~180		0.088	293	-1	
η	Cet A+2P	1 09.7	-10 04	3.46	1.16	K1.5 III CN1 [†]	26.3	0.6	124	0.257	123	+12V	<p>¶ Astron. Alm. (epoch 2021.5) assigns MK type K2- III CN 0.5</p>
β	And A	1 10.9	+35 44	2.07	1.58	M0 IIIa [†]	17	-1.8	200	0.209	123	+3 V	<p>slight var.? (AAVSO(VSX): 2.01-2.10 in V) Mirach ¶ Astron. Alm. (epoch 2021.5) assigns MK type M0⁺ IIIa</p>
δ	Cas A	1 27.2	+60 21	2.66v	0.16	A5 IV	32.8	0.2	99	0.301	99	+7 SB	<p>var.: 2.68-2.76 in V, 759 d Ruchbah ¶ E(B-V) =+0.27</p>
γ	Phe	1 29.3	-43 13	3.41v	1.54	K7 IIIa [†]	14	-0.9	230	0.209	185	+26 SB	<p>SB period 193.85 d; also var.: 3.39-3.49 in V, 194.1 d ¶ Astron. Alm. (epoch 2021.5) assigns MK type M0⁻ IIIa</p>
α	Eri	1 38.5	-57 08	0.45v [†] -0.16		B3 Vnpe (shell?) [†]	23	-2.7	140	0.095	114	+16 V	<p>ecl. var.: 0.40-0.46 in Hp, 1.263 d Achernar variable in λ Eri class (pulsation? or, rather, starspots?) ¶ brightest of the "Be phenomenon" stars (but first recognized as such only recently, ~1976); an active Be phase that began between 2012 Dec. and 2013 early Jan. (after a period of inactivity, with α Eri presumably diskless, over the previous 7 years) was first noted in amateur spectroscopy, in Brazil; Balmer hydrogen-α line indicates a slow, steady buildup of the Be disk, over a period of ~1.6 y, with polarization suggesting that disk was slightly less dense in 2014 than it had been in 2013; 2017A&A...601A..118D, a case study of α Eri, presents for the first time in astrophysics images of a disk forming around a Be-phenomenon star (with H-band (IR) emission from the disk extending to an outer radius of between 1.7 and 2.3 stellar equatorial radii, in good agreement with current computations in the general theory of the Be phenomenon); it is possible that plane of α Eri disk is inclined to plane of stellar equator; rapid variations in polarization indicate that in addition to its disk, α Eri possesses rings, due to episodic ejections of gas consignments from its photosphere ¶ α Eri is a notably rapid rotator (< 2.1 d) within the (currently small) population of stars interferometrically</p>

													resolved; period in or near the disk phases varies, either because gas is injected (“decreted”) from photosphere into Be disk or because Be-disk gas is re-accreted onto photosphere; interferometry as performed when α Eri is temporarily without its Be disk reveals oblateness (cf e.g. www.eso.org/public/unitedkingdom/news/eso0316/) ¶ although 2008A&A...484L..13K reports companion, at 2007 Dec. angular distance < 0.15”, WDS has not, as of 2020 Nov., asserted binarity; the orbital motion of this companion seems to not be correlated with the repeated formation and disappearance of the Be disk
τ	Cet A+4P	1 45.1	-15 50	3.49	0.73	G8 V [†]	~274.0	5.7	11.9	1.921 [†]	296	-16 [†] V	mass < 1 M _⊙ (unusual in Sample S, although typical in Population P) ¶ high space velocity, low metallicity: interloper from thick galactic disk ¶ on original Frank Drake (1960) SETI target list [THIS STAR ONLY IN ONLINE VERSION OF TABLE]
α	Tri A	1 54.3	+29 41	3.42	0.49	F6 IV	52	2.0	63	0.234	177	-13 SB	Moallahah
β	Ari A [†]	1 55.8	+20 55	2.64	0.16	A4 V	56	1.4	59	0.148	138	-2 SB [†]	Sheratan β Ari B (mag. 5.2) is SB companion of β Ari A exceptionally elongated orbit (0.08 AU min, 1.2 AU max, 107 d); the SB companion has been resolved interferometrically; one of only a few tens of binaries in which orbit is ascertainable both with spectroscopy and with micrometer astrometry (this duplication facilitates model testing)
ϵ	Cas	1 56.0	+63 46	3.35	-0.15	B3 IV:pe (shell?) [†]	8	-2.2	400	0.037	121	-8 V	instance of “Be phenomenon” ¶ He-weak (cp α And, α Tel)
α	Hyi	1 59.4	-61 28	2.86	0.29	F0n III-IV [†]	45	1.1	72	0.265	84	+1 V	rapid rotator (< 30 h) ¶ metal-rich
γ	And A	2 05.2	+42 26	2.10	1.37	K3 IIb [†]	9	-3.1	400	~0.065	~139	-12 SB	B: 5.0, B9 V, 12.0” (2019); C: 6.5, A0 V; BC 0.2” BC orbit: 63.7 y ¶ limb darkening observed interferometrically (disk 6.80 mas) ¶ Astron. Alm. (epoch 2021.5) assigns MK type K3- IIb calcium weak? Almach
α	Ari +1P	2 08.4	+23 34	2.01	1.15	K2 IIIab	~49.6	0.5	66	0.240	128	-14 SB	Hamal
β	Tri	2 10.8	+35 05	3.00	0.14	A5 IV	26	0.1	130	0.154	105	+10 SB2 [†]	SB orbit rather elongated (0.17 AU min, 0.42 AU max) ¶ IR excess (circumstellar matter? possible harbinger of planetesimals)
o	Cet Aa [†]	2 20.4	-2 53	6.47v [†]	0.97	M5-10 IIIe [†]	11 [†]	1.7	300 [†]	0.238	178	+64 V	LPV, 2-10.1; Ab (VZ Cet)WD, 10.4, 0.5”, ~500-600 y Mira [†] ¶ recent maxima Nov. 2019 (V~2.3), Sept.-Oct. 2020 (V~3.0), AAVSO visually mag. ~7.0 or ~6.5 2021 Jan. 12: 2009ApJ...691.1470T discusses variability, including variation in dominant (333 d) pulsation period and the question of longer-period variations ¶ times of maxima are, and times of minima are not, independent of wavelength: minima are at least coarsely correlated with maximum diameter of o Cet Aa ¶ 2016A&A...586A..69P discusses discrepancies in distance determinations (350 ly, 380 ly, 340 ly, and (least reliable?) <i>HIPPARCOS</i> 300 ly) ¶ prototype of the AGB variables, mass ~1 M _⊙ : the first O-rich AGB star with a CI detection (2018A&A...612L..11S) ¶ physical radius ~2 AU in visual, ~4 AU in IR, still greater upon taking instead the “radio photosphere,” which itself increases in radius as progressively longer radio wavelengths are selected: 2015ApJ...808..36M draws parallels with α Ori Aa, attributing radio inhomogeneities in both cases to convective cells (and cf also 2016A&A...592A..42K , which summarizes some recent radio work) ¶ nearest instance of (weak) symbiotic binarity, and the only symbiotic to be observed in all wavelength regimes from X-ray to (mm, also cm) radio; interferometry (in IR) is available from VLT, and <i>CASSINI</i> has yielded (via Saturn-ring occultations) tomographically recovered imaging (2016MNRAS.457.1410S); <i>GALEX</i> has found bow shock, tail (length 13 ly) in ISM: mass-loss rate ~2.5e-7 M _⊙ /y; asymmetric atmosphere is discussed in 2016MNRAS.457.1410S ¶ 2018A&A...620A..75K reports dust trail linking Aa,Ab (consistently with other reports of Aa-to-Ab mass transfer) ¶ 2016A&A...592A..42K (2017A&A...599A..59K) discusses o Cet Aa dust nucleation generally, with

γ	Cet A [†]	2 44.4	+3 20	3.54	0.09	A2 Va	41	1.5	80	0.207	225	-5 V	<p>reference to aluminum (resp. titanium) species: in o Cet Aa, it is silicates that dominate the spectrum (in contrast with less-evolved stars, in which alumina features are spectrally dominant); <u>2016A&A...590A.127W</u> discusses SiO gas, o Cet Aa inner dust shells: it seems still an open question whether o Cet Aa dust formation is cyclic, as part of the photometrically evident pulsation cycle, or proceeds independently of the pulsations ¶ X-ray emission from o Cet Aa was reported in 2005 (2005ApJ...623L.137K, as the first X-ray detection from an AGB star), and OH, SiO maser emission has also been reported (cf. e.g. 2017MNRAS.468.1703E); further, 2015A&A...577L...4V asserts a hot spot, proposing magnetic activity as the cause ¶ 2016A&A...590A.127W summarizes history of modelling: models generally agree that near o Cet are alternating circumstellar layers of infall and outflow, and that at greater radii is an accelerating outflow, from dust-driven winds: recent observations have tended to agree with overall results from running CODEX (e.g. 2014A&A...565A.119S) ¶ Aa,Ab orbit would, if better known, yield improved total mass of Aa,Ab system, thereby advancing the overall theory of AGB stars ¶ protoplanetary disk was detected around Ab in 2007 ¶ Astron. Alm. (epoch 2021.5) assigns MK type M5.5–9e III ¶ Fabricius noted variability in 1956; Hevelius proposed the name Mira in 1642 ¶ for entry-level briefing-with-bibliography, cf www.aavso.org/vsots_mira2.updpadding www.aavso.org/vsots_mira; and for summary of recent primary literature, cf first section of 2016MNRAS.460.673N B: 6.23, 2.0", PA:283°→299°, 1825→2015 orbit \geq 320 y B: 4.35, A1 Va, 8.2", PA:82°→90°, 1835→2020 Acamar low-amp. Cep., 4.0 d; B: 9.1, F3 V, 18.4" (2016) Polaris the brightest of the Cepheids, but not a classical Cepheid, matching instead the "s-Cepheid" light-curve phenomenology of 1995A&A...303..137B ¶ AAVSO(VSX) as at 2021 Jan. 15 gives V-mag. range 1.97–2.00, period 3.9696 d: period is increasing ~4.4–4.9s/y, with sudden change around 1963, and with <i>CORIOLIS</i> satellite suggesting a further recent change: period change is often in Cepheid theory linked to evolution, but this may not be the whole story here (in particular, pulsation-driven mass loss through stellar wind, as affirmed by some recent authors (denial also published) would increase the rate of period change) ¶ pulsation mode (1st overtone? 2nd? fundamental?), evolutionary history (1st crossing of IS? or 3rd crossing?), and distance are controverted by various 2010-through-2018 authors (we here use <i>Gaia</i> DR2 distance for α UMi B as a proxy, assuming with the current literature that B is indeed gravitationally bound with Aa,Ab) ¶ α UMi Aa is first Cepheid with mass determined through purely dynamical means (via the Aa,Ab orbit: Aa is single-lined SB, and Aa, Ab have been resolved with HST, as first announced (0.17") in 2008AJ...136.1137E (orbit ~30 y)) ¶ α UMi Aa is significant for general astrophysics as a possible anchor point for the Cepheid period-luminosity relation at the heart of extragalactic distance determinations, and is important also as a case study in the "Cepheid mass discrepancy" problem (Cepheid masses deduced from pulsation periods are found to be too low in comparison with masses from stellar-evolution modelling) ¶ strictly a three-star system, UMi Aa+ UMi Ab+UMi B; Aa,Ab has period 29.6 y, separation 6.7 AU min, 27 AU max, 17 AU average; B experiences Aa,Ab as essentially a point mass, with period \geq 42,000 y, separation at least 2400 AU; B is mag. 9.1, at angular distance 18" ¶ α UMi Aa, Ab,B is approaching NCP: closest approach will be 14', in ~2105 ¶ B has E(B–V)=0.0 ¶ 2018ApJ...863..187E summarizes recent work 2.45–2.54 in V Menkar mildly variable, in the "giant irregular" class ¶ radio source (due to stellar wind) ¶ notably deficient in carbon</p>
θ	Eri A	2 59.1	-40 13	3.28	0.17	A5 IV	30	0.5	100	0.057	293	+12 SB2	
α	UMi Aa [†]	2 59.3	+89 21	1.97v [†]	0.64	F5–8 Ib	7.5 [†]	-3.6	430 [†]	-0.046	-105 [†]	-17 SB	
α	Cet	3 03.4	+4 10	2.54v [†]	1.63	M2 III [†]	13	-1.9	250	0.078	188	-26 V	

γ	Per Aa,Ab [†]	3 06.4	+53 35	2.91 [†]	0.72	G8 III [†] + A2 V	13	-1.5	240	0.006	175	+3 SB2 [†]	<p>¶ Astron. Alm. (epoch 2021.5) assigns MK type M1.5 IIIa</p> <p>composite spectrum; orbit 14.6 y, next eclipse 2035 eclipse duration < 2 weeks, with AAVSO(VSX) ephemeris giving 2019 Dec. 25 as midpoint date; eclipse variation significantly above threshold of naked-eye detection, with AAVSO(VSX) giving V-mag. range 2.91–3.21 (and giving period 5346 d (14.64 y))</p> <p>¶ orbit is highly elliptical</p> <p>¶ Aa is mag. 3.6 and Ab is mag. 3.8</p> <p>¶ Astron. Alm. (epoch 2021.5) assigns MK type G5 III</p>
ρ	Per	3 06.6	+38 55	3.32v [†]	1.53	M4 II	11	-1.6	310	0.167	129	+28	semiregular variable: 3.3–4.0 in V, ~50 d period ~50 d, with possibly also a longer period
β	PerAa1 [†]	3 09.6	+41 02	2.09v [†]	0.00	B8 V [†]	36	-0.1	90	0.003	119	+4 SB	<p>Aa=compos. spectrum Aa1,2 ecl.;2.09-3.30 in V, 2.9 d Algol Aa2 is K2IV?</p> <p>¶ in older terminology, β Per Aa1 = β Per A, β Per Aa2 = β Per B, β Per Ab = β Per C; but WDS, following the current terminology, uses the “B” and “C” for other purposes (since there are optical neighbours B,C,D,E,F,G,H; all are between 5” and 100” from the Aa1,Aa2,Ab triple, and all are fainter than mag. 10); system is hierarchical, with outlying Ab experiencing the close (separation 14.14 R_o) Aa1,Aa2 pair as essentially a point mass; angular distance between Aa1,Aa2 and Ab is ~0.1” (WDS 1973, 2010)</p> <p>¶ among the most visually prominent of the eclipsing binaries, and for theoreticians the most familiar of the semidetached binaries (i.e., binaries in which one of the two Roche equipotential surfaces is fully occupied, the other not)</p> <p>¶ Aa2 is tidally locked, in a rapid circular orbit with Aa1; the consequent rapid spin of Aa2 causes dynamo action in Aa2 convection zone, with Aa2 consequently having complex magnetosphere (mass-transfer stream possibly even deflected out of Aa1,Aa2 orbital plane by magnetics; 2012ApJ...760...8R; Aa2 has additionally a meridional coronal loop, approximately as high as the diameter of Aa2 (the size exceeds what has been anticipated from modelling) believed pointing at all times to Aa1), X-ray emission, varying radio morphology (double-lobed when radio-brightest) and CME episodes (2017ApJ...850..191M) suggests the 1997 Aug. 30 superflare event supplies “arguably the best candidate” for a non-solar CME)</p> <p>¶ the (unsteady) Aa2-to-Aa1 mass transfer, while ongoing, and indeed responsible for an annulus around Aa1, is no longer copious (in contrast with the copious transfer still present in, e.g. β Lyr)</p> <p>¶ it is not the (now modest) unsteady mass transfer, but possibly instead the Applegate mechanism (1992ApJ...385..621A), implicating a stellar magnetic activity cycle, which dominates the Aa2,Aa1 period variation (increase-decrease-increase cycle, not quite strictly periodic, 32 y: there are additionally period modulations of 1.9 y and 180 y; as at 2021 Jan. 28, AAVSO(VSX) asserts period 2.86736 d); full amplitude of the Aa1 Aa2 period variation is ~0.8 s; such alternating period changes in binaries are still not, however, well understood</p> <p>¶ it is the (several My ago rapid and copious) mass transfer that resolves the “Algol paradox” of a lower-mass more evolved (in this case, sub-giant) star in orbit with a higher-mass less evolved (indeed MS) star; masses are well known in this particular case: 2015MNRAS.451.4150K, having disentangled the β Per Aa1, Aa1, Ab spectra, determines their masses within plus-minus 2%, corroborating 2012ApJ...752...20B</p>

													¶ β Per Aa2 elemental abundances below corona and flare (investigated in 2015MNRAS.451.4150K) are of special interest, since mass transfer has stripped off Aa2 outer layers, opening the Aa2 interior to spectroscopic inspection ¶ 1983ApJ...273L.85K reports discovery of Chandrasekhar eclipse-induced stellar limb polarization from β Per Aa1, in a wide optical passband ¶ MK type K2 IV is assigned to Aa2 in at least 3 recent papers, whereas the older 1993ApJ...410.808L has the slightly hotter MK type K0 IV; what is essential here is the agreed “IV” (as opposed to “V”), indicating evolution of this (secondary) star off the MS (and Astron. Alm. (epoch 2021.5) assigns MK type B8 V for Aa1, as we do here, and for one companion (is this Aa2, or is it Ab?) the uncertainty-flagged “F;” without luminosity class) ¶ β Per Ab, spectrally Am with some FIV characteristics, orbits the β Per Aa1,Aa2 binary with period ~680 d, without eclipsing ¶ 2012ApJ...752...20B presents CHARA interferometry (~0.5 mas, H (near-IR) band) of the β Per Aa1,Aa2,Ab system (finding orbital plane of Ab to be nearly perpendicular to Aa1,Aa2 orbital plane), and also summarizes earlier interferometry; an approx 55-frame “movie” from this paper can be conveniently viewed at https://en.wikipedia.org/wiki/Algol ¶ $E(B-V)=+0.03$ ¶ 2013ApJ...773...1J suggests that β Per system variability is documented in the “Cairo Calendar” papyrus (New Kingdom, dated to 1271–1163 BCE); al-Sufi (Persia, ca 964 CE) is, however, silent on question of β Per variability ¶ AAVSO has briefing notes, with some history, at www.aavso.org/vsots_betaper ; 1910ApJ...32..185S is the discovery paper for β Per Aa1,Aa2 secondary minimum, from the beginnings of photoelectric-cell photometry; 1998A&AT...15..357P analyzes “Algol paradox” history; https://arxiv.org/pdf/astro-ph/0611855.pdf . “Appendix B.” is a tabular history of β Per-pertinent investigations from antiquity to 1999; in this same K. Wecht 2006 Lehigh Univ PhD thesis, Table 2.5.1 summarizes 1966-through-1983 observational coverage, as tabulated in the less Web-accessible 1986 work of Budding in open cluster Mirfak
α	Per A	3 25.9	+49 56	1.79 [†]	0.48	F5 Ib	~6.4	-4.2	510	0.035	138	-2 V	near edge of HR diagram IS (slightly too hot to be a straightforward Cepheid)
δ	Eri	3 44.3	-9 42	3.52 _v	0.92	K0 IV	111	3.7	29.5	0.749	353	-6	[THIS STAR ONLY IN ONLINE VERSION OF TABLE]
δ	Per Aa	3 44.5	+47 51	3.01	-0.12	B5 IIIIn [†]	6	-3.0	500	0.050	149	+4 SB	variable in γ Cas class: 2.99–3.04 in V ¶ cluster affiliation is controverted ¶ Astron. Alm. (epoch 2021.5) assigns MK type B5 III ¶ $E(B-V)=+0.04$
γ	Hya	3 46.9	-74 10	3.26	1.59	M2 III	15.2	-0.8	~214	0.126	24	+16	evolutionary status is uncertain
η	Tau Aa	3 48.8	+24 10	2.85	-0.09	B7 IIIIne [†]	8	-2.6	400	0.048	156	+10 V?	brightest member of Pleiades Alcyone ¶ rapid rotator, with “Be” and “shell-spectrum” phenomena (BSC5: “rotationally unstable”) ¶ 1972JBAA...82..431K describes the 18.6-year 1940-through-2050 cycle of lunar occultation possibilities ¶ significant dimming by ISM dust; $E(B-V)=+0.03$ B: 9.16, B8 V, 12.9”, PA:205°→209°, 1824→2012 orbit $\geq 50,000$ y ¶ significant dimming by ISM dust; $E(B-V)=+0.33$ (pronounced reddening)
ζ	Per A [†]	3 55.5	+31 57	2.84	0.27	B1 Ib	4	-4.0	800	0.011	150	+20 SB	Ca, Cr weak Zaurak
γ	Eri A	3 59.0	-13 27	2.97	1.59	M1 IIIb [†]	16	-1.0	200	0.129	151	+62	¶ Kaler, at http://stars.astro.illinois.edu/ , writes, “must be one of the least-studied of the cooler bright stars” ¶ Astron. Alm. (epoch 2021.5) assigns MK type M0.5 IIIb Ca–I

ε	Per A [†]	3 59.3	+40 04	2.90v [†] -0.20	B0.5 IV [†]	5	-3.6	600	0.028	149	+1 SB2	B: 7.39, B9.5 V, 9.1", PA:10°→10°, 1821→2015 orbit ≥ 16,000 y ¶ variable possibly in the β Cep class (2.89–2.91 in V; one of the most extreme spectroscopic variables (periods 2.27 h and 8.46 h)) ¶ E(B–V) = +0.10	
λ	Tau	4 01.9	+12 33	3.41v [†] -0.10	B3 V	7	-2.4	480	0.017	209	+18 SB2 [†]	ecl.: 3.37–3.91 in V, 4.0 d; secondary is A4 IV AAVSO(VSX) as at 2021 Jan. 16 gives period 3.9529478 d ¶ shape distortion (mutual tides), reflection effect, some evidence of mass transfer	
α	Ret A	4 14.7	-62 25	3.33	0.92	G8 II–III	20.2	-0.1	162	0.065	40	+36 SB?	in evolutionary terms in “Red Clump,” fusing helium in Hyades; Aa,Ab 0.2", mags. ~3.6, ~6.0
ε	Tau Aa +1P [†]	4 29.9	+19 14	3.53	1.01	K0 III [†]	22.2	0.3	150	0.113	110	+39 V?	Ain ¶ metal-rich ¶ first known instance of a planet-host in an open cluster; unusually massive among the currently known planet-hosts ¶ Astron. Alm. (epoch 2021.5) assigns MK type G9.5 III CN0.5
θ	Tau Aa [†]	4 29.9	+15 55	3.40v [†]	0.18	A7 III	22	0.1	150	0.112	104	+40 SB [†]	[THIS STAR ONLY IN ONLINE VERSION OF TABLE] in Hyades, system Aa-plus-Ab is a.k.a. θ^2 Tau Chamukuy companion in elongated orbit (0.23 AU min, 1.3 AU max); the SB system θ Tau Aa,Ab forms wide double with the bright SB system θ Tau Ba-plus-Bb, a.k.a. θ^1 Tau (mag. 3.35–3.42; separation of Aa,Ab from Ba,Bb is 337") ¶ variable in the δ Sct class; 3.35–3.42 in V; 12 periods known, 1.64 h to 2.22 h, ranges 0.5 millimag. to 30 millimag A: 3.8; B: 4.3, B9 IV; 0.2" (2019); orbit 12 y orbit very elongated: 1.9 AU min, 17.5 AU max ¶ Astron. Alm. (epoch 2021.5) assigns MK temperature type A0p Si (and does not assign an MK luminosity class)
α	Dor A [†]	4 34.5	-55 00	3.30	-0.08	A0p V: (Si) [†]	19	-0.3	169	-0.059	~79?	+26	irregular var., 0.86–0.89 in V Aldebaran ¶ foreground star, not true Hyades member; among the nearest of the red giants; evolution has proceeded beyond the “FDU” stage which accompanies helium-core contraction on RGB ¶ 49 lunar occultations occurred over the period 2015 Jan. 29/2018 Sep. 03 (and yet there is a surprisingly large scatter in the occultation determinations of α Tau angular diameter; 1972JBA...82..431K describes the overall 18.6-year 1940-through-2050 cycle of lunar occultation possibilities) ¶ in contrast with its celestial-sphere neighbour α Ori, α Tau is of modest mass (with recent literature variously offering ~1.2 M_{\odot} , ~1.3 M_{\odot} , ~1.5 M_{\odot}): Appendix C of 2018ApJ...865L..20F tabulates values for mass, luminosity, radius, age, and several other parameters, on the strength of five separate 2008-through-2012 spectroscopy investigations ¶ 2013A&A...553A...30 reports “MOLsphere” (molecule-harboured atmosphere) inhomogeneities, from VLTI/AMBER, thereby helping advance the still poorly understood topic of RGB mass loss (especially in a context in which dust condensation might appear not to play a significant role; in general, it is RGB mass loss that is puzzling, AGB mass loss that is straightforward) ¶ recent literature proposes oscillations, and also proposes rotational modulation from modest photospheric-activity features (with possibly an activity cycle (2015A&A...580A..31H): the features could be (cool) starspots, but could alternatively be large convection cells; the general topic of activity in K giants is not yet well understood) ¶ Astron. Alm. (epoch 2021.5) assigns MK type K5+ III ¶ although 2019A&A...625A..22R casts doubt on 2018ApJ...865L..20F , 2015A&A...580A..31H exoplanet assertion, exoplanet is asserted in NASA exoplanet catalogue (as viewed 2021 Aug. 07)
π^3	Ori A	4 51.0	+7 00	3.19	0.48	F6 V	124	3.7	26.3	0.464	89	+24 SB2	Tabit
ι	Aur	4 58.4	+33 12	2.69v	1.49	K3 II	7	-3.2	500	0.016	155	+18V	Hassaleh var.: 2.63–2.78 in V; possibly 2 exoplanets ¶ X-ray “hybrid star” (unusual combination of (hot) corona, cool wind) ¶ dimming by ISM dust, ~0.6 mag.
ε	Aur A	5 03.5	+43 51	3.03v [†]	0.54	F0lab? [†]	~2 [†]	-8.0:~1450 [†]	~0.003	n.a.	-3 SB [†]	ecl.: 2.92–3.83 in V, ~27.1 y (dim ~700 d) more formally, period is 9896.0 ± 1.6 d: as again discussed	Almaaz

twice below, there are spectroscopic, as distinct from photometric, phenomena indicative of an eclipsing mass before the onset, and continuing after the end, of the photometric eclipse

¶ Astron. Alm. (epoch 2021.5) assigns MK type A9 Ia

¶ ϵ Aur B MK type is ~B5 V

¶ ϵ Aur ranks among the longest-period eclipsing binaries (exceeding even V383 Sco, with period 13.5 y; WW Vul, with period 13.9 y; and VV Cep, with period 20.3 y: the current long-period record is held by TYC 2505-672-1, at ~69.1 y (with dimming ~3.45 y))

¶ determination of orbit elements has proven troublesome, with [2012A&A...544A..91M](#) urging caution even in respect of recent careful studies

¶ it is remarkable that, even though the eclipsing entity is physically very extended (because the eclipse is protracted), and even though orbital dynamics indicates that the entity is quite massive, nevertheless no visible radiation from an eclipsing body is readily observable (i.e. it is remarkable that this SB is essentially a single-lined SB)

¶ although the (notably protracted) ϵ Aur eclipse is largely flat-bottomed, nevertheless even during eclipse the (dimmed) spectrum of the primary can be seen, with no visible-wavelength colour preference in the attenuation (except that there are absorption lines, as from a semi-transparent atmosphere around the eclipsing mass, at the start and the end of the dimming); the [1937ApJ...86..570K](#) explanation, postulating a large semitransparent totally eclipsing mass, with the non-selective opacity due to scattering off free electrons, is now universally abandoned in favour of the [1953AJ....58..219K](#) and [1965ApJ...141..976H](#) hypothesis of an almost edge-on ([2010ApJ...714..549H](#)) cool opaque gas-dust low-mass disk or disk-like entity (spiral arm? cf [2013PASP..125..775G](#)) (rotating while orbiting, and several AU in diameter, presenting a temperature gradient ~550 K to ~1150 K (representing the portions respectively farthest from and closest to the primary star), and in terms of its vertical development not a (thick) hockey puck but a (thin) wafer, of much larger radius than the primary star; [2013PASJ...65L...1S](#) gives evidence for clumping in the disk; [2015ApJS..220...14K](#) raises the possibility that the disk is slightly tilted out of the binary-system orbital plane, with consequent precession), shrouding a B-type star (B5V?) or star pair (the more dramatic hypothesis of a shrouded black hole is not now generally favoured: [2010AJ...140..595W](#), e.g. reports null result from X-ray search), with the disk geometry making the eclipses of the primary star, as observed from Earth, only partial; the disk may have been formed by mass transfer from the primary star, and indeed [2013PASP..125..775G](#) and [2018MNRAS.479.2161G](#) report putative spectroscopic detection of narrow mass-transfer stream; the former paper stresses that the detection of rare-earth elements within the putative stream spectrum (an indication that the primary is highly evolved?) now poses a fresh puzzle, in a system traditionally classed as puzzling

¶ [2012ApJ...748L..28H](#) and [2012MNRAS.423.2075M](#) discuss the question of gas-to-dust ratio in the disk; [2015ApJ...798...11P](#), [2012MNRAS.423.2075M](#), and [2010ApJ...714..549H](#) suggest not-very-small values

in the distribution of dust-particle diameters, with the first two of these three papers suggesting carbonaceous chemistry; additionally, [2011AJ...142..174S](#) spectroscopy finds CO absorption bands, symptomatic of sublimation, with indications that large particles dominate [¶ 2013ARep...57..991P](#) and [2013PASP..125..775G](#) document indications that the structure of the disk does not greatly change from one eclipse to the next [¶](#) the brightening around mid-eclipse has in the post-1970 papers repeatedly been attributed to a central opening in the postulated disk: however, (a) dissenter [2011A&A...530A.146C](#) has instead suggested intrinsic variability in the primary (which indeed has various quasi-periods or periods, with 67 d and 123 d prominent, with also variations in radial velocity, and (unblended) spectral line width, and other periodic or quasi-periodic behaviour, including possible orbitally excited non-radial pulsation; there seems as yet, however, to be no extensive astroseismology), and (b) dissenter [2011A&A...532L..12B](#) has instead suggested forward scattering by disk dust (a line of thought now supported by the key imaging-and-modelling paper [2015ApJ...220...14K](#)) [¶](#) *HIPPARCOS* yields π possibly < 2 mas, distance ~ 2000 ly; we now, however, choose to relinquish the *HIPPARCOS* determination, made at the limit of *HIPPARCOS* capabilities, in favour of [2019BVS.6258...1P](#), which deduces from *Gaia* DR2 $\pi = 2.4144 \pm 0.5119$ mas, and goes on to deduce from this, via supplementary (not straightforward, Bailer-Jones et al. [2018AJ...156..58B](#)) considerations what we express here as “ ~ 1450 ly” [¶](#) section 1 of [2012A&A...546A.123G](#) and section 1 of [2012A&A...544A..91M](#) summarize past controversies regarding mass of primary (low or high?), stemming from the difficulty in determining distance ([2012A&A...546A.123G](#) assigns a high distance, ~ 4900 ly, and consequently favours a high mass value, $\sim 20 M_{\odot}$; however, several post-2009 papers instead assign a modest mass to the primary, suggesting various values within the range $\sim 2 M_{\odot} - \sim 6 M_{\odot}$: [2014MNRAS.445.2884M](#), e.g. suggests $2.5 M_{\odot}$ for primary, $5.4 M_{\odot}$ for secondary (and suggests disk diameter 8.9 AU): evolutionary status of the primary has been correspondingly controverted (post-AGB star, now of modest mass, with much past shedding of mass, and consequent accumulation of the low-mass opaque disk around the secondary (a view taken by various papers, including recently [2019BVS.6258...1P](#)) or, rather, an evolutionally earlier supergiant (cf [2012JAVSO..40..647K](#)), even perhaps of high mass? – but it is clear that the primary is at any rate sufficiently evolved to have left the MS, and there are indications that it is pulsating and a wind source; angular diameter is 2.1 mas) [¶](#) most recent photometric eclipse started 2009 Aug. 12 ended 2011 Aug. 23 ± 15 d; next secondary (shallow, for the casual observer elusive) eclipse is possibly 2025 Dec. 20 through 2028 Mar. 29; next (deep, easy observable) primary photometric eclipse starts in 2036; monitoring even outside both the primary eclipse and the secondary eclipse is useful, in part because of intrinsic variations in the primary star (cf [2012JAVSO..40..647K](#)); in part because the postulated dense disk

													has an extended “atmosphere” yielding (e.g.) H α absorption even outside photometric eclipse (2011A&A...530A.146C), with spectral premonitions starting ~3 y before the onset of the photometric eclipse; and in part because the opaque primary-star-eclipsing disk is potentially liable to thermal changes, visible in mid-infrared outside primary and secondary eclipse (2011AJ...142.174S) ¶ the Kloppenborg et al. CHARA interferometric imaging of the eclipsing disk is perhaps the single largest 21st-century advance in ϵ Aur studies: 2010Natur.464.842G supplies journalistic background, including a recapitulation of 2010ApJ...714..549H modelling; 2010Natur.464.870K is the formal Kloppenborg et al. discovery paper (with the first spatially resolved image for any eclipsing binary during eclipse); and 2015ApJS...220...14K is a Kloppenborg-et-al update, with additional interferometry, now including also PTI and NPOI (and supplying also an overall history of ϵ Aur studies) ¶ news sources include http://mysite.du.edu/~rstencil/epsaur.htm (Prof. R. Stencel, Univ of Denver, on the Kloppenborg-2010 team) and https://twitter.com/epsilon_Aurigae ; 2012JAVSO...40..618S summarizes the 2009-2011 campaign from an AAVSO perspective; an 18-paper archive, of NSF-supported ~2009-through~2011 AAVSO eclipse campaign, is at www.aavso.org/citizen-sky-epsilon-aurigae-papers
ϵ	Lep	5 06.4	-22 21	3.19	1.46	K4 III	15	-0.9	210	0.076	164	+1	
η	Aur	5 08.0	+41 16	3.18v	-0.15	B3 V [†]	13	-1.2	240	0.075	155	+7 V?	Haedus
β	Eri A	5 08.9	-5 04	2.78 [†]	0.16	A3 IVn	36	0.6	89	0.112	228	-9	Cursa
μ	Lep	5 13.9	-16 11	3.29v? [†] -0.11		B9p IV: (HgMn) [†]	18	-0.5	190	0.050	109	+28	
β	Ori A [†]	5 15.6	-8 11	0.18 [†]	-0.03	B8 Ia	4	-6.9	900	0.001	69	+21 SB	Rigel
α	Aur Aa,Ab [†]	5 18.3	+46 01	0.08	0.80	G6:III + G2:III	76	-0.5	43	0.433	170	+30 SB2	Capella
η	Ori Aa [†]	5 25.6	-2 23	3.35v [†] -0.24		B0.5 Ve [†]	3	-4.0	1000	~0.004?	n.a.	+20 SB2	

evolutionary status is uncertain: RGB or AGB?
rotating ellipsoid var?: 3.16-3.18 in V, 2.5617 d
spectral variations also suggested
¶ weak magnetic field detected, ~2 \times strength of
Earth’s dipole field

unexplained brightening episode, over 2 h, by ~3 mag,
in 1985 (recalling the 1972 unexplained
brightening of ϵ Peg)
var?: 2.97-3.41 in V?, 2 d?
variable in α^2 CVn class? (variability so far
unconfirmed, and no CVn-class-appropriate
magnetic field detected yet?)
¶ among the brightest of the Hg-Mn stars
¶ Astron. Alm. (epoch 2021.5) assigns MK temperature type
B9p HgMn (and does not assign an MK luminosity class)
¶ X-ray emission noted from putative companion,
at angular distance 0.93”
B: 6.8, B5 V, 9.7” (2017); C: 7.6; BC: 0.1”
A-BC orbit \geq 25,000 y, BC orbit ~400 y
¶ variable in the α Cyg class (non-radial pulsator):
0.17-0.22 in Hp
¶ E(B-V) = +0.00
composite; Aa: 0.7, Ab: 0.9 0.0-0.1”
Aa,Ab are resp. mags. 0.08, 0.18
¶ under IAU rules, “Capella” designates Aa, not Ab
¶ orbit 104.0 y; first binary with orbit studied
interferometrically (Anderson-Pease, Mt Wilson,
1910); however, full system appears to be
 α Aur Aa+Ab+H+L, where H and L are
red dwarfs sharing the proper motion of Aa+Ab
and perhaps possessing further gravitationally
bound companions (with α Aur B, C, D,
E, F, G, I, J, K, however, being mere line-of-sight
coincidences); more recent interferometry is from
Mt Wilson “Mark III” 1994, Cambridge COAST
1995
¶ α Aur Ab is in rapid evolutionary transition,
currently crossing the Hertzsprung Gap
¶ system is among the brightest of X-ray sources
ecl.: 3.31-3.60 in V, 8.0 d; A: 3.6; B: 4.9, 1.8” (2019)
PA: 87 $^\circ$ -77 $^\circ$, 1848-2019, orbit \geq 2000 y;
so full system is η Ori Aa,Ab,B
¶ system also possibly presents β Cep variability
¶ Aa is an instance of “Be phenomenon”, and additionally
shell spectrum has been observed for Aa
¶ Astron. Alm. (epoch 2021.5) assigns MK type B1 IV

γ	Ori A	5 26.3	+6 22	1.64	-0.22	B2 III ^f	13	-2.8	250	0.015	212	+18 SB?	<p>¶ BSC5: “expanding circumstellar shell”</p> <p>BSC5: “expanding circumstellar shell”</p>	Bellatrix
β	Tau	5 27.7	+28 37	1.65	-0.13	B7 III ^f	24	-1.4	130	0.175	173	+9 V	<p>BSC5: “expanding circumstellar shell”</p> <p>¶ lunar occultations possible as far N as southern California</p> <p>¶ often, but not invariably, classified as Hg–Mn star: Mn 25× solar (and Ca, Mg only ~0.12× solar: radiative lofting, gravitational settling)</p> <p>¶ E(B–V)=0.00</p> <p>B: 7.5, 2.7”, PA:268°→10°, 1875→2017</p>	Elnath
β	Lep A [†]	5 29.2	-20 45	2.81	0.81	G5 II	~20.3	-0.6	160	0.086	183	-14 V?	<p>¶ β Lep B is possibly variable</p> <p>¶ duplicity now suspected also in β Lep A, through 2002 adaptive-optics observation at Haleakala: separation 2.58”</p>	Nihal
δ	Ori Aa [†]	5 33.1	-0 17	2.25v [†]	-0.18	O9.5 II	5	-4.4	700	0.001	137	+16 SB	<p>a good marker of celestial equator</p> <p>¶ eclipsing binary 2.14–2.26, 5.7 d, with secondary of MK temperature type B; since the Ori Aa SB is as yet unresolved, even by interferometry, WDS is not yet able to apply the names “Aa1”, “Aa2”;</p> <p>Ab, resembling Aa-pair secondary in being of MK type B, is at celestial-sphere distance 0.26” from the Aa SB pair</p> <p>¶ yielded first detection of ISM (Hartmann, 1904, through non-moving Ca line in the SB)</p> <p>¶ E(B–V)=+0.07</p>	Mintaka
α	Lep A	5 33.7	-17 48	2.58	0.21	F0 Ib [†]	1.5	-6.6	2000	0.004	72	+24	<p>evolutionary status unclear (has helium fusion already started in core?); helium-fusion past yields now abundances N 5× solar, Na 2× solar</p>	Arneb
β	Dor	5 33.8	-62 29	3.76v [†]	0.64	F7–G2 Ib	3.2	-3.7	1000	0.013	4	+7 V	<p>Cepheid variable: 3.41–4.08 in V, 9.8 d period not quite constant; evolutionary status uncertain</p> <p>¶ observed by <i>FUSE</i>, <i>XMM-Newton</i> missions</p> <p>[THIS STAR ONLY IN ONLINE VERSION OF TABLE]</p>	
λ	Ori A [†]	5 36.3	+9 57	3.39	-0.16	O8 III ^f	3	-4.2~1100		0.004	216	+34	<p>B: 5.45, B0 V, 4.3”, PA:45°→44°, 1779→2019</p> <p>¶ the dominant member of Collinder 69</p> <p>¶ within gas ring 150 ly in diameter (possibly, but not certainly, remnant from a Type II supernova)</p> <p>¶ E(B–V)=+0.12</p>	Meissa
ι	Ori Aa [†]	5 36.5	-5 54	2.75	-0.21	O9 III	~1.4	-6.5	2000	0.001	108	+22 SB2	<p>Aa,Ab 0.1”, mags. 3.0, 6.3</p> <p>B: 7.3, B7 IIIp (He wk), 12.5”, PA:134°→146°, 1779→2018, orbit \geq 700,000 y;</p> <p>ι Ori Aa,Ab 29 d, 0.11 AU min, 0.8 AU max; the elongated orbit, and the disparity in ages, suggest duplicity through many-body interaction-with-expulsion, rather than through co-genesis</p> <p>¶ colliding winds make ι Ori A a strong X-ray source</p> <p>¶ ι Ori B is variable</p> <p>¶ brightest member of Sword asterism</p> <p>¶ E(B–V)=+0.07</p>	Hatysa
ϵ	Ori A	5 37.3	-1 11	1.69	-0.18	B0 Ia	2	-7.2	2000	0.002	118	+26 SB	<p>luminosity (etc) controverted: Crowther (2006) 275,000 L_o, Searle (2008) 537,00 L_o, Puebla (2015) 832,000 L_o</p> <p>¶ E(B–V)=+0.08</p>	Alnilam
ζ	Tau	5 38.9	+21 09	2.97v [†]	-0.15	B2 IIIpe (shell) [†]	7	-2.7	400	0.020	175	+20 SB [†]	<p>2.80–3.17 in V, 133.0 d</p> <p>with also possible variability of the γ Cas type (γ Cas variability would be consistent with the observed Be-phenomenon-cum-shell, and is accepted by AAVSO(VSX), although not accepted throughout the literature)</p> <p>¶ the primary in the SB pairing is one of the best-known “Be phenomenon” stars, and is possibly one of the keys to the solution of currently unsolved Be-phenomenon problems; consistently with the shell-spectrum history, the disk is just 5° away from being seen edge-on</p> <p>(2013A&ARv..21...69R, p. 58n);</p> <p>although the disk gases move in Keplerian orbits, their orbits are not circular, and consequently the material has some nonzero radial velocity even at the midpoint of transit; a further consequence of this kinematics is that the orbiting gas is less dense at apastron than at periastron; shell spectrum underwent three full cycles of V/R variation from 1997 to 2010, with these cycles generally taken as making the precession, under gravitational influence of the elusive SB companion, of a one-armed density wave within the Be disk (for geometry and time variations)</p>	Tianguan

of disk, cf Fig. 7 of [2010AJ....140.1838S](#), Fig. 8 of [2015A&A...576A.112E](#); however, in more recent years, the V/R cycling has been absent; precession notwithstanding, the disk has been observed to be stable, and therefore must be being fed by decretion from the host-star photosphere at a nearly constant rate; as a step toward the eventual discovery of the excitation structure of some conveniently observable Be-phenomenon disk, [2012ApJ...744...19K](#) reports spectro-interferometry from two different ζ Tau primary-star radii, in hydrogen Brackett γ and in a set of hydrogen Pfund lines (while drawing also on hydrogen Balmer α data from previous literature); the emission is found to originate at roughly the same disk radius for hydrogen Balmer α and hydrogen Brackett γ , and at a smaller radius for the hydrogen Pfund lines; the [2012ApJ...744...19K](#) ζ Tau study provides some observational support for the viscous decretion-disk, Keplerian-rotation model prevalent in recent Be-phenomenon theorizing, and additionally supports the density-wave-in-disk hypothesis for the observed V/R cycles; modelling efforts are ongoing, with [2015A&A...576A.112E](#) serving as a progress report

¶ its rapid rotation and Be-phenomenon and shell-spectrum histories notwithstanding, the ζ Tau primary has already evolved some distance off the MS, to “giant” stage (in general, giants are not expected to be rapid rotators); [2012ApJ...744...19K](#) assumes an equatorial radius of $7.7 R_{\odot}$
 ¶ the nature of the elusive (low-flux? $\sim 1 M_{\odot}$ SB companion, of period 133.0 d, is unknown (could even be a neutron star); separation (with orbit nearly circular) is ~ 1.17 AU; since interferometry seems so far to have failed to resolve the companion, WDS, at any rate as of 2021 Jan. 25, is constrained to write “ ζ Tau” rather than “ ζ Tau A” and “ ζ Tau B”; the elusive companion may be producing a truncation in the Be-phenomenon disk, in the sense of a radical change in the dependence of disk density on radius ([2013A&ARv..21...69R](#)); under IAU rules, the name “Tianguan” applies only to the primary, not to the entire SB system

α	Col A	5 40.4	-34 04	2.65 [†]	-0.12	B7 IVe [†]	12	-1.9	260	0.025	176	+35 V?	Phact
													rapid rotator, with mass loss to disk, and so an instance of the “Be phenomenon”; variability, in γ Cas class, has been suspected: H α is variable, and H β profile varies rapidly; nevertheless, the Be disk is stable (unlike, e.g. the Be disk of γ Cas), indicating that the process of decretion-from-photosphere is in this case proceeding at a constant rate
ζ	Ori Aa	5 41.8	-1 56	1.74	-0.20	O9.5 Ib	4	-5.0	700	0.005	58	+18 SB [†]	Alnitak
													¶ E(B-V)=0.00 B: 4.2, B0 III, 2.4", PA:152 $^{\circ}$ →167 $^{\circ}$, 1822→2017 orbit ≥ 1500 y ¶ the secondary in the SB, namely ζ Ori Ab, is of MK type B1 IV, mag. 4.3 ¶ the brightest of the (rare) MK O-type stars ¶ vigorous mass ejection (consistently with membership in MK type O) ¶ E(B-V)=+0.09
ζ	Lep	5 47.9	-14 49	3.55	0.10	A2 Vann [†]	~46.3	1.9	~70.5	0.015	266	+20SB?	
													rapid rotator (period ~ 0.2 d or ~ 0.3 d) ¶ has debris disk, has first known extrasolar asteroid belt ¶ Astron. Alm. (epoch 2021.5) assigns MK type A2 Van ¶ approached to within ~ 4 ly or ~ 5 ly of Sun ~ 1 My ago [THIS STAR ONLY IN ONLINE VERSION OF TABLE]
κ	Ori	5 48.8	-9 40	2.07 [†]	-0.17	B0.5 Ia [†]	5	-4.4	600	0.002	131	+21 V?	Saiph
													evolutionary status unclear, high mass-loss rate; slight variability (0.04 mag) ¶ carbon-deficient (with metallicity otherwise unremarkable) ¶ E(B-V)=+0.07
β	Col	5 51.7	-35 46	3.12	1.15	K1.5 III [†]	37.4	1.0	87	0.408 [†]	8	+89 [†] V	Wazn
													high space velocity indicates that this is interloper from outside galactic thin disk, and yet it is richer than Sun in the elements beyond He
α	Ori Aa [†]	5 56.3	+7 25	0.45v [†]	1.50	M2 Iab [†]	7 [†]	-5.5	500 [†]	0.030 [†]	68	+21 [†] SB	Betelgeuse
													semireg., late-type supergiant var.: $\sim 0-1.7$ in V variability was discovered by J. Herschel in 1839; the latest minimum, early in 2020, at ~ 1.7 in V, exceeded even the minima of 1927 and 1941

(each ~1.2); journalism on this 2020 event includes www.sciencenews.org/article/betel-geuse-star-dim-supernova-death-what-happened; three currently offered explanations are dust cloud from mass ejection, (gigantic) starspot, and fortuitous coincidence of minima from three separate cyclical variations; recovery began 2020 Feb. 22, with a rise to ~0.3 in V by 2020 late April; AAVSO reports mag. 0.65 in V on 2021 Jan. 11; [2018A&A...615A.116M](#) suggests on basis of magnetic variations a scenario on which evolution of giant photospheric convective cells, generating magnetism through local dynamos, is responsible for the observed long secondary ~2100-day photometric period; there are additionally ~200- ~400-day photometric periodicities, plus a stochastic variation ascribed to photospheric granulation

¶ Astron. Alm. (epoch 2021.5) assigns MK type M1–M2 Ia–Iab

¶ brightest star in IR sky, also brightest in bolometric sky

¶ nearest RSG (contrast with o Cet, as AGB); greatest angular diameter of almost any star other than Sun (near-IR limb-darkened disk ~42 mas; but R Dor, having approx 1/3 radius of α Ori, is less distant, and so attains still greater angular diameter); https://en.wikipedia.org/wiki/List_of_largest_stars supplies context, giving radii for many supergiants; reduction of α Ori angular diameter over period 1993/2009 has been asserted

¶ [2017AJ...154...11H](#) reviews the longstanding α Ori distance problem: parallaxes, including *HIPPARCOS*, labour under the difficulty of accurately determining photocentre of visually extended object, awkwardly harbouring even plumes and hotspots; we now give in our table these authors' values for π (rounding from their 4.51 mas) and by implication for D (strictly 717 ly \pm 20%)

¶ very slow rotator (true period difficult; 8.4 y has been suggested)

¶ [2019A&A...628A.101H](#) announces dust halo with inner radius 1.5 R*;

[2016A&A...585A..28K](#) locates 3 R* as the interface between hot-gas and more outlying dust envelopes

¶ CO shells inner 50 R* to 150 R*, outer as far as 250 R*

¶ runaway star, exceeding local speed-of-sound in ISM; bow shock 6-7 arcmin, from stellar wind meeting ISM, plus linear bar at 9 arcmin (it has been suggested that the bar is a relic of collapsing wind from a previous BSG phase, and it also has been suggested that the bar is a feature intrinsic to the embedding ISM, unconnected with any α Ori Aa wind)

¶ although RSG stars pose a more serious mass-loss problem for astrophysics than do the AGB stars, since it is not immediately clear what mechanism is lifting RSG stellar material above the photospheres (convection? pulsation? magnetics?), there is now a possible partial resolution in this particular case: [2018A&A...609A..67K](#), using ALMA, finds α Ori anisotropic mass loss, with plume of ejecta; the authors suggest that plume is associated with strong "rogue" convection cell, observable as photospheric hot spot (in contrast with the cool spots encountered on such MS stars as the Sun)

¶ progenitor mass possibly ~20 M_o (making α Ori very massive), age since arrival on ZAMS possibly 8.0-8.5 My (making α Ori very young)

¶ present evolutionary status of α Ori uncertain: has this RSG previously been a BSG? (and [2017MNRAS.465.2654W](#) suggests history may have been complicated by a stellar merger)

¶ α Ori is SN Type II-P progenitor, the core collapse being due within,

													(perhaps much within) 1 My; although SN will plateau for several months, yielding a star visible even in daytime, with the brilliance of a half Moon or full Moon, the SN radiation from so distant a source will not constitute a terrestrial biohazard ¶ <i>Sky & Telescope</i> . feature article 2019-05 on α Ori can usefully be supplemented with Fig. 13 from 2018A&A...609A..67K (multi-wavelength composite, showing ejecta plume condensing to dust at a few R^* , and showing also two areas of local photospheric magnetic activity): AAVSO has backgrounder at http://www.aavso.org/vsots_alphaori ecl.: 1.89–1.98 in V, 3.96 d (mags. equal) ¶ under IAU naming rules, “Menkalinan” denotes Aa, not Ab
β	Aur Aa,Ab	6 01.1	+44 57	1.90v [†]	0.08	A1 IV + A1 IV	~40.2	-0.1	81	0.056	269	-18 SB2	Menkalinan
θ	Aur A	6 01.2	+37 13	2.65	-0.08	A0p II: (Si) [†]	~19.7	-0.9	166	~0.086	~149	+30 SB	Mahasim B: 7.2, G2 V, 4.2", PA: 7°→303°, 1871→2019 orbit \geq 1200 y, with separation \geq 185 AU ¶ A is magnetic, and an oblique rotator; there are abundance anomalies in photospheric patches, with Si and Cr 10 \times and 100 \times solar, respectively var.: 3.15–3.9 in V, 233 d; B: 6.2, 1.7" (2018) Propus orbit \geq 700 y ¶ variations in A have been variously ascribed either to binarity-eclipse or to Mira-like instability; A has finished core He fusion, and is beginning its ascent up the AGB ¶ liable to lunar, and also to very rare planetary, occultations
η	Gem A [†]	6 16.2	+22 30	3.31v [†]	1.60	M3 III	8	-2.0	400	~0.064	~259	+19 SB	Propus
ζ	CMa A	6 21.1	-30 04	3.02 [†]	-0.16	B2.5 V	9.0	-2.2	360	0.008	61	+32 SB [†]	Furud variability has been claimed (with membership claimed in the β Cep pulsator class) ¶ SB orbit 675 d
β	CMa A	6 23.6	-17 58	1.98v [†]	-0.24	B1 II–III	7	-3.9	~490	0.003	256	+34 SB	Mirzam var. in β Cep class: 1.97–2.00 in V, 0.25130 d (we give here the AAVSO(VSX) period and V-mag. range, as viewed 2021 Aug. 07); the brightest of the β Cep pulsators; has multiple modes, with beat period 50 d; it is not known why ϵ CMa, while physically similar, is not a pulsator ¶ near the boundary of the “Local Bubble” ISM cavity ¶ E(B–V) = +0.01 semiregular variable: 2.75–3.02 in V ¶ on AGB ¶ subject to lunar occultations
μ	Gem A	6 24.3	+22 30	2.87v	1.62	M3 IIIab	14	-1.4	230	0.124	153	+55 V?	Tejat
α	Car	6 24.4	-52 42	-0.62	0.16 [†]	A9 Ib [†]	11	-5.5	~310	0.031	41	+21	Canopus visible both in X-ray (magnetically heated corona; also rapid rotator, strongly convective) and in radio ¶ evolutionary status not fully clear, and colour unusual in its luminosity class ¶ Astron. Alm. (epoch 2021.5) assigns MK type A9 II rapid rotator, with period < 1.7 d shell spectrum has been suggested, with “central quasi-emission peak” (cf 1999A&A...348..831R) ¶ distance was ~27 ly 3.6 My ago
ν	Pup	6 38.4	-43 13	3.17	-0.10	B8 IIIIn [†]	9	-2.1	370	0.004	186	+28 SB	
γ	Gem Aa	6 39.0	+16 23	1.93	0.00	A1 IVs	30	-0.7	110	0.057	166	-13 SB [†]	Alhena SB in highly eccentric orbit, 12.6 y, average separation 8.5 AU; Ab is mag. ~7.5 ¶ the brightest star ever to be observed in an asteroid occultation (381 Myrrha, in 1991) ¶ E(B–V) = +0.03
ϵ	Gem A	6 45.3	+25 06	3.06	1.38 [†]	G8 Ib	4	-4.0	800	0.014	204	+10 SB	Mebstuta unusually yellow in the general population of supergiants ¶ among the few supergiants liable to lunar and planetary occultations B: 8.5, WDA: 10.7" (2016); orbit 50.1 y separation 8.2 AU min (3"), 31.5 AU max (11", in 2019) ¶ IRAS detected IR excess, a signature of dust (rather unexpected in a binary) ¶ Fe abundance of α CMa is ~2 \times or ~3 \times solar ¶ α CMa B is unusually massive for a WD (1.02 M_{\odot} ; Chandrasekhar Limit is, however, 1.4 M_{\odot} ; spectral type of α CMa B is DA (= hydrogen-only)) ¶ E(B–V) = -0.03
α	CMa A [†]	6 46.1	-16 45	-1.44	0.01	A0mA1 Va [†]	~379	1.5	8.6	~1.339	~204	-8 SB	Sirius
ζ	Gem	6 46.5	+12 52	3.35	0.44	F5 IV	56	2.1	58.7	0.223	211	+25 V? [†]	Alzirn possibly SB, with components of ~equal mass ¶ rapid rotator (but just barely over the internal-structure transition, or “F5 rotation break,” that causes some stars to rotate rapidly, others to experience braking through magnetics and winds) ¶ X-ray source (suggesting significant corona)

α	Pic	6 48.4	-61 58	3.24	0.22	A6 Vn [†]	~34	0.9	100	0.252	345	+21	rapid rotator; shell, with time-varying spectral absorption features ¶ X-ray emission suggests a companion, otherwise undetected
τ	Pup	6 50.5	-50 38	2.94	1.21	K1 III	18	-0.8	180	0.077	154	+36 SB [†]	SB period 1066.0 d, separation ~3 AU, orbit of low eccentricity; since the SB is as yet unresolved, even in interferometry, WDS is as yet unable to write “ τ Pup A” and “ τ Pup B”
κ	CMa	6 50.6	-32 32	3.50v [†]	-0.12	B1.5 IVne [†]	4.9	-3.0	700	0.010	293	+14	var. in γ Cas class: 3.40–3.97 in V (was at faint end of its range before 1963; AAVSO reports visually ~3.3 in 2021 Jan.); an instance of the “Be phenomenon”
ε	CMa A [†]	6 59.5	-29 00	1.50	-0.21	B2 II	8.0	-4.0	410	0.004	68	+27	binary (7.9”; B is mag. ~7.5) separation 900 AU, period at least 7500 y ¶ brightest known source of extreme UV (~75 nm) in Earth’s night sky; hydrogen Lyman α (121.6 nm) observed by NASA OAO-3 ¶ E(B–V) = +0.02
σ	CMa A	7 02.6	-27 58	3.49v	1.73	K7 Ib [†]	3	-4.2	1100	0.008	308	+22	irregular var.: 3.41–3.51 in V Unurgunite authorities are in some disagreement on MK type (possibly M, rather than K) [THIS STAR ONLY IN ONLINE VERSION OF TABLE]
ρ^2	CMa	7 03.9	-23 52	3.02 [†]	-0.08	B3 Ia	1	-6.6	3000	0.004	329	+48 SB	var. in α Cyg class: 2.98–3.04 in Hp, 24.44 d the α Cyg vars. are non-radial pulsators ¶ E(B–V) = +0.03
δ	CMa	7 09.3	-26 26	1.83	0.67	F8 Ia [†]	2	-6.6	2000	0.005	317	+34 SB	¶ slow rotator (possibly ~1 y); N 2 \times solar, Na 6 \times solar semireg. late-type var., 2.6–8.0 in V, 140.6 d Wezen HR2748
L ₂	Pup A	7 14.2	-44 41	4.42v	1.33	M5 IIIe	16	0.4	210	0.342	18	+53 V?	B: 7.9, 66”, PA: 214°→213°, 1826→2009
π	Pup Aa	7 17.9	-37 08	2.71v [†]	1.62	K3 Ib	4	-4.3	800	0.012	303	+16	¶ semiregular variable.: 2.70–2.85 B: 8.2, K3 V, 5.4”, PA: 198°→229°, 1822→2018 Wasat orbit 1200 y
δ	Gem A	7 21.4	+21 56	3.50	0.37	F0 IV [†]	54	2.2	60	0.018	237	+4 SB [†]	¶ lunar occultations possible; planetary occultations possible-yet-rare ¶ in evolutionary transition, having completed stable core-hydrogen fusion ¶ Astron. Alm. (epoch 2021.5) assigns MK type F0 V [†] [THIS STAR ONLY IN ONLINE VERSION OF TABLE]
η	CMa A	7 24.9	-29 21	2.45v [†]	-0.08	B5 Ia	2	-6.5	2000	0.007	325	+41 V	B: 6.8, 178” (2010) is mere optical companion Aludra ¶ variable in α Cyg class of non-radial pulsators; AAVSO(VSX) as at 2021 Jan. 28 gives mag. range 2.36–2.50 in V, period 4.70433 d ¶ strong wind; ejected circumstellar mass inferred from IR excess ¶ E(B–V) = +0.02
β	CMi A	7 28.3	+8 15	2.89v [†]	-0.10	B8 Ve [†]	~20.2	-0.6	~162	0.064	234	+22 SB	Gomeisa rapid rotator, possibly ~1 d, with modest variability in the hydrogen Balmer emission; disk of ejected matter has diameter ~4 \times diameter of β CMi itself (BSC5: “rotationally unstable”); an instance of the “Be phenomenon”; although GCVS and AAVSO(VSX) assertion of γ Cas-type variability has not been corroborated, <u>2007ApJ...654..544S</u> reports, using <i>MOST</i> , millimagnitude “slowly pulsating B-type” variability; AAVSO(VSX) as viewed 2021 Jan. 16 gives V-mag. range 2.84–2.92; in contrast with e.g. the Be-phenomenon star γ Cas A, the Be disk is in this case considered very stable (<u>2013A&ARv.21...69R</u>), indicating constancy in the process of decretion from the host-star photosphere; <u>2012ApJ...744...19K</u> reports confirmation of Keplerian rotation in the Be disk (an important follow-on to the discovery of Keplerian rotation in Be-phenomenon star α Ara A) B: 8.8, G5: V, 22.1”, PA: 90°→74°, 1826→2015 orbit \geq 27,000 y, separation \geq 1300 AU; SB is eclipsing, of β Lyr type, with orbit 257.8 d, with very modest alternating primary (0.04 mag) and secondary (0.03 mag) minima; the SB primary component shows slow irregular variability ¶ system has high space velocity orbit 445 y; max = 6.5”, in 1880; min = 1.8”, in 1965; 5.5” (2019); separation 71 AU min, 138 AU max; C mag. 9.8; AC PA: 162°→163°, 1822→2017, 70”, orbit \geq 14,000 y; C has variable-star name YY Gem (an eclipsing binary, and additionally a variable of the BY Dra class, with flaring); not only C, but also each of A, B is itself SB,
σ	Pup A	7 29.9	-43 21	3.25 [†]	1.51	K5 III	17	-0.6	190	0.198	342	+88 SB [†]	
α	Gem A [†]	7 36.0	+31 50	1.93	0.03	A1mA2 Va	63	0.9	52	-0.254	~234	+6 SB	
α	Gem B [†]	7 36.0	+31 50	2.97	0.03	A2mA5 V:	63	2.0	52	-0.254	~234	-1 SB	Castor

α	CMi A [†]	7 40.4	+5 10	0.40 [†]	0.43	F5 IV–V	285	2.7	11.5	~1.259	~215	–3 SB	Procyon
													<p>making ABC a hierarchical 6-star system (Kaler at http://stars.astro.illinois.edu/sow/castor.html writes, “certainly the sky’s ranking sextuple”); https://en.wikipedia.org/wiki/Castor_(star) has a diagram summarizing this sextuple hierarchy, on the basis of 2012MNRAS.423.493H; since the A SB is not yet resolved (even interferometrically) and since the B SB is not yet resolved (even interferometrically), WDS is not yet able to write “Aa,Ab” and is not yet able to write “Ba,Bb”</p> <p>¶ Castor–Pollux comparison is a helpful test of naked-eye night colour response</p> <p>B: 10.8, WD; 3.8” (2014); orbit 41 y separation 8.9 AU min, 21.0 AU max</p> <p>¶ astroseismology of A is somewhat uncertain (<i>MOST</i> mission 2004 did not find pulsations, and yet <i>WIRE</i> mission 1999 and 2000 did)</p> <p>¶ the WD Procyon B is physically unlike the WD Sirius B, attaining only ~0.2 of the Sirius B density, and being of rare spectral type DQZ</p>
β	Gem A+1P [†]	7 46.6	+27 58	1.16	0.99	K0 IIIb	97	1.1	33.8	0.628	266	+3 V	Pollux
													<p>the nearest of the giants; unusual in being a giant known to harbour an exoplanet (and the brightest known exoplanet host in Earth’s sky); as of 2015, exoplanet is IAU-named “Thestias”</p> <p>¶ subject to rare lunar occultations, for observers S of Earth’s equator</p> <p>¶ Castor–Pollux comparison is a helpful test of naked-eye night colour response</p>
ξ	Pup A [†]	7 50.2	–24 55	3.34	1.22	G6 Iab–Ib [†]	3	–4.5	1200	0.005	260	+3 SB [†]	Azmid
													<p>full system is SB with B (mag. 13, ~5”, orbit $\geq 26,000$ y)</p> <p>¶ SB primary has high metallicity, with exact evolutionary status uncertain</p> <p>¶ SB primary is near, but is a little too cool to lie within the HR diagram Instability Strip</p>
χ	Car	7 57.3	–53 02	3.46 [†]	–0.18	B3 IV(p?) [†]	7	–2.3	500	0.035	304	+19 V	
													<p>Si II anomalous strength now discounted</p> <p>¶ suggestion of variability now discounted, via <i>HIPPARCOS</i></p> <p>¶ the MK luminosity class “IV” (phenomenologically “giant”) notwithstanding, χ Car is in astrophysical terms in the last part of its stable core-hydrogen-fusion phase; Astron. Alm. (epoch 2021.5) assigns MK temperature type B3p Si without assigning an MK luminosity class</p>
ζ	Pup	8 04.3	–40 04	2.21 [†]	–0.27 [†]	O5 Iafn [†]	3.0 [†]	–5.4	1080 [†]	0.034 [†]	299	–24 [†] V?	Naos
													<p>¶ rapid rotator (1.78 d), despite ~2300 km/s stellar wind (in which spiral structure was announced in 2017 by <i>BRITTE</i> mission team), with mass loss rate $> 1e-6 M_{\odot}/y$</p> <p>¶ high space velocity (impelled by past nearby supernova? or, rather, impelled by multibody gravitational interactions in its stellar birth family?); possibly ejected from Trumpler 10 OB association</p> <p>¶ distance has been controverted</p> <p>¶ He, N overabundant</p> <p>¶ has been suspected of being a variable in the α Cyg class</p>
ρ	Pup A	8 08.5	–24 22	2.83v [†]	0.46	F2mF5 II: (var) [†]	51.3	1.4	64	0.095	299	+46 SB	Tureis
													<p>¶ E(B–V) = +0.04</p> <p>var.: 2.68–2.87 in V, 0.14 d</p> <p>prototype of the “p Pup stars” (these combine δ Sct variability with Am-like abundance anomalies); main period is ~3.3 h (0.15 mag.); photosphere temperature is notably low in the overall population of stars presenting δ Sct variability</p> <p>¶ Astron. Alm. (epoch 2021.5) assigns MK type F5 (Ib–II)p</p>
γ	Vel Aa [†]	8 10.2	–47 24	1.75v	–0.14	O7.5 III-I [†]	3 [†]	–5.9	~1100 [†]	0.012	330	+35 SB2 [†]	
													<p>¶ IR excess (circumstellar ring, at separation 50 AU?) eruptive var.: 1.81–1.87 in V; Aa,Ab system is a.k.a. γ^2 Vel</p> <p>¶ strictly a quadruple system, comprising the SB Aa,Ab pair (period 78.5 d) and the tighter SB Ba,Bb pair (a.k.a. γ^1 Vel, period 1.48d); separation of these two pairs, i.e. of “AB”, 42.9”\rightarrow41.2”, 1826\rightarrow2017; PA: 222$\circ$$\rightarrow221\circ$, 1826$\rightarrow$2017</p> <p>¶ Astron. Alm. (epoch 2021.5) assigns MK type “O9 I:”</p> <p>¶ the (carbon-rich) WR component γ Vel Ab, of spectral type WC8, is the nearest and visually brightest of all WR stars, and is an exceptionally massive WR</p>

(9.0 M_{\odot} ; but at birth, $> 30 M_{\odot}$);
the Aa,Ab pair is the best studied of all O-WR binaries:
in the SB γ Vel Aa,Ab pair
(orbit 78.5 d, separation 0.8 AU min?
1.6 AU max), γ Vel Ab
dominates spectrally, making
the γ Vel Aa,Ab SB the “Spectral
Gem of the Southern Skies,” and a notable sight within
the broader “Vela complex” (dominated by the
the Gum Nebula, within which lie the Vela SNR, the *IRAS*
Vela shell, and the Vela
pulsar: some literature, including [2011A&A...525A.154S](#),
indeed proposes intersection
between the Vela SNR and a γ Vel SWB,
taking the *IRAS* Vela shell as marking the meeting of
SNR and SWB); nevertheless,
the V-band light is overwhelmingly
from the more massive (28.5 M_{\odot})
O-type component γ Vel Aa
¶ like η Car (bright to mag. ~ 0 for several years after 1837,
but now too faint, and now too lacking in firm
future-outburst prognoses, to qualify
for the RASC Handbook “Brightest Stars” list),
 γ Vel Aa,Ab is a colliding-wind pair
([2017MNRAS.468.2655L](#) Fig. 1 sketches the collision
geometry), and in consequence is a UV and X-ray source
(and in consequence may also possibly resemble η Car
in being a γ -ray source
(cf [2017ApJ...847...40R](#); as of
at any rate 2017, it seems that
no other colliding-winds-binary
stellar γ -ray sources are known));
it is the wind from Ab that dominates, with mass-loss
rate at least 100 \times greater than for Aa;
the Ab wind may feature some clumping,
but is to a good approximation
spherically symmetric until it
encounters the γ Vel Aa wind;
orbital motion of Aa,Ab around centre of mass
yields a spiral structure in the wind-collision area,
particularly salient during periastron
¶ [2017ApJ...847...40R](#) summarizes recent observations
of Aa,Ab Vel, in radio and IR
and optical, including interferometry,
noting inter alia discrepancies in the available determinations
of mass-loss rates from Ab (a copious 3e-6 M_{\odot} /y?
or a still more copious 8e-5 M_{\odot} /y?)
¶ notable among recent observational studies are
[2017MNRAS.468.2655L](#) (VLTI/AMBER
near-IR spectro-interferometry,
with also 3-D hydrodynamic modelling)
and [2012MNRAS.427.581R](#)
¶ likely destiny of γ Vel Ab is as (exotic) stripped-core
SN (same prognosis as for η Car; this
contrasts with α Ori, which
will for its part instead explode as a (not exotic)
hydrogen-spectrum SN)
¶ dust emission is absent (even though
formation of circumstellar dust is common
in stars that, like γ Vel Ab, undergo
copious mass outflow)
¶ distance ~ 1200 ly, in contrast with our
 ~ 1100 ly, has also been recently asserted,
on basis of VLTI/AMBER
¶ we take MK type for γ Vel Aa from
[1999A&A...345..163D](#) (as what
must be considered an emendation
of our (slightly cooler)
Garrison-approved MK type
from earlier editions of this table;
admittedly, MK determination of γ Vel Aa is still
difficult, because the raw spectrum
is a composite comprising
not only the two stars Aa, Ab, but also emission from the
wind-collision zone)
¶ neither the traditional Suhail al-Muhlif nor
the modern Regor (devised within NASA, to
commemorate 1967 fire victim Roger Chaffee) is
presently IAU-approved name for any of the four stars
 γ Vel Aa, Ab, Ba, Bb

													as contamination from defunct companion (but no companion remnant has been found)
													¶ Astron. Alm. (epoch 2021.5) assigns MK type K4 III Ba 0.5
													[THIS STAR ONLY IN ONLINE VERSION OF TABLE]
ε	Car A	8 23.0	-59 35	1.86v [†]	1.20	K3:III	5	-4.5	600	0.034	311	+2	ecl.: 1.82–1.94 in V period possibly 785 d; separation ~4 AU, precluding mass transfer; B is mag. ~3.9, MK type (uncertainty-flagged) “B2: V” var.?: 3.30?–3.36 in V? ¶ currently in rapid evolutionary transition, crossing the Hertzsprung Gap ¶ despite high space velocity, a member of the galaxy thin disk Avior
o	UMa A+1P	8 32.0	+60 39	3.35v [?]	0.86	G5 III	~18.2	-0.3	~179	0.172 [†]	231	+20 [†]	¶ Aa, Ab brightest known ecl. binary (dim by ~0.4) B: 5.0, 0.8”, PA: 177°→195°, 1894→2019 AB orbit 142 y (min angular distance was in 2000) ¶ Aa, Ab resolved both interferometrically and with VLT adaptive optics; orbit 45.15 d, average separation 90.61 AU Ashlesha
δ	Vel Aa [†]	8 45.3	-54 47	1.93v [†]	0.04	A1 Va	40	0.0	81	~0.107	~164	+2 V [?]	composite A: 3.8; B: 4.7, 0.2” (2018); C: 7.8, 2.9” (2020); B is of poorly known MK type “A:”; AB orbit 15.09 y, AB+C orbit 590 y ¶ C is SB, orbit 9.9 d
ε	Hya A [†]	8 47.9	+6 20	3.38	0.68	G5:III	25	0.4	130	~0.232	259	+36 SB [†]	¶ Astron. Alm. (epoch 2021.5) assigns MK type G9 IIIa A+BC 2.4”, PA: 349°→90°, 1831→2017 A+BC orbit 818 y; BC 0.9”, period ~39 y; A is itself SB, orbit 4028 d, making this a quadruple system; the system is not, as in many cases of multiplicity, hierarchical and stable, but kinematically unstable (disruption in ~0.1 My?); B mag. 9.9 M1 V, C mag. 10.1 M1 V; since the A SB has not yet been resolved, even interferometrically, WDS is not yet able to write “ι UMa Aa”, “ι UMa Ab” semireg. var.: 2.14–2.30 in V ¶ probably on or approaching AGB, but could still be on RGB ¶ Astron. Alm. (epoch 2021.5) assigns MK type K4.5 Ib ¶ has slow wind, whose origins are said to be poorly understood ecl.: 3.41–3.44 in V ¶ orbit 6.74 d, with light curve indicating tidal distortion; since the SB is as yet unresolved, even in interferometry, WDS is not yet able to write “a Car A,” “a Car B” ¶ there is some uncertainty whether observable light is solely from primary, or whether primary and secondary make approximately equal contributions ¶ not to be confused with α Car Suhail
ζ	Hya	8 56.5	+5 52	3.11	0.98	G9 II–III	~19.5	-0.4	~167	0.101	279	+23	rapid rotator (< 2.1 d), despite having finished stable core hydrogen fusion ¶ quasi-periodic variation, ~0.5 h, in hydrogen Balmer lines var.: 2.23–2.28 in V ¶ despite being slow rotator, has magnetic activity (as inferred from X-rays) ¶ not to be confused with I (letter el) Car B: 8.8, 223”, PA: 33°→43°, 1823→2016 suspected var., mag. 3.12–3.17 (beginning to evolve into a Mira?) orbit 116.65 d, average separation possibly ~1.1 AU; since the SB has not yet been resolved (even interferometrically), WDS is not yet able to write “κ Vel A” and “κ Vel B” ¶ mass loss rate ~1e–9 M _⊙ /y ¶ system is X-ray source ¶ ISM absorption has varied over the years (ISM cloud in transit?) Markeb
ι	UMa A [†]	9 00.7	+47 57	3.12	0.22	A7 IVn	~68.9	2.3	47.3	~0.491	~244	+9 SB [†]	slow rotator (possibly 2.4 y), with Ba mildly overabundant ¶ astroseismology has been studied ¶ α Hya B (mag. 9.7; 284”, PA: 55°→155°, 1833→2015) might be a true binary component (with orbit ≥ 870,000 y, separation ≥ 15,700 AU) semiregular variable, 3.12–3.18 in V, 82.0 d ¶ evolutionary status uncertain (helium core fusion impending, or already ended?) Talitha
λ	Vel A	9 08.8	-43 31	2.23v	1.66	K4 Ib–IIa [†]	6.0	-3.9	540	0.028	299	+18	luminosity class, and also SB status, have been controverted, with postulated SB companion
a [†]	Car	9 11.5	-59 03	3.43v [†]	-0.19	B2 IV–V	7	-2.3	500	0.022	312	+23 SB2 [†]	
β	Car	9 13.4	-69 48	1.67	0.07	A1 III	28.8 [†]	-1.0	113	0.191	305	-5 V [?]	
ι [†]	Car	9 17.7	-59 22	2.21v	0.19	A7 Ib	4.3	-4.6	800	0.022	302	+13	
α	Lyn A	9 22.4	+34 18	3.14 [†]	1.55	K7 IIIab	16	-0.8	~203	0.224	274	+38	
κ	Vel	9 22.8	-55 06	2.47	-0.14	B2 IV–V	6	-3.8	600	0.016	315	+22 SB [†]	
α	Hya A [†]	9 28.6	-8 45	1.99	1.44	K3 II–III [†]	18	-1.7	180	0.038	336	-4 V [?]	
N	Vel	9 31.9	-57 08	3.16	1.54	K5 III	13.6	-1.2	240	0.033	280	-14	
θ	UMa A	9 34.3	+51 35	3.17	0.48	F6 IV [†]	74.2	2.5	44.0	1.088	241	+15 SB [†]	

o	Leo Aa [†]	9 42.3	+9 48	3.52v	0.52	F5 II [†]	25	0.5	130	0.148	255	+27 SB [†]	<p>remaining undetected in speckle interferometry eclipsing binary 14.5 d</p> <p>orbit 14.5 d, separation 0.165 AU (interferometrically resolved), with Ab poorly known (MK possibly A5)</p> <p>¶ a rare instance of a star that has ended core hydrogen fusion, and yet in which the convection typical of an evolved star has not yet removed the chemical peculiarities possible in a core-hydrogen fuser (where still-quiet atmosphere facilitates radiative lofting and gravitational settling)</p> <p>[THIS STAR ONLY IN ONLINE VERSION OF TABLE]</p>	Subra
l [†]	Car	9 45.8	-62 36	3.69v [†]	1.01	F9-G5 Ib	2	-4.7	2000	0.015	302	+3 V	<p>Cepheid variable: 3.28-4.18 in V, 36 d</p> <p>AAVSO(VSX) viewed 2021 Jan. 18 gives period 35.551609 d; an exceptionally luminous, and consequently exceptionally slow, Cepheid (compare both the visual brightness and the intrinsic luminosity with less dramatic δ Cep A (in this table), η Aql A (Okab; in this table), and ζ Gem Aa (Mekbuda; almost, but not quite, bright enough for inclusion in this table); Kaler remarks that "if Carina had been in the northern hemisphere, the collection of these variables might well have been called the 'Carinids'"); radius, in its pulsation cycle, has been measured as 160 R_o min, 194 R_o max</p> <p>¶ circumstellar envelope of ejected matter, radius 10 AU-100 AU</p> <p>¶ lower-case ell Car; not to be confused with i (lower-case i) Car (HR 3663), ι Car (HR 3699), L Car (HR 4089), I (upper-case i) Car (HR 4102) (and note additionally that Bayer nomenclature does not use the label "λ Car")</p> <p>[THIS STAR ONLY IN ONLINE VERSION OF TABLE]</p>	HR 3884
ε	Leo	9 47.1	+23 40	2.97 [†]	0.81	G1 II	13.2	-1.4	250	0.047	259	+4 V?	<p>slow rotator, period possibly as long as 200 d</p> <p>¶ currently residing in the Hertzsprung Gap?</p> <p>¶ variability has been studied (cf Andrievsky 1998; pulsation as in Cepheids?)</p> <p>¶ the Arabic or quasi-Arabic name Algenubi (more classically, al Ras al Asad al Janubiyah et al.) is not presently IAU-official</p>	
v	Car A	9 47.6	-65 10	2.92	0.29	A6 II	2.3 [†]	-5.3~1400 [†]		0.028	307	+14	<p>A: 3.01; B: 5.99, B7 III, 5.1", PA:126°→128°, 1836→2015 orbit ≥ 19,500 y, separation ~2000 AU</p> <p>¶ the duplicity causes parallax to be poorly known</p>	
φ	Vel A	9 57.6	-54 40	3.52	-0.07	B5 Ib	2.0	-4.9	1600	0.014	285	+14	<p>[THIS STAR ONLY IN ONLINE VERSION OF TABLE]</p>	
η	Leo A	10 08.5	+16 39	3.48	-0.03	A0 Ib [†]	3	-4.5	1300	~0.003	n.a.	+3 V	<p>B: 8.4, 0.4", PA:84°→239°, 1937→2015</p> <p>¶ mass-loss rate ~5e-8 M_o/y (> 10,000× solar mass-loss rate); BSC5: "chromospheric shell"</p> <p>¶ a lunar occultation has suggested duplicity, but this is unconfirmed</p>	
α	Leo A [†]	10 09.5	+11 52	1.36	-0.09	B8 IVn [†]	41	-0.6	79	0.249	271	+6 SB [†]	<p>α Leo A is SB orbit 40.11 d, with the secondary in the pair that is α Leo A now detected (2011BVS.5987....1R reported null photometry result from MOST, at the high precision of ~0.5 millimag., but a spectroscopic detection is reported in 2020ApJ...902...25G; since the secondary is not yet resolved, even interferometrically, WDS is not yet able to write "α Leo Aa" and "α Leo Ab")</p> <p>¶ the primary in α Leo A is an exceptionally rapid rotator (15.9 h), making the star an oblate spheroid (R_{pol} ~3.14 R_o, but Req ~4.16 R_o) and rendering the photosphere equator ~3000 K cooler than the photosphere poles (and possibly inducing meridional flow in the envelope); this is the first rotating star not in an eclipsing binary system to have its gravitational low-latitudes darkening detected, and the first to have its inclination angle and low-latitudes darkening measured through a direct application of spectroscopy-constrained interferometry (inaugural science run of CHARA, 2005ApJ...628.439M); in contrast with pole-on rapid rotators such as α Lyr A, the α Leo A primary is seen nearly equator-on; in 2011ApJ...732...68C, Fig. 5 presents an image as fitted to CHARA interferometry (the luminosity contours display the disturbing effect of limb darkening upon the rotation-induced gravity darkening; since the poles are near the limb, the brightest regions, as viewed from Earth, do not quite coincide with the poles); the aperture-synthesis imaging of 2017NatAs...1..690C Fig. 5 displays the photosphere temperature variation</p>	Regulus

(a joint consequence of limb darkening and oblateness), along with oblateness and axis orientation; according to [2017NatAs...1..690C](#), the α Leo A primary (*i*) has attained 96.5% of its breakup speed (earlier literature had suggested 86%), and (*ii*) is the first rapid rotator found to exhibit Chandrasekhar rotation-induced stellar limb polarization (the related phenomenon of eclipse-induced stellar limb polarization was admittedly detected earlier, with β Per, as reported in [1983ApJ...273L..85K](#))

¶ despite the large rotation-induced latitude variation in photospheric effective temperature, at all latitudes the envelope is radiative (since the photospheric effective temperature, even at the equator, never falls so low as to approach the ~ 8300 K radiation-to-convection transition value); rotation-induced meridional circulation, on the other hand, disturbs the usual radiative-equilibrium picture of a radiative envelope ([2011ApJ...732...68C](#), p. 11a); since meridional circulation transfers angular momentum, the envelope cannot be presumed to be in solid-body rotation

¶ the rapid rotation, the membership in MK type B, and the near-MS evolutionary status notwithstanding, the question of Be-phenomenon behaviour is answered in the negative by [2005ApJ...628..439M](#); the authors do, however, remark on p. 446 that the historical record contains a lone report of marginal hydrogen Balmer- α emission, from February 1981 (might amateur-spectroscopist monitoring now be advisable?)

¶ [2011ApJ...732...68C](#) revises the mass of the primary upward, offering $4.15 M_{\odot}$ in place of the [2005ApJ...628..439M](#) determination of $\sim 3.5 M_{\odot}$

¶ A+BC almost unchanged since 1779 (179"; PA: $307^{\circ} \rightarrow 304^{\circ}$, 1779 \rightarrow 2019); separation ≥ 4200 AU, orbit $\geq 125,000$ y; BC combined light is mag. ~ 8.2 ; BC is no longer underobserved (PA: $89^{\circ} \rightarrow 94^{\circ}$, $4.0'' \rightarrow 2.20''$, 1867 \rightarrow 2019, with orbit ≥ 880 y)

¶ a puzzling discrepancy between the ages of the α Leo SB primary and α Leo B (surely condensed from the same ISM cloud, at the same time) is perhaps to be explained by the peculiarities in the evolution of rapid rotators

¶ we adopt here the MK classification of [2003AJ...126.2048G](#), while recalling that earlier editions of our RASC brightest-stars table used instead B7 Vn, essentially in accordance with [1953ApJ...117..313](#); Astron. Alm. (epoch 2021.5) likewise assigns MK type B7 Vn

¶ the α Leo system is occasionally occulted by Mercury, Venus (e.g. 1959 Jul. 07, 2044 Oct. 01), Moon (e.g. 2017 Sep. 18; [1972JBA...82..431K](#) describes the 18.6-year 1940-through-2050 cycle of possibilities), and asteroids (e.g. 166 Rhodope 2005 Oct. 19 ([2008mgm.conf.2594S](#) reports GTR effect of light bending, not only from general solar gravitational field but also from Rhodope field), 163 Erigone 2014 Mar. 20 (cloud-defeated 2014 Erigone campaign is documented at <https://occultations.org/regulus2014>))

¶ E(B-V) = +0.01 or "IIIne"; shell star

¶ rapid rotator (< 1.2 d, $\sim 85\%$ of breakup speed); instance of "Be phenomenon"; photometric variation (cp γ Cas, δ Sco, ...) might be expected, and yet seems undocumented; BSC5 does report variable hydrogen Balmer- α irregular variable: 3.36–3.44 in V

¶ metallicity is uncertain

¶ evolutionary state is uncertain (has core already started He fusion?)

¶ Astron. Alm. (epoch 2021.5) assigns MK type K2.5 II B (6.0, 331" in 2015) is mere optical companion Adhafera

ω	Car	10 14.2 –70 09	3.29 [†] –0.07	B8 III n [†]	9.5 –1.8 340	0.037 281	+7 V
q	Car A	10 17.8 –61 26	3.39 [†] 1.54	K3 II a [†]	5.0 –3.1 660	0.026 286	+8
ζ	Leo A	10 17.9 +23 19	3.43 0.31	F0 III a [†]	12 –1.2 270	0.020 110	–16 SB

HR 4050

λ	UMa	10 18.4 +42 48	3.45	0.03	A1 IV [†]	24	0.3	140	0.186	256	+18 V	<p>¶ Astron. Alm. (epoch 2021.5) assigns MK type F0 III</p> <p>¶ in rapid evolutionary transition, currently residing in Hertzsprung Gap</p> <p>Tania Borealis</p> <p>despite MK luminosity class “IV”, has not yet finished core hydrogen fusion</p> <p>¶ mildly metallic, being insufficiently metallic to warrant MK “Am”</p> <p>¶ seems mild IR excess (indicating circumstellar debris)</p>
γ	Leo A +1P [†]	10 21.2 +19 44	2.61 [†]	1.13	K1 IIIb Fe-0.5 [†]	26	-0.3	130	-0.333 [†]	~118	-37 [†] SB	<p>4.7” (2020), PA:99° → 127°, 1820→ 2020 (510.3 y);</p>
γ	Leo B	10 21.2 +19 44	3.16 [†]	1.42	G7 III Fe-1 [†]	26	0.2	130	-0.346 [†]	~118	-36 [†] V	<p>max = ~5”, around 2100</p> <p>Algieba</p> <p>separation \geq 170 AU, orbit > 500 y, orbital parameters not yet well known</p> <p>¶ A, B are of mildly unequal masses, and therefore are of mildly disparate evolutionary stage (Kaler http://stars.astro.illinois.edu/: “best understood as being in different stages of gianthood”; cf this same source for further discussion of the uncertainties in various γ Leo parameters, including the respective mags of A and B)</p> <p>¶ γ Leo A “+ 1P” is an exception to the tendency for exoplanets to be found around the more metallic stars (but the “+1P” could be modelled as a brown dwarf); and indeed even “+2P” is now considered possible</p> <p>¶ high space velocity of the γ Leo AB pair, plus their low metallicity, suggests system is interloper from more remote galactic region</p> <p>¶ γ Leo AB, and indeed also the next “Sickle” star ζ Leo, serve to mark the radiant of the Leonids meteor shower</p>
μ	UMa	10 23.6 +41 23	3.06v [†]	1.60	M0 IIIp [†]	14	-1.2	230	0.089	293	-21 SB [†]	<p>Tania Australis</p> <p>var. 3.03–3.10 in V</p> <p>AAVSO(SVX), viewed 2021 Jan. 16, considers the system to be presenting both eclipse variability resembling the β Lyr case and slow irregular evolved-star variability</p> <p>¶ SB period 230 d; since the SB is not yet resolved, even interferometrically, WDS cannot yet write “μ UMa A” and “μ UMa B”;</p> <p>¶ Ca II emission</p> <p>¶ Astron. Alm. (epoch 2021.5) assigns MK type M0 III</p> <p>¶ Kaler (http://stars.astro.illinois.edu/sow-taniaas.html) terms this “a rare ‘hybrid star’” (in the sense of blowing both a fast-and-thin wind and a slower-and-dense wind), and additionally notes the puzzle posed by X-ray emission in the presence of cool photosphere</p>
ρ	Car	10 32.8 -61 48	3.30v	-0.09	B4 Vne [†]	7	-2.6	500	0.021	304	+26	<p>var. in γ Cas class: 3.22–3.55 in V</p> <p>HR 4140</p> <p>and instance of the “Be phenomenon”</p> <p>¶ fast rotator;</p>
θ	Car	10 43.7 -64 30	2.74	-0.22	B0.5 Vp	7	-3.0	460	0.022	303	+24 SB [†]	<p>BSC5: shell; variable hydrogen Balmer-line profiles chemically anomalous</p> <p>SB period 2.2 d is unusually short, suggesting that mass transfer could be the culprit in the anomalies</p> <p>¶ since the SB is as yet unresolved, even interferometrically, WDS is not yet able to write “θ Car A,” “θ Car B”</p> <p>¶ the primary is the brightest of the “blue stragglers”; at http://stars.astro.illinois.edu/sow/thetacar.html, Kaler discusses difficulties in determination of the primary’s temperature and of its (short) rotation period</p>
μ	Vel A [†]	10 47.7 -49 32	2.82	1.07	G5 III [†]	28	-0.1	~117	0.083	131	+6 SB	<p>¶ E(B-V) = +0.06</p> <p>A: 2.82; B: 5.65, 2.8”, PA:55°→57°, 1880→2016 period variously given as 116.24 y (Hoffleit) and 138 y (Heintz); separation possibly 8 AU min, 93 AU max, 51 AU average</p> <p>¶ B is of MK type F8:V</p> <p>¶ A is in rapid evolutionary transition, having recently finished core hydrogen fusion</p> <p>¶ A is magnetic, and an X-ray emitter, with hot corona, and with violent 2-day X-ray flare detected in 1998 by IUE</p>
ν	Hya	10 50.7 -16 18	3.11	1.23	K2 III [†]	23	-0.1	144	0.220 [†]	25	-1 [†]	<p>slow rotator (but \leq 619 d)</p> <p>¶ low metallicity and high space velocity suggest interloper, born outside Sun’s neighbourhood</p> <p>¶ Astron. Alm. (epoch 2021.5) assigns MK type K1.5 IIIb Hδ-0.5</p>
β	UMa	11 03.1 +56 16	2.34	0.03	A0mA1 IV-V	~40.9	0.4	80	0.088	68	-12 SB	<p>Merak</p> <p>debris disk first detected via IR excess, now marginally resolved by <i>Herschel Space Observatory</i> (2010A&A...518L135M)</p>
α	UMa A [†]	11 05.0 +61 38	1.81 [†]	1.06	K0 IIIa	27	-1.1	120	0.139	255	-9 SB	<p>A: 1.86; B: 5.0, A8 V, 0.8” (2017), PA 342°</p> <p>Dubhe</p> <p>orbit 44 y</p> <p>¶ the first cool star found to have multimodal oscillations (WIRE camera; 2000ApJ...532L133B suggests</p>

													fundamental mode 6.35 d)
													¶ the most distant of the seven Big Dipper stars (and, like η UMa at the other extreme of the Big Dipper, not a member of the same-age association that is the UMa Moving Group)
ψ	UMa	11 10.9 +44 23	3.00	1.14	K1 III	22.6	-0.2	145	0.068	246	-4 V?		slow rotator (but ≤ 2.6 y)
δ	Leo A	11 15.2 +20 24	2.56 [†]	0.13	A4 IV	56	1.3	58	0.193	132	-20 V		Zosma
													rapid rotator (< 0.5 d)
θ	Leo	11 15.4 +15 19	3.33	0.00	A2 IV [†]	~19.8	-0.2	165	0.099	217	+8 V		¶ suspected δ Sct variable
													Chertan
													rotation rather slow for MK type A (but < 9 d); quiet atmosphere renders Ca, Sc underabundant, and Fe, Sr, Ba overabundant; Ca II K-line is variable
v	UMa A	11 19.6 +32 59	3.49	1.40	K3 III Ba0.3 [†]	~8.2	-1.9	400	0.039	317	-9 SB		¶ IR excess (debris disk?) B: 10.1, 7.5", PA: 145°→149°, 1827→2018 Alula Borealis orbit ≥ 12,000 y; separation ≥ 950 AU
ξ	Hya Aa	11 34.1 -31 59	3.54	0.95	G7 III	~25.2	0.5	130	0.214	259	-5 V		¶ Astron. Alm. (epoch 2021.5) assigns MK type K3- III [THIS STAR ONLY IN ONLINE VERSION OF TABLE]
													La Silla CORALIE ~2001 detected multimodal oscillations, not all radial, with periods ~3 h [THIS STAR ONLY IN ONLINE VERSION OF TABLE]
λ	Cen Aa [†]	11 36.8 -63 08	3.11	-0.04	B9.5 IIIn [†]	8	-2.4	400	0.034	258	-1 V		despite possible fast rotation (< 2.7 d?), Fe is overabundant, with Si and C mildly underabundant
													¶ at http://stars.astro.illinois.edu/sow/lambda_cen.html , Kaler discusses questions of visual binarity (λ Cen Aa, Ab, B)
β	Leo A	11 50.2 +14 27	2.14 [†]	0.09	A3 Va	91	1.9	36	0.511	257	0 V		Denebola
													rapid rotator (< 0.65 d)
													¶ debris disk resolved by <i>Herschel Space Observatory</i> (2010A&A...518L135M), disk structures differentiated with ground-based interferometry (2010ApJ...724.1238S)
γ	UMa A	11 55.0 +53 35	2.41	0.04	A0 Van [†]	39	0.4	83	0.108	84	-13 SB		¶ assertion of δ Sct variability now seems erroneous
													Phecda
													rapid rotator: although in MK temperature class A, nevertheless an instance of the "Be phenomenon" (the term "Ae star" is sometimes used for this rare class)
δ	Cen Aa [†]	12 09.5 -50 51	2.58 ^v	-0.13	B2 IVne [†]	8	-2.9	400	0.050	262	+11 V		¶ E(B-V)=0.00 var. in γ Cas class: 2.51-2.65 in V
													¶ rapid rotator (< 1.3 d), with shell spectrum; 2008A&A...488L...67M summarizes recent research, and as part of a wider VLT investigation into the "Be phenomenon" not only discusses the circumstellar ejecta, but also reports discovery of binarity (Ab at angular distance 68.7 mas)
ε	Crv	12 11.2 -22 44	3.02	1.33	K2 III [†]	~10.3	-1.9	320	0.072	278	+5		slow rotator (but ≤ 3.9 y)
													¶ metals somewhat overabundant
													¶ evolutionary status uncertain (core helium fusion starting, in progress, or finished?)
													¶ Astron. Alm. (epoch 2021.5) assigns MK type K2.5 IIIa
													¶ the etymologically Arab name "Minkar" is of merely modern origin, and is not currently IAU-official variable. in β Cep class: 2.78-2.84 in V, 0.15 d
δ	Cru	12 16.3 -58 52	2.79 ^v	-0.19	B2 IV [†]	9.4	-2.3	350	0.037	254	+22 V?		¶ rapid rotator (< 1.3 d; BSC5: "expanding circumstellar shell")
													Imai
δ	UMa A	12 16.5 +56 55	3.32	0.08	A2 Van	40.5	1.4	81	0.104	86	-13 V		Megrez
													possesses debris disk, of unusually low radius (Wyatt et al 2007; Pointing-Robertson drag?)
γ	Crv	12 16.9 -17 40	2.58	-0.11	B8 III [†]	21	-0.8	154	0.160	278	-4 SB		spectral variable?
													Gienah
													¶ rather rapid rotation notwithstanding (BSC5: "expanding circumstellar shell"), Hg and Mn are overabundant, with some other elements underabundant (but rotational line broadening makes abundance determinations difficult); Astron. Alm. (epoch 2021.5) assigns MK temperature type B8p Hg Mn, and does not assign an MK luminosity class
α	Cru A [†]	12 27.8 -63 13	1.25 [†]	-0.20	B0.5 IV	10	-3.7	~320	0.037	251	-11 SB		5.4" (1826); 3.5" (2020)
α	Cru B [†]	12 27.8 -63 13	1.64 [†]	-0.18	B1 Vn	10	-3.3	~320	0.037?	251?	-1		Acrux
													PA: 114°→111°, 1826→2020 orbit ≥ 1300 y, separation ~430 AU;
													A is SB pair Aa, Ab (75.78 d, separation ~0.5 AU min, ~1.5 AU max);
													C (itself an SB pair) at ~90" from AB, imperfectly sharing the AB proper motion, is possibly (not assuredly) gravitationally bound with AB (if bound, then > 130,000 y, with separation ≥ 9,000 AU);
													WDS additionally documents D, E, F, G, H, I, J, K
													¶ duplicity makes individual mag. determinations for A, B somewhat controverted
δ	Crv A [†]	12 31.0 -16 38	2.94	-0.01	B9.5 IVn	~37.6	0.8	87	0.252	237	+9 V		B: 8.26, K2 V, 24.2", PA: 216°→216°, 1782→2020 Algorab orbit ≥ 9400 y; although A, B have common proper

γ	Cru A	12 32.4 –57 14	1.59v [†] 1.60	M3.5 III [†]	37	–0.6 89	0.267 174	+21	<p>motion, disparity in age estimates has caused binarity to be questioned</p> <p>¶ Kaler suggests B is young post-T-Tauri star, with surrounding dust as yet uncleared</p> <p>Gacrux</p> <p>semiregular variable: 1.60–1.67 in V</p> <p>although has been classified as semiregular var., at least 6 pulsation periods have been documented</p> <p>¶ the nearest of the M giants, radius > 0.5 AU; evolutionary status uncertain (is core He fusion now finished?)</p> <p>¶ cause of the observed Ba overabundance is unknown (undetected evolved companion?)</p>
β	Crv	12 35.5 –23 31	2.65 [†] 0.89	G5 II [†]	22	–0.6 146	0.057 179	–8 V	<p>Kraz</p> <p>slight variability has been reported (2.60–2.66 in V)</p> <p>¶ slow rotator (but \leq 180 d)</p> <p>¶ possibly in evolutionary transition (He core about to ignite?)</p> <p>¶ assertion of weak Ba-star status is perhaps erroneous</p> <p>¶ Astron. Alm. (epoch 2021.5) assigns MK type G5 IIB</p> <p>variable. in β Cep class: 2.68–2.73 in V, 0.090 d</p> <p>classification of the low-amplitude variability as β Cep has been questioned</p> <p>¶ rapid rotator (< 2 d)</p> <p>orbit 84 y; 0.4" (2010), 0.3" (2019); max = 1.7";</p> <p>separation 8 AU min, 67 AU max, 37 AU average</p> <p>¶ Arabic name Muhlifain is not currently IAU-official</p> <p>A: 3.48; B: 3.53; 0.8" (2007); 3.0" (2020)</p> <p>Porrima</p> <p>orbit 169 y;</p> <p>separation 5 AU min (most recently 1836 and 2005), 81 AU max, 43 AU average, with plane of orbit inclined 31° to plane of sky; for discussion of orbit, with observations plot showing error bars (binary astrometry being now old enough to archive data for one full orbit), cf Kaler at http://stars.astro.illinois.edu/sow/porrima.html</p> <p>¶ lunar occultations possible, planetary occultations possible-yet-rare</p> <p>A: 3.52; B: 3.98, 1.0", PA:317°→56°, 1880→2019 orbit 194 y; average separation uncertain (101 AU, or only ~80 AU?); orbit map, showing error bars, given by Kaler at http://stars.astro.illinois.edu/sow/betamus.html, with Kaler's accompanying discussion of orbit-modelling problems, underscores limitations in current β Mus AB knowledge</p> <p>¶ β Mus A is rapid rotator (< 1 d)</p> <p>¶ β Mus B is of MK type B2.5 V</p> <p>¶ a runaway system, in the sense of presenting a high velocity relative to the general galactic rotation</p>
α	Mus Aa	12 38.5 –69 15	2.69 [†] –0.18	B2 IV–V	10.3	–2.2 320	0.042 252	+13 V	<p>variable in β Cep class: 2.68–2.73 in V, 0.090 d</p> <p>classification of the low-amplitude variability as β Cep has been questioned</p> <p>¶ rapid rotator (< 2 d)</p> <p>orbit 84 y; 0.4" (2010), 0.3" (2019); max = 1.7";</p> <p>separation 8 AU min, 67 AU max, 37 AU average</p> <p>¶ Arabic name Muhlifain is not currently IAU-official</p> <p>A: 3.48; B: 3.53; 0.8" (2007); 3.0" (2020)</p> <p>Porrima</p> <p>orbit 169 y;</p> <p>separation 5 AU min (most recently 1836 and 2005), 81 AU max, 43 AU average, with plane of orbit inclined 31° to plane of sky; for discussion of orbit, with observations plot showing error bars (binary astrometry being now old enough to archive data for one full orbit), cf Kaler at http://stars.astro.illinois.edu/sow/porrima.html</p> <p>¶ lunar occultations possible, planetary occultations possible-yet-rare</p> <p>A: 3.52; B: 3.98, 1.0", PA:317°→56°, 1880→2019 orbit 194 y; average separation uncertain (101 AU, or only ~80 AU?); orbit map, showing error bars, given by Kaler at http://stars.astro.illinois.edu/sow/betamus.html, with Kaler's accompanying discussion of orbit-modelling problems, underscores limitations in current β Mus AB knowledge</p> <p>¶ β Mus A is rapid rotator (< 1 d)</p> <p>¶ β Mus B is of MK type B2.5 V</p> <p>¶ a runaway system, in the sense of presenting a high velocity relative to the general galactic rotation</p>
γ	Cen A [†]	12 42.7 –49 05	2.95 –0.02	A1 IV	25	–0.1 130	–0.194 –267	–6	<p>orbit 84 y; 0.4" (2010), 0.3" (2019); max = 1.7";</p> <p>separation 8 AU min, 67 AU max, 37 AU average</p> <p>¶ Arabic name Muhlifain is not currently IAU-official</p> <p>A: 3.48; B: 3.53; 0.8" (2007); 3.0" (2020)</p> <p>Porrima</p> <p>orbit 169 y;</p> <p>separation 5 AU min (most recently 1836 and 2005), 81 AU max, 43 AU average, with plane of orbit inclined 31° to plane of sky; for discussion of orbit, with observations plot showing error bars (binary astrometry being now old enough to archive data for one full orbit), cf Kaler at http://stars.astro.illinois.edu/sow/porrima.html</p> <p>¶ lunar occultations possible, planetary occultations possible-yet-rare</p> <p>A: 3.52; B: 3.98, 1.0", PA:317°→56°, 1880→2019 orbit 194 y; average separation uncertain (101 AU, or only ~80 AU?); orbit map, showing error bars, given by Kaler at http://stars.astro.illinois.edu/sow/betamus.html, with Kaler's accompanying discussion of orbit-modelling problems, underscores limitations in current β Mus AB knowledge</p> <p>¶ β Mus A is rapid rotator (< 1 d)</p> <p>¶ β Mus B is of MK type B2.5 V</p> <p>¶ a runaway system, in the sense of presenting a high velocity relative to the general galactic rotation</p>
γ	Cen B [†]	12 42.7 –49 05	2.85 –0.02	A0 IV	25	–0.2 130	–0.194 –267	–6	<p>orbit 84 y; 0.4" (2010), 0.3" (2019); max = 1.7";</p> <p>separation 8 AU min, 67 AU max, 37 AU average</p> <p>¶ Arabic name Muhlifain is not currently IAU-official</p> <p>A: 3.48; B: 3.53; 0.8" (2007); 3.0" (2020)</p> <p>Porrima</p> <p>orbit 169 y;</p> <p>separation 5 AU min (most recently 1836 and 2005), 81 AU max, 43 AU average, with plane of orbit inclined 31° to plane of sky; for discussion of orbit, with observations plot showing error bars (binary astrometry being now old enough to archive data for one full orbit), cf Kaler at http://stars.astro.illinois.edu/sow/porrima.html</p> <p>¶ lunar occultations possible, planetary occultations possible-yet-rare</p> <p>A: 3.52; B: 3.98, 1.0", PA:317°→56°, 1880→2019 orbit 194 y; average separation uncertain (101 AU, or only ~80 AU?); orbit map, showing error bars, given by Kaler at http://stars.astro.illinois.edu/sow/betamus.html, with Kaler's accompanying discussion of orbit-modelling problems, underscores limitations in current β Mus AB knowledge</p> <p>¶ β Mus A is rapid rotator (< 1 d)</p> <p>¶ β Mus B is of MK type B2.5 V</p> <p>¶ a runaway system, in the sense of presenting a high velocity relative to the general galactic rotation</p>
γ	Vir AB [†]	12 42.8 –1 34	2.74 0.37	F1 V + F0mF2 V	85	2.4 39	–0.619 –276	–20	<p>orbit 169 y;</p> <p>separation 5 AU min (most recently 1836 and 2005), 81 AU max, 43 AU average, with plane of orbit inclined 31° to plane of sky; for discussion of orbit, with observations plot showing error bars (binary astrometry being now old enough to archive data for one full orbit), cf Kaler at http://stars.astro.illinois.edu/sow/porrima.html</p> <p>¶ lunar occultations possible, planetary occultations possible-yet-rare</p> <p>A: 3.52; B: 3.98, 1.0", PA:317°→56°, 1880→2019 orbit 194 y; average separation uncertain (101 AU, or only ~80 AU?); orbit map, showing error bars, given by Kaler at http://stars.astro.illinois.edu/sow/betamus.html, with Kaler's accompanying discussion of orbit-modelling problems, underscores limitations in current β Mus AB knowledge</p> <p>¶ β Mus A is rapid rotator (< 1 d)</p> <p>¶ β Mus B is of MK type B2.5 V</p> <p>¶ a runaway system, in the sense of presenting a high velocity relative to the general galactic rotation</p>
β	Mus Aa	12 47.6 –68 14	3.04 –0.18	B2 V [†]	~9.6	–2.1 340	–0.043 [†] ~258	+42 [†] V	<p>variable in β Cep class: 2.68–2.73 in V, 0.090 d</p> <p>classification of the low-amplitude variability as β Cep has been questioned</p> <p>¶ rapid rotator (< 2 d)</p> <p>orbit 84 y; 0.4" (2010), 0.3" (2019); max = 1.7";</p> <p>separation 8 AU min, 67 AU max, 37 AU average</p> <p>¶ Arabic name Muhlifain is not currently IAU-official</p> <p>A: 3.48; B: 3.53; 0.8" (2007); 3.0" (2020)</p> <p>Porrima</p> <p>orbit 169 y;</p> <p>separation 5 AU min (most recently 1836 and 2005), 81 AU max, 43 AU average, with plane of orbit inclined 31° to plane of sky; for discussion of orbit, with observations plot showing error bars (binary astrometry being now old enough to archive data for one full orbit), cf Kaler at http://stars.astro.illinois.edu/sow/porrima.html</p> <p>¶ lunar occultations possible, planetary occultations possible-yet-rare</p> <p>A: 3.52; B: 3.98, 1.0", PA:317°→56°, 1880→2019 orbit 194 y; average separation uncertain (101 AU, or only ~80 AU?); orbit map, showing error bars, given by Kaler at http://stars.astro.illinois.edu/sow/betamus.html, with Kaler's accompanying discussion of orbit-modelling problems, underscores limitations in current β Mus AB knowledge</p> <p>¶ β Mus A is rapid rotator (< 1 d)</p> <p>¶ β Mus B is of MK type B2.5 V</p> <p>¶ a runaway system, in the sense of presenting a high velocity relative to the general galactic rotation</p>
β	Cru A	12 49.0 –59 48	1.25v [†] –0.24	B0.5 III [†]	12	–3.4 300	0.046 249	+16 SB [†]	<p>variable in β Cep class: 1.23–1.31 in V, 0.24 d</p> <p>SB period 5 y, separation 5.4 AU min, 12.0 AU max; Kaler at http://stars.astro.illinois.edu/sow/mimosa.html</p> <p>discusses other possible companions, including an X-ray visible, and yet optically invisible, object interpreted as a pre-MS star</p> <p>¶ β Cru A is believed to be a rapid rotator (possible ~3.6 d)</p> <p>¶ β Cru A is a multiperiodic β Cep variable</p> <p>¶ β Cru A, its MK luminosity class "III" notwithstanding, is only about halfway through its career of stable-core hydrogen fusing</p> <p>variable in α^2 CVn class: 1.76–1.78 in V, 5.1 d</p> <p>Alioth</p> <p>the brightest of the Ap stars</p> <p>(in the specific case of ϵ UMa A, the magnetic-dipole axis is believed to be nearly perpendicular to rotation axis, yielding Cr bands nearly perpendicular to equator; dipole strength is unusually low) (but it has also been suggested that a substellar companion of mass ~14.7× Jupiter, at average separation 0.055 AU, orbit 5.1 d, rather than a 5.1-d stellar rotation, is the source of the observed variability period)</p>
ϵ	UMa A	12 55.0 +55 51	1.76v [†] –0.02	A0p IV: (CrEu)	~39.5	–0.3 83	0.112 94	–9 SB?	<p>variable in α^2 CVn class: 1.76–1.78 in V, 5.1 d</p> <p>Alioth</p> <p>the brightest of the Ap stars</p> <p>(in the specific case of ϵ UMa A, the magnetic-dipole axis is believed to be nearly perpendicular to rotation axis, yielding Cr bands nearly perpendicular to equator; dipole strength is unusually low) (but it has also been suggested that a substellar companion of mass ~14.7× Jupiter, at average separation 0.055 AU, orbit 5.1 d, rather than a 5.1-d stellar rotation, is the source of the observed variability period)</p>
δ	Vir A	12 56.7 +3 17	3.39 [†] 1.57	M3 III [†]	16	–0.5 ~198	0.473 [†] 264	–18 [†] V?	<p>Minelauva</p> <p>semireg. var. (multiperiod pulsator), mag. 3.32–3.40 in V</p> <p>¶ high space velocity relative to galactic neighbours</p> <p>¶ evolutionary status uncertain (helium fusion recently started, or already finished?)</p> <p>¶ Astron. Alm. (epoch 2021.5) assigns MK type M3⁺ III</p> <p>B:5.5, F0 V, 19.5", PA:234°→230°, 1777→2018</p> <p>Cor Caroli</p> <p>orbit \geq 8300 y (common proper motion indicates true binarity); separation \geq 675 AU; prototype for the α^2 CVn var. class (chemically anomalous photospheric regions yielding spectroscopic variability, and with</p>
α	CVn A [†]	12 57.0 +38 12	2.85v [†] –0.06	A0 Vp (SiEu)	28	0.1 110	0.241 283	–3 V	<p>variable in α^2 CVn class: 1.76–1.78 in V, 5.1 d</p> <p>Alioth</p> <p>the brightest of the Ap stars</p> <p>(in the specific case of ϵ UMa A, the magnetic-dipole axis is believed to be nearly perpendicular to rotation axis, yielding Cr bands nearly perpendicular to equator; dipole strength is unusually low) (but it has also been suggested that a substellar companion of mass ~14.7× Jupiter, at average separation 0.055 AU, orbit 5.1 d, rather than a 5.1-d stellar rotation, is the source of the observed variability period)</p>

ε	Vir A	13 03.2 +10 51	2.85	0.93	G9 IIIab [†]	29.8	0.2	110	0.275	274	-14 V?	<p>magnetism yielding large spots; in the particular case of α^2 CVn, rotation period is 5.46939 d, with consequent spot-driven mag. range 2.84–2.98)</p> <p>¶ two correct, potentially confusing, designations are α CVn A (signalling that this is the brighter of the binary pair) and α^2 CVn (signalling that α^1 crosses the local meridian before α^2, lying further W); the Latin “heart-of-Charles” designation, official at IAU as of 2016, honours the “martyr king” Charles I (although Charles II is sometimes cited in error)</p> <p>Vindemiatrix</p> <p>one of the most notable X-ray sources in our table (X-ray luminosity, although far below α Aur, is nevertheless almost 300× solar)</p> <p>¶ Astron. Alm. (epoch 2021.5) assigns MK type G8 IIIab</p>
γ	Hya A	13 20.1 -23 17	2.99	0.92	G8 IIIa	-24.4	-0.1	134	0.081	121	-5 V?	<p>slow rotator (but ≤ 240 d)</p> <p>evolutionary state uncertain (core helium fusion impending, or already in progress?)</p>
ι	Cen	13 21.8 -36 50	2.75	0.07	A2 Va [†]	55	1.5	59	0.352	256	0	<p>rapid rotator (< 2d)</p> <p>¶ low metallicity</p> <p>¶ debris disk (unusually luminous, given evolutionary state of ι Cen)</p>
ζ	UMa Aa [†]	13 24.8 +54 49	2.23	0.06	A1 Va [†]	40	0.1	90	0.123	100	-6 SB2 [†]	<p>B:3.88, A1mA7 IV–V, 14.6"; period >5000 y? Mizar</p> <p>not only are A+B a true binary; it is now additionally argued (controversy possibly continues) that Alcor is gravitationally bound to A+B (Bob King, <i>Sky & Telescope</i> 2015 Mar. 25); although ζ UMa A and ζ UMa B (both chemically anomalous) are universally accepted as themselves individually SB (yielding quartet ζ UMa Aa, Ab, 20.538 d (cf NPOI trial, 1997), ζ UMa Ba, Bb, 175.6 d; both SB orbits are highly elliptical), the old, widely repeated claim that Alcor is itself SB requires scrutiny (pro, F. Heard <i>ApJ</i> 1949; contra, www.leosondra.cz/en/mizar, specifically rebutting Heard; once again pro, but now on new basis (discovering elusive red-dwarf companion), 2010ApJ...709..733Z;</p> <p>this Leos Ondra web source should be consulted also (a) for details on Mizar-Alcor multiplicity-studies history, including Galileo and Michelson (Ondra, citing inter alia Fedele 1949, seems to establish that it was Galileo pupil Castelli, rather than (as widely asserted) Riccioli, who discovered Mizar’s visual duplicity) and (b) for a 15' map documenting around 20 of the field stars, including mag. 7.58 “Stella Luoviciana” (“Sidus Ludovicianum”))</p> <p>¶ Astron. Alm. (epoch 2021.5) assigns MK type “A1 Va+ (Si)”</p>
α	Vir Aa [†]	13 26.3 -11 16	0.98v [†]	-0.24	B1 V [†]	13	-3.4	250	0.052	234	+1 SB2 [†]	<p>variable: 0.96–1.00 in V, 4.0 d; mags 3.1, 4.5, 7.5 Spica</p> <p>¶ this SB (separation 0.12 AU, 4.0145 d; the geometry is close to achieving a grazing eclipse) is the brightest of the rotating ellipsoid variables (by definition no eclipse, and by definition with the SB’s total presented luminous area varying, through geometrical asymmetry, as the orbital motion proceeds); the Aa,Ab orbit is highly eccentric; Aa (a rapid rotator, at ~0.3 breakup speed) is itself a pulsating variable of the β Cep type (0.1738 d; shortly after the ~1970 discovery of the β Cep variability, photometric and spectroscopic variations were present; the photospheric variations soon ceased, but the spectroscopic (radial-velocity, i.e. pulsational) variations continued; 2016MNRAS.458.1964T, incorporating precision <i>MOST</i> photometry, reports for Aa one radial and two non-radial pulsation modes, with one of the non-radial modes tidally induced)</p> <p>¶ in an early application of interferometry, 1971MNRAS.151..161H argues with the example of α Vir Aa,Ab that given supporting spectroscopy and photometry, orbit and distance of a double-lined SB can be deduced (the SB distance notably</p>

													without recourse to parallax measurements and without recourse to spectroscopic determination of luminosity classes) ¶ the tidal-interactions studies 2016A&A...590A..54H and 2013A&A...556A..49P stress the importance of the α Vir Aa,Ab double-lined SB for critically testing the (astrophysically foundational) assumption that the individual components x , y of a binary, of determined masses, rotation periods, and chemical compositions, resemble in their photospheres, and even in their interiors, solitary stars x' , y' possessing the same masses, rotation periods, and chemical compositions (could tidal effects, e.g. change internal temperature structure?); additionally, the tidal effects in the α Vir Aa,Ab SB are judged in 2009ApJ...704..813H to be responsible for large-scale shearing horizontal photospheric motions, spectroscopically observable as modifiers of line profiles (but 2016MNRAS.458.1964T questions the judgement) ¶ assignment of individual MK types to Aa, Ab is challenging: the rather-unevolved-B MK types (1971MNRAS.15L..161H B1.5 IV-V + B3V, 2007AAS...211.6301A B0.5 III-IV + B2.5-B3V) are in any case consistent with rather high masses (10.9 M_{\odot} + 6.8 M_{\odot} , 10.25 M_{\odot} + 6.97 M_{\odot} , for these two respective papers) ¶ as is to be expected from the failure of Aa,Ab to be tidally locked, the system is young (with 2016MNRAS.458.1964T assigning as age 12.5 ± 1 My) ¶ the Aa,Ab binary is a polarimetric variable (ISM material entrained?), and a strong X-ray source (colliding winds?) ¶ α Vir Ab is one of the few stars known to exhibit Struve-Sahade variation (https://en.wikipedia.org/wiki/Struve%E2%80%93Sahade_effect) in its spectral line strengths ¶ 1972JBA...82..431K describes the 18.6-year 1940-through-2050 cycle of lunar occultation possibilities ¶ Aa was measured in 1975 to lie 0.50" from Ac ; although the Aa,Ab pair is at all times very close, an angular distance of 0. 1" is reported from 1975 ¶ $E(B-V)=+0.03$ a good marker of celestial equator (precession placed ζ Vir exactly onto equator in February 1883) ¶ rapid rotator (< 0.5 d; this renders puzzling the possible evidence for chemical anomalies, which would presuppose a quiet atmosphere) ¶ Astron. Alm. (epoch 2021.5) assigns MK type A2 IV- ¶ elusive red-dwarf companion ζ Vir B is reported in 2010ApJ...712..421H (0.168 M_{\odot} , possibly accounting for the X-ray emission observed by <i>ROSAT</i> : as a star of a spectral type lacking strong winds and lacking convection at photosphere, ζ Vir A would not itself be expected to emit X-rays)
ζ	Vir A	13 35.8	-0.42	3.38	0.11	A2 IV [†]	44	1.6	74	0.285	280	-13	Heze
ε	Cen Aa	13 41.3	-53 34	2.29 [†]	-0.17	B1 III [†]	8	-3.3	400	0.019	233	+3	slight variability (mag. 2.29–2.31 in V; multiperiodic; in β Cep class) ¶ rapid rotator (< 2.7 d) ¶ metals underabundant ¶ although we here assign MK luminosity class "III", Kaler at http://stars.astro.illinois.edu/sow/epsce.html discusses uncertainty
η	UMa	13 48.4	+49 12	1.85	-0.10 [†]	B3 V [†]	31	-0.7	104	0.122	263	-11 SB?	Alkaid resembles α UMa, at the other extreme of the Big Dipper, in not belonging to UMa Moving Group; 1921LicOB..10..110T asserts

													membership in what was at that time called the “Pleiades Group” ¶ rapid rotator (< 21 h), with some line variability (circumstellar ejecta disk?) ¶ X-ray source ¶ colour and mean temperature are anomalous for the MK type ¶ unusually young in our Sample S (< 15 My) ¶ E(B–V)=+0.02
v	Cen	13 50.8 –41 48	3.41 [†] –0.22	B2 IV [†]	~7.5 –2.2 440	0.034	233	+9 SB [†]					SB period is 2.622 d; system is rotating ellipsoidal variable (not eclipsing, but varying in light as the presented surface area changes); additionally, the primary is a pulsator in the β Cep class (mag. 3.40–3.42 in V) ¶ MK luminosity class “IV” notwithstanding, primary is still a stable fuser of core hydrogen ¶ possible weak instance of “Be phenomenon” (with the outbursts possibly temporary)
μ	Cen Aa	13 50.9 –42 35	3.47v [†] –0.17	B2 IV–V pne [†]	~6.4 –2.5 510	0.031	232	+9 SB					variable in γ Cas class: 2.92–3.47 in V rapid rotator, and (consistently with the γ Cas behaviour) an instance of the “Be phenomenon”; additionally said to be a multiperiodic non-radial pulsator; BSC5: “line profiles of MgII 4481 change in period 0.505 d, about five times the period of weaker absorption”; variable Hα; “variable line profiles”; short-term photometric and polarimetric variability has also been reported (cf p. 46 of 2013A&ARv..21...69R , which notes a rapid rise, over just a few days, in photometric brightness or line-emission intensity, with a subsequent slower decline) ¶ Astron. Alm. (epoch 2021.5) assigns MK type “B2 IV–Vpne (shell)”
η	Boo A	13 55.7 +18 17	2.68 0.58	G0 IV [†]	88 2.4 37	0.361	190	0 SB					Muphrid unusually metal-rich ¶ an X-ray source (hot corona) ¶ 2007ApJ...657.1058V discusses recent work (<i>MOST</i> helioseismology, PTI interferometry)
ζ	Cen	13 56.9 –47 24	2.55 –0.18	B2.5 IV [†]	8.5 –2.8 380	0.073	232	+7 SB2					SB period 8.02 d; SB as yet unresolved, even by interferometry (so WDS not yet able to write “Cen A,” “Cen B”) ¶ primary is a rapid rotator (<1.5 d) (BSC5: “expanding circumstellar disk”, and yet not presently (Jan. 2021) catalogued as an instance of the “Be phenomenon” in Paris-Meudon BeSS database) ¶ MK luminosity class “IV” notwithstanding, primary is possibly only halfway through its core hydrogen fusing ¶ E(B–V)=–0.02
β	Cen Aa,Ab	14 05.4 –60 29	0.58v [†] –0.23	B1 III + B1 III	8 [†] –4.8 400 [†]	0.041	235	+6 SB [†]					Hadar B:3.94, A1mA7 IV–V, 0.3” (2019) Aa, Ab are of mags. ~1.3, ~1.4, respectively; IAU name “Hadar” designates Aa, not the Aa+Ab SB; entire (triple) system comprises SB Aa+Ab (357 d, separation 0.53 AU min, 5.5 AU max, 4 AU average) with B; at least one, and perhaps both, of Aa, Ab are multiperiod variables in the β Cep class; Kaler discusses uncertainty in distance (etc.) of the triple at http://stars.astro.illinois.edu/sow/hadar.html ¶ E(B–V)=+0.02
π	Hya	14 07.6 –26 47	3.25 1.09	K2 IIIb [†]	~32.3 [†] 0.8 ~101 [†]	0.148 [†]	163	+27 [†] V					negative cyanide ion lines are anomalously weak relative to metal lines, consistent with this star’s anomalously high velocity relative to Sun (suggesting interloper in our own galactic region; however, π Hya is more metal-rich than the celebrated interloper α Boo (Arcturus)) ¶ Astron. Alm. (epoch 2021.5) assigns MK type K2– III Fe–0.5 ¶ in evolutionary terms, in “Red Clump” of core-He fusers (but uncertain whether recent arrival in clump or longtime denizen)
θ	Cen A	14 08.0 –36 28	2.06 1.01	K0 IIIb [†]	55 0.8 59	0.734 [†]	225	+1 [†]					Menkent high velocity with respect to Sun suggests interloper status (and yet metallicity is approximately solar) ¶ Astron. Alm. (epoch 2021.5) assigns MK type K0– IIIb
α	Boo A	14 16.6 +19 04	–0.05 1.24	K1.5 III Fe–0.5 [†]	89 –0.3 37	2.279 [†]	209	–5 [†] V					high space velocity a metal-poor interloper (from galactic thick disk? but galaxy-merger scenario has also been suggested), and member of Arcturus Moving Group (2009IAUS..254..139W)

												<p>¶ a magnetic cycle (< 14 y?) has been detected</p> <p>¶ still ascending RGB, with He flash impending? (but a later evolutionary stage has also been suggested)</p> <p>¶ publication of α Boo A line atlas 1968pmas.book.....G (R.Griffin) was a major event in postwar spectroscopy</p> <p>¶ α Boo A has been studied in recent astroseismology</p> <p>¶ there may be a companion, at margin of <i>HIPPARCOS</i> detectability; and perhaps separately from this, WDS reports a 1991 observation of “α Boo B”, 0.30”, at mag. 3.49; we are therefore constrained to write here “α Boo A” rather than “α Boo,” while stressing that presence of celestial-sphere neighbour is currently far from confirmed)</p>
ι	Lup	14 20.8 -46 09	3.55 -0.18	B2.5 IVn [†]	~9.6 -1.5 340	0.013	249	+22				<p>rapid rotator (possibly ~0.9 d), and yet no evidence of circumstellar disk, and in particular no Be-phenomenon spectral features</p> <p>¶ the MK luminosity class “IV” notwithstanding, still performing stable core-hydrogen fusion</p> <p>[THIS STAR ONLY IN ONLINE VERSION OF TABLE]</p>
γ	Boo Aa [†]	14 32.9 +38 13	3.04v [†] 0.19	A7 IV ⁺	37.6 0.9 87	0.190	323	-37 V				<p>variable in the δ Sct class</p> <p>¶ IR excess (from circumstellar debris, so far unexplained)</p> <p>¶ Aa,Ab resolved in speckle Interferometry, angular distance 70 mas</p> <p>γ Cas and λ Eri var.: 2.29–2.47 in V, multiperiodic BSC5: Hα variable, Hβ “sometimes bright, sometimes dark and double or multiple”; consistently with γ Cas variability, a rapid rotator (< 1 d) and an instance of “Be phenomenon”;</p> <p>Astron. Alm. (epoch 2021.5) assigns MK type “B1.5 IVpne (shell)”</p>
η	Cen	14 36.9 -42 15	2.33v [†] -0.16	B1.5 IV pne [†]	11 -2.5 310	0.048	227	0 SB				<p>AB 4.4” (2017) orbit 79.9y</p> <p>min = 2” (1955); max = 22”; PA (2017) = 325°; separation 11.2 AU min, 35.6 AU max; Kaler at http://stars.astro.illinois.edu/sow/rigil-kent.html has map of AB orbit (note further here that Kaler’s green, violet, blue denote micrometry, photography, interferometry, respectively: as the error bars suggest, the AB orbit is one of the most precisely known in visual-binary astrometry; Kaler also discusses some uncertainties in the ABC physical properties)</p> <p>¶ magnetic activity of α Cen A is in deep decline since 2005</p> <p>¶ 2005A&A...442...315R reports a flare on magnetically active α Cen B</p> <p>¶ 2012 B-exoplanet claim now discounted, and yet an exoplanet Bc now considered possible in 2015MNRAS.450.2043D</p> <p>¶ Einstein-ring event expected with 45% probability in 2028, early in May</p>
α	Cen B [†]	14 41.1 -60 55	1.35 0.9	K1 V [†]	750 5.7 4.3	~3.703	~283	-21 V?				<p>Toliman</p>
α	Cen A [†]	14 41.1 -60 55	-0.01 0.71	G2 V [†]	750 4.4 4.3	~3.710	~277	-22 SB				<p>Ca (Proxima), 12.4, M5e, 212°, Cb (exopl) Rigil Kentaurus gravitational binding of AB+C was finally established with high probability in 2017A&A...598L...7K (~550,000 y: min > 4300 AU, max 13,000 AU)</p> <p>¶ 2016Natur.536.437A announces an approx. Earth-mass exoplanet (mass < ~3× Earth?) around α Cen C (unfortunately, however, for exobiology, α Cen C suffers superflares);</p> <p>https://en.wikipedia.org/wiki/Proxima_Centauri_b discusses astrobiology pros and cons;</p> <p>http://breakthroughinitiatives.org/initiative/3 advocates nanocraft exploration</p> <p>¶ all of A, B, C are metals-rich</p>
α	Lup A	14 43.4 -47 29	2.30 [†] -0.15	B1.5 III	7 -3.5 460	0.032	221	+5 SB				<p>variable in β Cep class: 2.29–2.34 in V, 0.26 d actually multiperiodic, with primary period (unusually long) 0.2598466 d given by AAVSO(VSX) as viewed 2021 Jan. 16</p>
α	Cir A [†]	14 44.3 -65 04	3.18 [†] 0.26	A7 Vp (Sr)	60.4 2.1 54.1	0.303	220	+7 SB?				<p>B: 8.6, K5 V, 15.7”, PA:263°→224°, 1826→2016</p> <p>AB probably true binary, with orbit \geq 2600 y</p> <p>¶ Astron. Alm. (epoch 2021.5) assigns MK temperature type A7p Sr Eu and does not assign an MK luminosity class</p> <p>¶ the brightest variable of the AAVSO “rapidly oscillating Ap” class (features of the type include rapid non-radial pulsation with a stellar-rotation signal), with V mag. range 3.17–3.19; magnetically an oblique rotator (4.4790 d, with field strength ~500× solar);</p> <p>2009MNRAS.396.1189B discusses the rotation, two notably stable putative equatorial chemical-</p>

ϵ	Boo A	14 45.9 +26 59	2.58	1.34	K0 II–III [†]	16 [†]	-1.6	200 [†]	0.044	288	-17 V	<p>anomaly regions, and astroseismology, with history and fresh <i>WRE</i>+SAAO observations</p> <p>B:4.81, 2.8", PA:318°→347°, 1822→2018 orbit well over 1000 y</p> <p>¶ ϵ Boo B is a rapid rotator</p> <p>¶ ϵ Boo B is of MK type A0 V</p> <p>¶ http://stars.astro.illinois.edu/sow/izar.html discusses difficulties in determination of the individual magnitudes and of the binary system's distance</p> <p>¶ F.G.W. von Struve: "pulcherrima" ("the loveliest")</p> <p>Izar</p>
β	UMi A+1P [†]	14 50.7 +74 04	2.07	1.46	K4 III [†]	24.9	-0.9	131	0.035	289	+17 V	<p>useful for aligning small equatorial mount</p> <p>(since NCP, although not quite coincident with α UMi, does lie near the great-circle arc linking β UMi with α UMi:</p> <p>http://arksky.org/Kochab.htm)</p> <p>¶ Fe underabundant, Ba possibly slightly overabundant</p> <p>¶ 2008A&A...483L..43T suggests (via <i>CORIOLIS</i>-SMEI) two short-lived radial-pulsation mods</p> <p>¶ 2014A&A...566A...67L announces exoplanet</p> <p>Kochab</p>
α	Lib Aa [†]	14 52.1 -16 08	2.75	0.15	A3 III–IV [†]	43	0.9	76	0.126	237	-10 SB [†]	<p>Zubenelgenubi</p> <p>angular distance from α Lib B, which shares the proper motion of α Lib A entails separation ≥ 5500 AU; if B and A are gravitationally bound, then their period is $\geq 200,000$ y; alternative names for the α Lib Aa,Ab pairing and the single star α Lib B are α^2 Lib and α^1 Lib, respectively, with "1" signalling the fact that α^1 Lib, lying to W of α^2 Lib, although fainter than "2", is the earlier of the two in its crossing of the local meridian</p> <p>¶ one of α Lib Aa, α Lib Ab is overabundant in some metals, perhaps due to influence of its very close SB companion (angular distance ~10 mas, separation a few tenths of 1 AU)</p> <p>¶ lunar occultations are possible, planetary occultations possible yet rare</p>
β	Lup	15 00.0 -43 13	2.68 [†]	-0.18	B2 IV [†]	9	-2.7	380	0.054	222	0 SB	<p>has been claimed to be low-amplitude (β Cep) var.: dominant period 0.232 d</p> <p>¶ fast rotator (< 3.4 d)</p> <p>¶ low metallicity</p>
κ	Cen Aa [†]	15 00.6 -42 11	3.13 [†]	-0.21	B2 V	9	-2.2	400	0.029	218	+8 SB	<p>strictly a triple system, Aa+Ab+B; B mag. 11.5; AB 4", PA: 84°→83°, 1926→2000, separation ≥ 470 AU; ≥ 3000 y; Aa+Ab separation possibly ~10 AU, period possibly ~10 y (http://stars.astro.illinois.edu/sow/kappacen.html discusses various physical uncertainties)</p> <p>¶ line profiles vary; although the Aa+Ab binarity has made variability classification difficult, κ Cen Aa is classified by AAVSO(VSX), viewed 2021 Jan. 16, as a variable of the β Cep class (3.13–3.14 in V, 0.095325 d)</p> <p>Nekkar</p>
β	Boo	15 02.8 +40 18	3.49	0.96	G8 IIIa [†]	14.5	-0.7	230	0.049	234	-20 V?	<p>Ba 0.4, Fe -0.5</p> <p>1995A&A...296..509H discusses the puzzling flare seen by <i>ROSAT</i> 1993 Aug. 08 (unusual for a lone M giant; it is possible, but seems unlikely, that flare came instead from an undetected M-dwarf companion; the mild Ba enhancement is, admittedly, consistent with presence of such a companion); slow rotator (~200 d)</p> <p>[THIS STAR ONLY IN ONLINE VERSION OF TABLE]</p> <p>semireg. var.: 3.20–3.46 in V, mean period 20 d</p> <p>Brachium there is also rapid microvariability</p> <p>¶ highly evolved (on AGB, with dead carbon-oxygen core)</p> <p>Nekkar</p>
σ	Lib	15 05.3 -25 22	3.25v [†]	1.67	M2.5 III	11	-1.5	290	0.083	239	-4	<p>semireg. var.: 3.20–3.46 in V, mean period 20 d</p> <p>Brachium there is also rapid microvariability</p> <p>¶ highly evolved (on AGB, with dead carbon-oxygen core)</p>
ζ	Lup A [†]	15 13.8 -52 11	3.50	0.92	G8 III	~27.8	0.6	117	0.133	238	-10	<p>B: 6.74; 71.60" (2016), PA: 249°→249°, 1826→2016 separation ≥ 2600 AU; shared proper motion suggests true binarity (period possibly $\geq 68,000$ y)</p> <p>¶ ζ Lup A is in evolutionary terms on "Red Clump" (Sun-like when still on MS, but helium flash now finished, core helium fusion now underway)</p>
δ	Boo A [†]	15 16.4 +33 14	3.46	0.96	G8 III Fe-1 [†]	~26.8	0.6	122	0.140	143	-12 SB	<p>a very wide double: B is mag. 7.89, 105" PA: 84°→78°, 1780→2017, separation ≥ 3800 AU, period 120,000 y (with shared proper motion indicating true binarity)</p> <p>¶ δ Boo A is CN weak; δ Boo B could be a subdwarf, consistently with the observed low metallicity of δ Boo A</p> <p>¶ δ Boo A is in evolutionary terms a "Red Clump" star (core helium fusion now underway)</p>
β	Lib	15 18.2 -9 28	2.61 [†]	-0.07	B8 IIIIn	~17.6	-1.2	190	0.100	259	-35 SB	<p>Zubeneschamali</p> <p>flagged by AAVSO(VSX), viewed 2021 Jan. 16, as "suspected variable lacking</p>

												deeper studies,” with V mag. 2.60–2.62 (and yet Eratosthenes, resp. Ptolemy, asserted β Lib to be brighter than, resp. equal to, α Sco)
γ	UMi	15 20.7 +71 45	3.00 [†]	0.06	A3 III [†]	6.7	-2.9	490	0.025	315	-4 V	<p>‡ rapid rotator</p> <p>‡ E(B-V) = -0.02</p> <p>Pherkad</p> <p>a rapid rotator, and (despite being in MK type A, not B) said to be a variable shell star (cf 2000A&A...354...157H; BSC5: “shell possibly variable,” H and CaII variable); AAVSO(VSX), however, viewed 2021 Jan. 16, classifies this as a low-amplitude variable in the δ Sct group</p>
γ	TrA	15 20.9 -68 45	2.87	0.01	A1 IIIIn [†]	17.7	-0.9	184	0.074	244	-3 V	<p>has been asserted to be chemically anomalous (Eu overabundance), and also, not quite consistently, has been classed as a rapid rotator (< 1.2 d)</p> <p>‡ although we here give MK luminosity class III, class V has also been asserted;</p> <p>Astron. Alm. (epoch 2021.5) assigns MK type A1 III</p> <p>‡ IR excess has been asserted (circumstellar disk?)</p>
δ	Lup	15 22.8 -40 43	3.22 [†]	-0.23	B1.5 IVn	4	-3.9	900	0.032	218	0 V?	<p>rapid rotator (< 2.4 d)</p> <p>‡ a (low-amplitude) variable in the β Cep Group, 3.2–3.24 in V (AAVSO(VSX), viewed 2021 Jan. 16), with a single period known, 0.16547 d (cf 2007MNRAS.377..645S)</p>
ϵ	Lup Aa [†]	15 24.2 -44 46	3.37 [†]	-0.19	B2 IV-V	6	-2.6	500	-0.030	-230	+8 SB2	<p>A: 3.56; B: 5.04, 0.1”, PA:285°→53°, 1883→2019 orbit 737 y;</p> <p>in more detail, a (probable) hierarchical quadruple; although B experiences A as essentially a point mass, in fact A is SB, for which 2005A&A...440..249U gives SB period 4.55970 d (classifying primary as a suspect β Cep variable and secondary as a new β Cep variable), experiencing AB, on the other hand, as essentially a point mass is the (probably) gravitationally bound C (lying at angular distance 26.1” in 2016; separation \geq 4100 AU; if gravitationally bound, then period \geq 60,000 y); in its stable kinematics, this putative hierarchical quadruple may be contrasted with the unstable, nonhierarchical θ Ori system, and in its detailed organization with the stable, hierarchical, but mere “double-double” ϵ Lyr system</p>
ι	Dra A+1P [†]	15 25.4 +58 53	3.29	1.17	K2 III [†]	32.2	0.8	101	0.019	334	-11	<p>2002ApJ...576..478F announces substellar-mass companion and discusses possibility of transits; this is the first discovery of a planet or brown dwarf (IAU name: Hypatia) orbiting a star that has finished stable core-hydrogen fusion; http://exoplanet.eu/catalog/HIP%2075458_b/ may from time to time have updates;</p> <p>its substellar companion notwithstanding, ι Dra has metallicity only slightly greater than solar ecl.: 2.21–2.32 in B band, 17 d</p> <p>(more precisely, from AAVSO(VSX) as viewed 2021 Jan. 16, 17.359907 d): a detached binary, with neither component filling its Roche lobe; separation 0.13 AU min;</p> <p>as with β Per, so also with α CrB, instrumental photometry reveals both the primary and the secondary eclipse;</p> <p>components have not been interferometrically resolved (so WDS-conformant designation is still “α CrB”, not “α CrB A” and “α CrB B”)</p> <p>‡ individual MK types are difficult: primary possibly A0 V, secondary possibly G5</p> <p>‡ primary has IR excess (debris disk?)</p> <p>‡ secondary is X-ray visible and is a rather rapid rotator (~9 d or ~7 d or less, so not tidally locked)</p> <p>‡ non-IAU name Gemma denotes α CrB as “gem of the Northern Crown”</p>
α	CrB	15 35.6 +26 39	2.22v [†]	0.03	A0 IV (composite) [†]	43	0.4	75	0.150	127	+2 SB [†]	<p>Alphecca</p>
γ	Lup A [†]	15 36.6 -41 14	2.80	-0.22	B2 IVn [†]	8	-2.8	400	-0.030	-212	+2 V	<p>A: 3.0; B: 4.4; similar spectra 0.8” (2019)</p> <p>PA: 94°→275°, 1835→2019;</p> <p>max angular distance 1980, min ang. dist. 2075;</p> <p>orbit 190 y: γ Lup AB orbit is seen nearly edge-on; separation 41 AU min, 128 AU max, 84.5 AU average;</p> <p>http://stars.astro.illinois.edu/sow/gammalup.html has an orbit map, showing that observational coverage is imperfect (green for micrometry (with large error bars), violet for photography, blue for interferometry);</p> <p>γ Lup A is itself SB (2.8081 d), making this a</p>

α	Ser A	15 45.3 +6 22	2.63v [†] 1.17	K2 IIIb CN1 [†]	44	0.9	74	0.141	71	+3 V?	<p>hierarchical triple system, with the primary in the γ Lup A pairing a fast rotator (< 1 d, so not tidally locked)</p> <p>¶ BSC5 asserts expanding circumstellar shell, and (citing 1987 Vainu Bappu spectra) notes emission peaks in Hα profiles, says possibly in transition from B to Be semiregular variable (low amplitude)</p> <p>¶ a “strong-lined giant” (although [Fe/H] metallicity is not very much above solar)</p> <p>¶ a modest X-ray source</p> <p>¶ has borne also the (not IAU-official) name Cor Serpentis (“Heart of the Serpent”), despite being the principal luminary of Serpens Caput (“Serpent Head”)</p> <p>Unukalhai</p>
μ	Ser A	15 50.7 -3 30	3.54 -0.04	A0 III	19	0.0	170	0.104	255	-9 SB [†]	<p>announced in 2010NewA...15..324G as astrometric binary, 36 y</p> <p>[THIS STAR ONLY IN ONLINE VERSION OF TABLE]</p>
β	TrA A	15 57.1 -63 30	2.83 0.32	F0 IV [†]	~80.8	2.4	40.4	0.444	205	0	<p><i>Spitzer Space Telescope</i> finds IR excess (debris disk?)</p> <p>¶ rapid rotator (slightly < 1 d), with detectable magnetic field</p> <p>¶ metals vary widely (some overabundant, some underabundant)</p>
π	Sco A	16 00.2 -26 10	2.89 [†] -0.18	B1 V [†]	6	-3.4	600	0.029	203	-3 SB2	<p>A: ecl. binary.: 3.4 & 4.5, 1.57 d, 2.88–2.91 in V (more precisely, from AAVSO(VSX) viewed 2021 Jan. 16, 1.570103 d), circular orbit, possibly tidally locked, separation possibly ~0.07 AU; although system has been said to be of β Lyr type, the AAVSO(VSX) classification is, rather, “rotating ellipsoidal variable” (the stars so close as to be gravitationally distorted into ellipsoids, but neither star deformed into the teardrop shape possible in one β Lyr scenario (the β-Lyr variability-type scenario, namely, in which a component becomes so grossly distended as to fill its Roche lobe; in any β Lyr variable, the shape distortion is by definition so severe as to leave no constant-light segments in the light curve)); inspection of AAVSO archive, 2021 Jan. 19, indicates a longstanding shortage of photometry (and Kaler at http://stars.astro.illinois.edu/sow/pisco.html additionally discusses some difficulties in astrophysical modelling)</p> <p>¶ π Sco B is of MK type B2 V</p> <p>¶ E(B–V) = +0.08</p> <p>Fang</p>
T	CrB A	16 00.4 +25 52	9.8v [†] 1.34	M3 III [†]	—	0.6	2500?	0.011	329	-29 SB	<p>recurrent nova 1866&1946 mags 3&2; ~9.9 2021 Jan. 11 only ten galactic recurrent novae are currently known (2010ApJ...187..275S; these are by definition novae known to recur, and yet lacking the short periods of dwarf novae)</p> <p>¶ T CrB A partner in the recurrent-nova activity, T CrB B, is WD with hot circumstellar accretion (dominating the aggregate T Cr AB signal in UV) of MK type Bep, orbit 227 d or 228 d, separation ~0.5 AU, with angular distance measured in 1946 as 0.3”</p> <p>¶ long documented in Handbook as mag. 10.08, T CrB AB (combined light) brightened from February 2015, attaining ~9.2 in April 2015 (with mag. 9.8, on the other hand, reported on 2018 Nov. 15); Bob King in <i>Sky & Telescope</i> 2016 Apr. 20 gives recent history, and AAVSO has a backgrounder at https://www.aavso.org/t-crb;</p>
η	Lup A [†]	16 01.6 -38 27	3.37 -0.21	B2.5 IVn [†]	7	-2.2	440	0.033	211	+8 V	<p>next eruption 2026, or earlier?</p> <p>B: 7.70, 15.0”, PA:22°→19°, 1834→2016 orbit \geq 26,000 y:</p> <p>a hierarchical system, with remote outlier D at angular distance 135” (separation \geq 18,000 AU, period \geq 750,000 y), with D experiencing the AB pair as essentially a point mass; η Lup C is not part of this (triple) system, C’s angular proximity to AB being a mere line-of-sight coincidence</p> <p>¶ although η Lup A is a rapid rotator (< 1.1 d), there is no evidence of a circumstellar disk, and in particular there seems to be no documentation of “Be phenomenon” spectral behaviour</p>
δ	Sco A [†]	16 01.6 -22 41	2.29 [†] -0.12	B0.3 IVe [†]	7	-3.6	500	-0.037	-196	-7 SB	<p>¶ η Lup B is chemically peculiar</p> <p>periastron outbursts 2000, 2011</p> <p>Dschubba</p> <p>(AB orbit is very elongated) as instance of “Be phenomenon” (but the Be-phenomenon intermittent equatorial disk of gaseous ejecta is observed to be present)</p>

													<p>even before periastron); www.aavso.org/delta-scorpi has recent forum discussion, notably on choice of comparison stars for visual photometric estimates; two typical recent AAVSO V-filter photometry reports, from one and the same observer, are 2020 June 20, mag. 1.86 and 2020 June 26, mag. 1.81; classified at AAVSO(VSX) as a variable of the γ Cas type; consistently with this classification, the primary is a rapid rotator; www.aavso.org/vsots_delsco covers 2000–2011 ¶ AB: 10.8 y, 0.2" (2019); B is of MK type B3 V; http://stars.astro.illinois.edu/sow/dschubba.html discusses multiplicity (in all, possibly quadruple, with hierarchical organization; AB period is 20 d, separation ~0.4 AU) ¶ E(B–V)=+0.16</p>
β	Sco Aa [†]	16 06.7 –19 52	2.56 –0.06	B0.5 V		8 –2.9 400	0.025 192 –1 SB						<p>Aa: 2.62; B: 10.6, 0.3" (2019); C: 4.52, 13" Acrab in gross terms a visual binary (as AB), with separation \geq 2200 AU, period > 16,000 y; but in fact putatively a sextuplet; https://en.wikipedia.org/wiki/Beta_Scorpii summarizes the sextuplet's hierarchy in a diagram (Aa with Ab (6.82 d), and B experiencing Aa+Ab as essentially a point mass (610 y); Ea with Eb (10.7 d), and C experiencing Ea+Eb as essentially a point mass (39 y); the B+AaAb triple is in a wide, > 16,000-y orbit with the C+EaEb triple, around the centre of mass shared by this pair of triples, thereby delivering the gross visual-binary phenomenology) ¶ lunar occultations possible, planetary occultations possible yet rare (1971 May 14 occultation by Jovian satellite Io) ¶ the name Graffias is not IAU-official var.? (2.72–2.75 in V?) Yed Prior[†] ¶ slow rotator ¶ high metallicity ¶ although δ Oph has finished core hydrogen fusion, its exact evolutionary state is uncertain (cf http://stars.astro.illinois.edu/sow/yedprior.html) ¶ Astron. Alm. (epoch 2021.5) assigns MK type M0.5 III ¶ naked-eye neighbour Yed Posterior is a mere optical companion, too greatly separated in space for true binarity; the "prior" and "posterior" in the traditional, and as of 2016 IAU-official, names denote the order in which these two (physically unrelated) stars cross the local meridian ¶ listed in NSV as a suspected variable, and in AAVSO(VSX), viewed 2021 Jan.16, as unobserved; 1992IBVS.3792...1P finds no variability, but says that since NSV V-amplitude is just 0.03 mag., variability cannot be excluded</p>
δ	Oph A	16 15.5 –3 45	2.73 [†] 1.58	M1 III [†]		~19.1 –0.9 171	0.150 198 –20 V						<p>Yed Posterior cyanogen and carbon notably underabundant, suggesting that ε Oph is an interloper from outside the galactic thin disk; Astron. Alm. (epoch 2021.5) assigns MK type G9.5 IIIb Fe–0.5</p>
ε	Oph A	16 19.5 –4 45	3.23 0.97	G9.5 IIIb [†]		31 0.7 106	0.093 64 –10 V						<p>var.: 2.86–2.94, 0.25 d; B: 8.4, B9 V, 20.5" (2019) Alniyat recent studies, including lunar occultation measures, show σ Sco to be a quadruple system, with σ Sco Aa1,Aa2 in fact a spectroscopic binary (33.0 d), and the entire σ Sco A configuration in orbit with a B7 star at angular distance 9.4" (period > 100 y); 2007MNRAS.380.1276N announces interferometric solution for the SB orbit, proposing for primary and secondary the respective MK types B1 III, B1 V; in the SB pair, the primary is a variable of the β Cep type (AAVSO(VSX) viewed 2021 Jan. 16 gives V-mag. range 2.86–2.94, period 0.246839 d; 1992A&A...261..203P discusses period changes) ¶ photography shows σ Sco to be embedded in diffuse nebula ¶ E(B–V) =+0.4 (pronounced reddening) B: 8.2, 4.4", PA: 150°→143°, 1843→2015 Athebyne ¶ a "Red Clump" resident (evolved, presently stable, performing core helium fusion)</p>
σ	Sco Aa1 [†]	16 22.5 –25 39	2.91v [†] 0.13	B1 III [†]		5 –3.7 700	0.019 213 +3 SB [†]						
η	Dra A	16 24.3 +61 28	2.73 [†] 0.91	G8 IIIab		35.4 0.5 92	0.059 343 –14 SB?						

α	Sco A [†]	16 30.7 -26 29	1.06v [†]	1.86	M1.5 Iab [†]	6	-5.1	600	0.026	207	-26 SB	<p> [†] believed to be a slow rotator (~400 d) [†] a modest X-ray source [†] listed by NSV (Kukarkin et al.) as a suspected variable [†] near the radiant of the η Draconids meteor shower smreg. var.: 0.75–1.21, 5.97 y; B: 5.40, 3.2" (2016) Antares[†] PA: 273°→276°, 1847→2016; orbit 2500 y? AAVSO(VSX): semireg. (with some discussion of period; cf also 2013AJ...145...38P, where a true period is found for radial-velocity variations, and the detected variation is judged to be more likely of pulsational than of orbital origin), V mag. 0.75–1.21 (but the variability has also been called irregular); 2018AujAn..29..89H reports that variability was observed by, and incorporated into the oral tradition of, aboriginals in southern Australia; asserted by Eratosthenes to be fainter than β Lib, and by Ptolemy to equal β Lib [†] Astron. Alm. (epoch 2021.5) assigns MK type M1.5 Iab–Ib [†] radius has been studied interferometrically and via lunar occultations (up to 3.4 AU; however, even apart from the problem of pulsation, radius determination of highly evolved red stars is wavelength-dependent); one of the two first-mag. supergiants (the other being α Ori Aa (Betelgeuse)) [†] significant stellar wind, with mass loss almost 1e-6 M\odot/y, within which α Sco B has created a locally ionized region [†] the most massive member of the Sco-Cen Association (the nearest OB association) [†] B shares in the proper motion of A, indicating true binarity: AB separation is \geq 530 AU, period possibly ~1200 y [†] location (within zodiac) makes the classical Greek name for “rival of Mars” appropriate not only as regards naked-eye colour but also as regards sky geometry [†] 1972BAAs...82..43IK describes the 18.6-year 1940-through-2050 cycle of lunar occultation possibilities </p>
β	Her Aa	16 31.1 +21 27	2.78 [†]	0.95	G7 IIIa [†]	23	-0.4	140	0.100	261	-26 SB [†]	<p> SB period computed 1908, and again 2008, in both cases ~410 d; 1977ApJ...214L..79B announces speckle-interferometry resolution of the β Her Aa,Ab SB, with angular distance 43 mas [†] suspected variable (NSV (Kukarkin, et al., online) suggests V-mag. range 2.76–2.81, and AAVSO(VSX) viewed 2021 Jan. 16 concurs) [†] X-ray emission from the SB primary indicates magnetic activity [†] Astron. Alm. (epoch 2021.5) assigns MK type G7 IIIa Fe–0.5 [†] Kaler, noting that primary has N enhanced relative to C, says in his overall summation “a very normal star for its state of age” [†] “Kornephoros” = Gk “club-bearer,” in reference to the weapon of Hercules (compare α Her, which in the pictorial-atlas tradition, marks the hero’s head) Kornephoros[†] </p>
τ	Sco	16 37.2 -28 16	2.82	-0.21	B0 V [†]	7	-3.0	500	0.025	203	+2 V	<p> intrinsically more luminous than σ Sco, but more heavily obscured by ISM [†] anomalous in its UV lines (P Cyg profile) [†] O and Fe are underabundant [†] 2006MNRAS.370..629D discusses τ Sco magnetic topology (poloidal, with also a warped toroidal component of modest strength), including both its origin (more likely a fossil field from the star’s (recent) birth than a dynamo effect) and its connection with winds and with the observed hard X-ray emission; the authors note that the topology is stable over the 1.5-y period of their observations (in contrast with a strongly differential-rotation star, such as Sun); in additionally announcing a (refined) rotation period of 41.033 d, the authors comment, “the second-slowest rotator so far known among high-mass stars” [†] Kaler: “among the most-observed stars in the sky” [†] E(B-V)=+0.06 [†] the τ Sco name Paikauhale[†] </p>

ζ	Oph	16 38.3 –10 37	2.54 [†]	0.04	O9.5 Vne [†]	9	-2.7	370	0.029 [†]	32	-15 V [†]	<p>IAU-approved in 2018 Aug. 10; the not-IAU-official “Al Niyat,” or “the arteries of the Heart,” on the other hand, denotes σ Sco and τ Sco jointly, as flanking α Sco the nearest O-type star (and consistently with this extreme temperature, resident in an H II region) ¶ unusual in being an “Oe”, i.e. an O-star instance of the “Be phenomenon” ¶ “runaway star” (consistently with this extreme speed-relative-to-LSR, forming bow shock in ISM), perhaps formerly the secondary in a binary pair whose primary perished in a supernova; 2011AN...332..147H confirms magnetic field, discusses X-ray properties, suggests PSR B1919+10 as remnant of the hypothesized defunct companion ¶ line of sight to ζ Oph is one of the most used in spectroscopic studies of ISM ¶ 2014MNRAS.440.1674H is a recent discussion of variability, from radial and non-radial pulsation modes; AAVSO(VSX), assigning magnitude range 2.56–2.58 in V, follows GCVS in treating ζ Oph as a variable with Be-phenomenon behaviour, and yet lacking the history of outbursts founds in the γ Cas class; ζ Oph is, on the other hand, classified as γ Cas-variable (and is termed a shell star) in BSC5; still elsewhere, ζ Oph has been treated as a prototype for the “ζ Oph variables” ¶ E(B-V)=+0.32 (pronounced reddening; if ISM were not present, ζ Oph would reach nearly first mag.) ¶ recapitulations of recent ζ Oph studies include 2012MNRAS.427L..50G (MK classification problem, also mass-loss rate in context of “weak-wind problem”), 2014MNRAS.440.1674H (rotation, pulsation, Hα emission episodes, inferred circumstellar decretion disk, satellite-based photometry), 2015ApJ...800..132C (distance, age, mass, effective temperature, bow shock in ISM, ...); additionally, 2012A&A...543A..56D is among the papers describing not only the specific interaction of ζ Oph with ISM, but also the quite general ISM bow-shock topic (noting inter alia that not all runaway stars produce bow shocks) B: 5.40, G7 V, 1.6” (2019), orbit 34.45 y orbit well studied since F.G.W. von Struve 1826 micrometry (however, it was Herschel, not von Struve, who discovered the binarity); separation 8 AU min, 21 AU max, 15 AU average, 34.45 y; one of the few binaries in which ratio of B mass to sum of A and B masses can be studied both astrometrically and spectroscopically ¶ Astron. Alm. (epoch 2021.5) assigns MK type G0 IV ¶ ζ Her A is unusual in its evolutionary phase, being in the Hertzsprung Gap (and so in rapid evolutionary transition) ¶ 2001A&A...379..245M summarizes previous work, presents detailed physical modelling for A and B, and discusses astroseismology, remarking in conclusion that “among the binaries to be calibrated with some confidence, ζ Herculis is one of the most interesting owing to the difference of evolutionary state of components”</p>
ζ	Her A [†]	16 42.1 +31 34	2.62	0.65	G1 IV [†]	93	2.7	35	-0.575	-307	-70 [†] SB	

η	Her A	16 43.6 +38 53	3.48	0.92	G7.5 IIIb Fe-1 [†]	30.0	0.9	109	0.092	157	+8 V?	<p>¶ high velocity relative to Sun</p> <p>in evolutionary terms a resident of the “Red Clump” (fusing helium in stable core)</p> <p>¶ Astron. Alm. (epoch 2021.5) assigns MK type G7 III Fe-1</p> <p>¶ Fe is notably underabundant</p>
α	TrA A	16 51.0 -69 04	1.91	1.45	K2 IIb-IIIa [†]	~8.4	-3.5	390	0.036	150	-3	<p>Atria</p> <p>anomalous for its MK type, with flares and X-ray emission, perhaps from as-yet-undetected magnetically active companion (a companion would indeed be indicated by the claimed “barium star” status of α TrA;</p> <p>http://stars.astro.illinois.edu/sow/atRIA.html, in discussing the possibility of a companion, also remarks, however, “the classic ‘hybrid star,’ a giant that shows evidence for blowing a cool wind from its surface, yet having a hot surrounding magnetic corona at the same time”;</p> <p>https://ntrs.nasa.gov/archive/nasa/casi.ntrs.nasa.gov/20040086627.pdf further discusses both α TrA and β Dra, as (solitary) stars, which are in this particular posited sense “hybrid”</p>
ϵ	Sco	16 51.6 -34 20	2.29	1.14	K2 III	51	0.8	64	0.666 [†]	247	-3 [†]	<p>Larawag</p> <p>slow rotator (possibly even 1.3 y)</p> <p>¶ evolved, and yet not a clump star;</p> <p>http://stars.astro.illinois.edu/sow/epsSCO.html discusses the uncertainty in evolutionary stage (brightening, with He core as yet awaiting ignition? dimming, with He core fusion in progress? or brightening, with dead C-and-O core, He-core fusion now over?)</p> <p>¶ high velocity relative to Sun indicates origin outside the galactic thin disk (and metal underabundances are consistent with such an origin)</p>
μ^1	Sco A	16 53.3 -38 05	3.00v [†] -0.20		B1.5 IVn	7	-2.9	500	0.024 [†]	206-25 [†]	SB2 [†]	<p>Xamidimura</p> <p>(more precisely, in AAVSO(VSX) viewed 2021 Jan. 16, 1.44626907 d); semidetached, partially eclipsing, binary system, with mass transfer, resembling β Lyr in its never-constant light and in exhibiting both primary and secondary minima;</p> <p>1948MNRAS.108.398S gives the light curve, and also discusses early observational history (this is the third SB discovery in astronomy (made by Bailey, 1896)); separation is ~0.07 AU</p> <p>¶ μ^1 Sco and μ^2 Sco are not gravitationally bound, although both belong to the (gravitationally unbound) “Upper Sco” subgroup of the Sco-Cen Association</p>
κ	Oph	16 58.7 +9 21	3.19 [†]	1.16	K2 III	36	1.0	91	0.292 [†]	268	-56 V [†]	<p>slow rotator (possibly as slow as 1.6 y)</p> <p>¶ historical assertion of variability may be due to a confusion between κ Oph and χ Oph; completely apart from this historical problem, however, 2001BaltA...10..593A discusses the possible variability both of κ Oph and of other Red Clump stars</p> <p>¶ high velocity relative to Sun suggests origin outside the galactic thin disk</p>
ζ	Ara	17 00.4 -56 01	3.12	1.55	K4 III	7	-2.7	490	0.041	206	-6	<p>one of the rather rare instances of a giant excessively bright in far IR</p> <p>(1997A&A...323..513P suggests that such giants are more likely to be radiating their IR excess from circumstellar debris disks than from winds, and so are to be considered evolved-star analogues of the unevolved (and IR-bright) α Lyr)</p>
ζ	Dra A	17 08.9 +65 41	3.17	-0.12	B6 III [†]	10	-1.8	330	0.028	314	-17 V	<p>Aldhibah</p> <p>A,B: mags 3.2, 4.2, respectively; 0.10" (1994)</p> <p>¶ given the recent formation of the ζ Dra system, Fe is anomalously underabundant</p> <p>¶ E(B-V) = +0.03</p>
η	Oph AB [†]	17 11.6 -15 45	2.43	0.06	A1 IV + A1 IV? [†]	37	0.3	90	-0.107	~22	-1 SB	<p>Sabik[†]</p> <p>A: 3.0; B: 3.3, A3 V, 0.5" (2019), orbit 87.6 y highly eccentric orbit: separation 2 AU min, 65 AU max</p> <p>¶ under IAU rules, “Sabik” designates η Oph A, not η Oph B</p> <p>¶ our present assignment of MK types (confident for η Oph A, tentative for η Oph B) is from the literature; our Handbook predecessor R.F. Garrison, however, himself favoured “A2.5 Va,” perhaps for the AB composite; Astron. Alm. (epoch 2021.5), perhaps again for the AB composite, assigns MK type “A2 Va+ (Sr)”</p> <p>¶ it is possible that A, or B, or both A and B, are superabundant in metals</p>

η	Sco	17 13.7 -43 16	3.32	0.44	F5 IV [†]	-44.4	1.6	73	0.290	175	-28	<p>Garrison legacy dwarf MK type “F2 V:p(Cr)” (complexity of the legacy type hints at difficulties in classification; even “dwarf barium star” has been asserted elsewhere; Astron. Alm. (epoch 2021.5) assigns the same MK type as legacy-Garrison), NASA NStars, in work summarized at 2006AJ...132..161G (with Garrison the third author) is our authority for subgiant MK type F5 IV</p> <p>¶ rapid rotator (< 1 d); the observed X-ray emission is consistent with magnetic effects (including coronal heating?) stemming from rapid rotation semireg. var.: 2.73–3.60 in V; B: 5.4, 5.0” (2019)</p>
α	Her Aa [†]	17 15.6 +14 22	2.78v [†]	1.16	M5 Ib–II	9	-2.4	400	0.032	347	-33 V	<p>Rasalgethi[†]</p> <p>AB PA: 117°→103°, 1777→2019; orbit > 3000 y; α Her A is strictly SB Aa+Ab (~10 y), and α Her B also strictly SB Ba+Bb (51.578 d, separation 0.4 AU), making α Her at least a (kinematically stable, hierarchically organized) quadruplet; more distant α Her C and α Her D are not necessarily gravitationally bound to the quadruplet; separation of Aa+Ab and Ba+Bb (each of these binaries experiencing its distant companion binary as essentially a point mass) is > 500 AU</p> <p>¶ 1956ApJ...123..210D discusses mass loss from (in 21st-century nomenclature) α Her Aa, and the consequent circumstellar material (so copious as to encompass even Ba+Bb)</p> <p>¶ https://en.wikipedia.org/wiki/List_of_largest_stars shows ranking of α Her Aa (radius ~1.5 AU, or more) in the overall known cosmic population of giants, supergiants, and hypergiants</p> <p>¶ in classification of AAVSO(VSX), α Her Aa is a “semi-regular late-type giant”, with V-mag. range 2.73–3.60; 2001PASP...113..983P reports timescales both of 80–140 d and of 1000–3000 d, while underscoring the complexity of the variation (in their Figure 5, authors show light curve)</p> <p>¶ in the pictorial-atlas tradition, α Her marks the head of hero Hercules (with β Her marking his club; for summer-evening observers in the northern hemisphere, the hero is to be visualized inverted, with feet high in the sky, club and head lower)</p>
π	Her	17 15.8 +36 47	3.16 [†]	1.44	K3 IIab [†]	8.7	-2.2	380	0.027	276	-26 V?	<p>¶ Astron. Alm. (epoch 2021.5) assigns MK type K3 II</p> <p>¶ low-amplitude photometric variations with low-amplitude radial-velocity variations, 613 d, perhaps favour the hypothesis of non-radial pulsation over the competing hypotheses of an undetected low-mass companion and of rotation with starspots</p>
δ	Her Aa [†]	17 15.9 +24 49	3.12	0.08	A1 Vann	43.4	1.3	75	-0.158	-188	-40 SB [†]	<p>Sarin</p> <p>B: 8.3, 13.7” (2019) is mere optical companion δ Her A, being SB (and also resolved as a binary in interferometry, with angular distance 60 mas; separation \geq 1.45 AU, period \geq 335 d), is strictly δ Her Aa,Ab</p> <p>¶ δ Her Aa is a fast rotator (< 9 h)</p> <p>¶ as with δ Her B, so also δ Her C and δ Her D, at respective angular distances 174” (2013) and 192” (2009), are most likely mere optical companions</p>
θ	Oph A [†]	17 23.3 -25 01	3.27v [†]	-0.19	B2 IV	-7.5	-2.4	440	0.025	197	-2 SB [†]	<p>variable. in β Cep class: 3.25–3.31 in V, 0.14 d http://stars.astro.illinois.edu/sow/thetaoph.html discusses uncertainties in multiplicity: perhaps SB with outlying, gravitationally bound, companion at angular distance 0.15”;</p> <p>the primary in the putative SB is a β Cep variable, 0.140531 d</p> <p>¶ occasional lunar occultations</p>
β	Ara	17 27.1 -55 33	2.84	1.48	K3 Ib–IIa [†]	5	-3.6	600	0.027	199	0	<p>slow rotator (possibly as much as 2.33 y)</p> <p>¶ high metallicity</p> <p>¶ not gravitationally bound to γ Ara AB</p>
γ	Ara A	17 27.2 -56 24	3.31	-0.15	B1 Ib	-2.9	-4.4	1100	0.016	182	-3 V	<p>broad lines for Ib</p> <p>¶ γ Ara A is rapid rotator (both “~4.8 d” and “< 2.5 d” have been asserted, and yet rapid rotation is unusual for the (evolved) γ Ara A luminosity class)</p> <p>¶ 1997A&A...318..157P</p>

β	Dra A	17 30.9 +52 17	2.79 [†]	0.95 [†]	G2 Ib-IIa	8.6	-2.5	380	0.020	308	-20 V	Rastaban
												<p>finds via <i>IUE</i> spectroscopy that, consistently with this rapid rotation, the stellar wind of γ Ara A may be equatorially enhanced (and more generally, that the wind is variable, and is structured with two components, its structure being not typical of stars in this portion of the HR diagram)</p> <p>¶ γ Ara AB is not gravitationally bound to β Ara</p> <p>¶ E(B-V)=+0.08</p> <p>in evolutionary terms, β Dra A is somewhat unusual, as a yellow more-than-giant (having been a stable core-hydrogen fuser just 0.5 My ago, the star is in transition to being redder, and of still larger radius)</p> <p>¶ it is also odd that β Dra A, while lying in the HR diagram Instability Strip, has not been observed to pulsate</p>
ν	Sco	17 32.2 -37 19	2.70	-0.18	B2 IV [†]	6	-3.5	600	0.030	185	+8 SB	Lesath
												<p>although we here give spectral type B, type Be has also been asserted</p> <p>¶ ν Sco and λ Sco are not gravitationally bound (although both belong to the (gravitationally unbound) Sco-Cen OB association, and have as an optical double been called the "Cat's Eyes")</p> <p>¶ E(B-V)=+0.02</p>
α	Ara A	17 33.5 -49 53	2.84	-0.14	B2 Vne [†]	12 [†]	-1.7	300 [†]	0.075	206	0 SB	
												<p>an instance of the "Be phenomenon," with (since the star, with its equatorial ejecta, is seen nearly equator-on) "shell" spectrum:</p> <p>2007A&A...464...59M says, "For the first time, we obtain the clear evidence that the [equatorial ejecta] disk is in Keplerian rotation, closing a debate that has continued since the discovery of the first Be star γ Cas by Father Secchi"; on the authors' modelling, α Ara is rotating near breakup speed (and consequently is oblate), with an enhanced wind from its poles; the authors assert the possibility that equatorial ejecta disk is truncated by an unseen companion at 32 stellar radii</p> <p>¶ IR excess is unusually high for a Be star</p> <p>¶ for problem of distance (the <i>HIPPARCOS</i> distance given here may be too high) cf 2005A&A...435..275C and 2007A&A...464...59M</p>
λ	Sco Aa,Ab [†]	17 35.1 -37 07	1.62v [†]	-0.23	B1.5 IV + n.a.	>6 [†]	-4.6	400 [†]	0.032	195	-3 SB2	Shaula
												<p>eclipsing?, variable 1.59-1.65 in V, 0.2137 d</p> <p>λ Sco Aa,Ab are respectively of mags. ~2.1, ~2.8; strictly a hierarchical triple system, with orbits studied interferometrically in 2006MNRAS.370..884T; the narrow λ Sco Aa,Ab pair has period 5.9525 d, with eclipsing, and the wider λ Sco AB pairing has period ~1000 d (B is elusive, at mag. ~15)</p> <p>¶ although <i>HIPPARCOS</i> parallax entails distance (to one significant figure) 600 ly, 2006MNRAS.370..884T entails instead (to one significant figure) 400 ly: generally speaking, <i>HIPPARCOS</i>, like other fine-grained parallax measures of distance, risks degradation if a star has a stellar-mass gravitationally bound companion</p> <p>¶ λ Sco Aa is a β Cep variable; since full orbital coverage is available in this case (as also with β Cep itself; in most or all other β Cep-class cases, full orbital coverage is presently unavailable), mass determination becomes feasible, making the λ Sco Aa,Ab binary important in β Cep-variable research; λ Sco Ab is itself of interest, as a possible pre-main-sequence star (this would be consistent with the observed X-ray emission)</p> <p>¶ 1975MNRAS.173..709L gives some photometry</p> <p>¶ a flare was observed in vicinity of λ Sco on 1975 Jun. 01</p> <p>¶ 2004A&A...427..581U summarizes previous work on λ Sco, discusses masses, discusses tidal effect on β Cep pulsation</p> <p>¶ λ Sco and ν Sco are not gravitationally bound (although both belong to the (gravitationally unbound) Sco-Cen OB association, and have as an optical double been jointly called the "Cat's Eyes")</p> <p>¶ E(B-V)=+0.03</p>

α	Oph A [†]	17 35.9 +12 33	2.08 [†]	0.16	A5 Vnn	67	1.2	49	0.247	154	+13 SB?	low-amplitude var. (δ Set type, γ Dor type) Rasalhague \P α Oph A is a fast rotator (oblateness has been imaged interferometrically), seen nearly equator-on; the binary system has become better understood with the recent, 2011ApJ...726..104H , determination of masses and orbit geometry, through coronagraph and adoptive optics (period 3148.4 d, angular distance at periastron passage \sim 50 mas; the now-achieved determination of masses in this particular system has implications for astrophysics generally, since it potentially facilitates the refining of numerical models for rapidly rotating hot stars) \P astroseismology mission <i>MOST</i> has identified \sim 50 pulsational modes in α Oph A \P α Oph B is mag. 5, PA: 243 $^{\circ}$ \rightarrow 238 $^{\circ}$, 1982 \rightarrow 2018
ξ	Ser Aa [†]	17 38.8 -15 25	3.54 [†]	0.26	F0 IIIb [†]	31	1.0	105	0.073	215	-43 SB	hierarchically organized triple system, comprising ξ Ser Aa and ξ Ser Ab, experienced as essentially a point mass by the outlying ξ Ser B; period of single-lined SB Aa+Ab is 2.29 d, with angular distance (in 1987) 0.30"; B is mag. 13.0, at angular distance 24.5" (2015), with PA: 81 $^{\circ}$ \rightarrow 78 $^{\circ}$, 1943 \rightarrow 2015, and period possibly \sim 15,000 y \P ξ Ser Aa has been asserted to be very slightly hotter than Garrison's "F0 IIIb", and moreover to be chemically peculiar, being on this (more recent?) determination of MK type A9 IIIp Sr; additionally, somewhere in the literature, δ Set variability has been asserted or conjectured (Kaler comments at http://stars.astro.illinois.edu/sow/xiser.html : "the star /.../ remains cryptic") [THIS STAR ONLY IN ONLINE VERSION OF TABLE]
θ	Sco A [†]	17 38.9 -43 01	2.0	0.41	F1 III	\sim 11	-3.0	300	0.006	119	+1	Sargas[†] rapid rotator, in the sense that $v \sin i$ is (according to 2005yCat.3244...0G) 125.0 km/s; since, however, θ Sco is a (rapidly evolving) giant, its high $v \sin i$ may correspond to a not-spectacularly short rotation period, of up to 10 d; if, as asserted in literature, θ Sco A truly is a rapid rotator, it will resemble β Cas A (a rapid rotator that, strikingly, has evolved beyond the MS) \P optical companion, mag. 5.36, 6.50"; PA: 322 $^{\circ}$ \rightarrow 315 $^{\circ}$, 1896 \rightarrow 1991 \P although it is the Sumerian name Sargas that is IAU-official as of 2016 Aug. 21, θ Sco, like κ Sco, has also been known under the different, not IAU-official, name Girtab (originally applied by the Sumerians to an entire asterism)
κ	Sco [†]	17 44.0 -39 02	2.41 [†]	-0.17	B1.5 III	7	-3.5	480	0.026	193	-14 SB	low-amplitude variability: 2.41-2.42 in V \P since the SB has not yet been resolved, even interferometrically, WDS cannot yet write " κ Sco A" and " κ Sco B"; the SB has orbital period 195.65 d, with separation 1.7 AU \P the SB primary is a rapid rotator (1.9 d), and additionally is a variable of β Cep type (0.1998303 d in AAVSO(VSX) as viewed 2021 Jan. 16; 1975MNRAS.173..709L gives some photometry, confirming a beat period) \P κ Sco, as a single naked-eye object, has (like θ Sco) been known under the different, not IAU-official, name Girtab (originally applied by the Sumerians to an entire asterism)
β	Oph	17 44.5 +4 34	2.76 [†]	1.17	K2 III [†]	\sim 39.8	0.8	82	0.165	345	-12 V	Cebalrai \P Astron. Alm. (epoch 2021.5) assigns MK type K2 III CN 0.5 \P 1996ApJ...468..391H finds multiple pulsation periods, in behaviour paralleling α Boo ("it may well be that these [two] stars represent a new class of radially and unradially pulsating stars"), and also a possible long period of 142.3 d; the authors suggest that if the latter is real, then although the more likely explanation is a 142.3-d rotation, nevertheless gravitational pull from an unknown exoplanet is conceivable

μ	Her Aa [†]	17 47.3 +27 43	3.42	0.75	G5 IV	~120.3	3.8	27.1	0.804	201	-16 V	BC: 9.78, 35.5", PA:240°→249°, 1781→2015 orbit \geq 3700 y: stable quadruple system, in fact the third-closest quadruple star system to the Sun; system is organized as a "double double," being one of the best-studied such systems (2016AJ...151..169R, consequently says μ Her "serves as an archetype for understanding stellar system formation"): μ Her Aa,Ab is in tight orbit, and μ Her BC is in tight orbit, with each of these two pairs experiencing the other as essentially a point mass; 2016AJ...151..169R gives Aa,Ab a period of ~100 y, with wide uncertainties, concluding also that Ab is an M-dwarf (as opposed to a mere substellar object); BC has period 43.127 y, separation 1.5 AU min, 3.6 AU max, 2.2 AU average; the separation of the Aa,Ab centre of mass from the BC centre of mass is \geq 300 AU, with orbital period \geq 3700 y ¶ despite having finished core-hydrogen fusion, μ Her Aa is a fast rotator, and is consequently magnetically active and an X-ray source
ι^1	Sco A	17 49.1 -40 08	2.99	0.51 [†]	F2 Ia	2 [†]	-5.9	2000 [†]	0.006	180	-28 SB	B: mag.13, 38" (2000) ¶ a rare instance of a yellow supergiant (dead helium core; the star is now cooling, and is now in transition to the less exotic status of red supergiant) ¶ radius estimates vary; CADARS (2001A&A...367..521P) value is ~1.9 AU ¶ mass loss ~1e-7 M _⊙ /y ¶ slow rotator (\geq 0.5 y) ¶ distance and mass are rather uncertain ¶ the modest angular distance of ι^1 Sco from ι^2 Sco is the result of a mere line-of-sight coincidence (with ι^2 ~2 times as distant as ι^1 ; again by coincidence, not ι^1 alone, but also ι^2 , is a supergiant)
G	Sco A	17 51.3 -37 03	3.19	1.19	K2 III	25.9	0.3	126	0.049	56	+25	HR6630, Fuyue although masses of K giants are in general uncertain, in this particular case the mass is known via WIRE salvage-mission astroseismology (being determined in 2008ApJ...674L..53S as 1.44 M _⊙ , with just a 15% uncertainty)
γ	Dra A	17 57.1 +51 29	2.24	1.52	K5 III [†]	21.1	-1.1	154	0.024	200	-28	Eltanin in 1728, James Bradley used γ Dra to demonstrate aberration of light ("velocity aberration"); his demonstration strongly confirmed the heliocentric (and thus non-Ptolemaic) kinematics of the Solar System ¶ Fe is slightly underabundant brown-dwarf companions with masses \leq 24 \times Jupiter and \leq 27 \times Jupiter (deuterium fusion begins at a lower mass, 13 \times Jupiter), periods 530.3 d and 3190 d (Quirrenbach et al. 2011, and additionally 2012PASJ...64..135S; the latter paper suggests formation in circumstellar disk, with subsequent migration, in a scenario reminiscent of planet and exoplanet formation): this is the third star found to be hosting two brown dwarfs ¶ slow rotator (\leq 234 d) ¶ far-IR variability has been suspected ¶ CN underabundant, Fe overabundant ¶ Astron. Alm. (epoch 2021.5) assigns MK type G9 IIIa
ν	Oph +2P [†]	18 00.2 -9 46	3.32	0.99	G9.5 IIIa [†]	22	0.0	150	0.117	185	+13	Alnasl metals underabundant ¶ Astron. Alm. (epoch 2021.5) assigns MK type K0 ⁺ III ¶ ϵ Sgr and the γ^2 - γ^1 Sgr pair serve as pointers to Baade's Window ¶ angular proximity of γ^1 Sgr (= W Sgr; ~50 arcmin, to ~N of γ^2 Sgr) is a mere line-of-sight coincidence
γ^2	Sgr	18 07.2 -30 25	2.98	0.98	K0 III [†]	34	0.6	97	0.189	197	+22 SB	Alnasl metals underabundant ¶ Astron. Alm. (epoch 2021.5) assigns MK type K0 ⁺ III ¶ ϵ Sgr and the γ^2 - γ^1 Sgr pair serve as pointers to Baade's Window ¶ angular proximity of γ^1 Sgr (= W Sgr; ~50 arcmin, to ~N of γ^2 Sgr) is a mere line-of-sight coincidence
η	Sgr A [†]	18 19.1 -36 45	3.10 ν^{\dagger}	1.5	M3.5 IIIab	22	-0.2	~146	0.211	218	+1 V?	irreg. var.: 3.05-3.12; B: 8.00, G8: IV.; 3.5" (2016) PA:100°→110°, 1879→2016; orbit \geq 1270 y, separation \geq 165 AU ¶ η Sgr A is variously asserted to be on the (very highly evolved) HR diagram AGB or at the tip of the RGB ¶ η Sgr A is in the AAVSO(VSX)

δ	Sgr A	18 22.4 -29 49	2.72	1.38	K2.5 IIIa [†]	9	-2.4	350	0.041	128	-20 V?	<p>classification an “LB,” i.e. a slow irregular variable</p> <p>¶ temperature of η Sgr A not yet well determined? Kaus Media[†]</p> <p>¶ Astron. Alm. (epoch 2021.5) assigns MK type K2.5 IIIa CN 0.5</p> <p>¶ possibly a weak barium (Ba) star, δ Sgr A possesses (as expected for a Ba star) a WD companion</p> <p>¶ temperature of δ Sgr A not yet well determined?</p> <p>¶ “Kaus” is Arabic “bow,” with Kaus Borealis (λ Sgr), Kaus Media (δ Sgr), and Kaus Australis (ϵ Sgr) the three delineating stars of the archer’s bow; by coincidence, the archer turns out to be aiming rather close both to Baade’s Window and (prolonging the line of firing) to the Sgr A* black hole at the galaxy’s centre</p>
η	Ser A	18 22.4 -2 54	3.23	0.94	K0 III-IV [†]	54	1.9	~60.5	0.890	218	+9 V?	<p>slow rotator (but ≤ 1.9 y)</p> <p>¶ high velocity relative to Sun suggests that η Ser is an interloper (born outside the galactic thin disk? consistently with this conjecture, Fe is underabundant)</p>
ϵ	Sgr A	18 25.6 -34 22	1.79	-0.03	A0 II n (shell?) [†]	23	-1.4	~143	0.130	198	-15	<p>Kaus Australis[†]</p> <p>fast rotator (consistent with shell-star classification); as might be predicted for a fast rotator, a magnetic field, and also X-ray emission, have been detected</p> <p>¶ Astron. Alm. (epoch 2021.5) assigns MK type “A0 II n (shell)”</p> <p>¶ has been classified as a λ Boo star, apparently in error</p> <p>¶ IR excess indicates debris disk (possibly also detected in polarimetry), at average separation 155 AU; and yet a companion is also asserted, surprisingly present within this radius</p>
α	Tel	18 28.6 -45 57	3.49 [†]	-0.18	B3 IV [†]	12	-1.2	280	0.056	198	0 V?	<p>http://stars.astro.illinois.edu/sow/alphatel.html remarks that MK luminosity class IV notwithstanding, α Tel is still on the astrophysical (as opposed to the MK-phenomenological) main sequence (in other words, is still fusing core hydrogen)</p> <p>¶ said in 2005ApJS...158..193J to be among the (rare) He-rich stars; these authors list α Tel as a candidate-and-unconfirmed β Cep variable, and say they suspect it is a variable in the slowly pulsating B-star class; although α Tel has <i>HIPPARCOS</i> microvariability (0.909 d), it is absent from the AAVSO(VSX) database as at 28 Jan. 2021</p> <p>[THIS STAR ONLY IN ONLINE VERSION OF TABLE]</p>
λ	Sgr A	18 29.3 -25 24	2.82	1.02	K1 IIIb	~41.7	0.9	78	0.191	194	-43 V?	<p>Kaus Borealis[†]</p> <p>modest X-ray emission indicates some magnetic activity (not usual in a duly evolved, stable core-He-fusing, HR diagram “clump star”</p> <p>¶ lunar occultations are possible, planetary occultations possible yet rare; most recent planetary occultation was by Venus, on 1984 Nov. 19</p> <p>¶ unusual in occupying fully three roles in the Western pictorial traditions: as northernmost star of the Archer’s Bow, as westernmost (handle-tip) star of the Little Milk Dipper, and as uppermost (lid-knob) star in the Teapot</p>
α	Lyr A	18 37.7 +38 48	0.03 [†]	0.00	A0 Va [†]	130	0.6	25.0	0.350	35	-14 V	<p>Vega</p> <p>pole-on rapid rotator with circumstellar disk</p> <p>¶ pole-on rotators are useful for astroseismology, since all but the axisymmetric modes (whether radially symmetric or radially not symmetric) are helpfully rendered invisible to photometry (and in a rather analogous way, equator-on rotators are also useful through suppression of some modes); α Lyr pole-on orientation represents an extreme on a continuum whose other extreme is represented by the equator-on rapid rotator α Leo A (Regulus); the α Lyr A rapid rotation (2015A&A...577A..64B) now confidently asserts 0.68 d) yields oblate spheroid shape (here as with α Leo A); it is this, with consequent latitude-varying photosphere (severe temperature and luminosity gradients along</p>

the arcs of photospheric longitude,
 with equator coolest and darkest),
 rather than any evolution
 beyond core-hydrogen-fusion stage, that
 explains the anomalously
 high luminosity (α Lyr A
 is in MK luminosity class Va,
 rather than in the slightly dimmer V class
 that would be observed if
 its orientation was equator-on)
 ¶ α Lyr A is now known to harbour
 all three of the classical
 circumstellar-dust regimes (~1500 K, near-IR;
 ~120-170 K, mid-IR, as an
 analogue of our own zodiacal dust; and
 ~50 K, far-IR, as an analogue
 of our own Kuiper Belt:
 for regimes overview without
 specific reference to α Lyr,
 cf [2013ApJ...763..118S](#), section 1):
[2013ApJ...763..118S](#) is the paper
 announcing discovery of the second
 of these around α Lyr
 (with sections 5.1 and 5.2,
 respectively, summarizing
 previous α Lyr A work on the
 first and third of the three regimes):
 a question of recent interest
 is the origin of the α Lyr A
 exozodiacal (warm-regime, mid-IR)
 dust (episode analogous to our own
 planetary system's Late Heavy
 Bombardment? or, rather,
 some steady-state replenishment mechanism?);
 efforts at detecting exoplanet(s)
 to account for the complex
 inferred, and indeed in some
 wavelengths also now directly imaged,
 disk structure have not yet succeeded
 ¶ [2007ASPC...364...305G](#),
 reviewing the history of α Lyr A
 photometry, considers modest variability likely,
 the historical use of α Lyr A as a photometric
 standard notwithstanding
 (and indeed α Lyr A
 is described at AAVSO(VSX) as
 a low-amplitude δ Sct variable,
 in the now-obsolete AAVSO(VSX) "DSCTC"
 classification bin, with range
 -0.02-0.07 in V, with period 0.19 d)
 ¶ [2010A&A...523A..41P](#),
 with [2014A&A...568C...2P](#) corrigendum,
 is a recent discussion of α Lyr A magnetism
 (the authors note that α Lyr A
 "may well be the first confirmed
 member of a much larger, as yet
 unexplored, class of weakly-magnetic
 stars now investigatable with
 the current generation of stellar
 spectropolarimeters"; for origin, they
 somewhat favour dynamo over fossil,
 and radiative dynamo over core dynamo):
 consistently with magnetism,
[2015A&A...577A..64B](#) finds,
 via line-profile variations,
 multiple (bright, not dark) star spots,
 in some undetermined
 complex pattern (authors comment
 that this is "first strong
 evidence that standard A-type
 stars can show surface structure");
[2015A&A...577A..64B](#) is
 additionally one of several papers
 summarizing recent work on an
 interrelated complex of α Lyr A
 themes, comprising (in addition
 to magnetism) rotation, spots,
 photovariability, and pulsation
 ¶ in [2010ApJ...725.2401E](#), Fig. 8 with its accompanying
 discussion summarizes studies on elemental abundances
 (important because α Lyr A, as a rather
 "normal" star for MK temperature type A,
 might serve as a benchmark for

ϕ	Sgr [†]	18 47.0 -26 58	3.17 -0.11	B8 III	14	-1.2 240	0.051	89	+22	<p>appraising chemically peculiar A stars) ¶ E(B-V)=0.00 apparent duplicity now discounted (erroneous lunar-occultation observation) ¶ http://stars.astro.illinois.edu/sow/phisgr.html discusses some difficulties in physical modelling (if pole-on rotator, then there will be troublesome temperature and luminosity gradients along the arcs of photospheric longitude)</p>	
β	Lyr Aa1 [†]	18 50.9 +33 23	3.52v [†] 0.00	B7 Vpe (shell) [†]	~3.4 [†]	-3.8 ~960 [†]	0.004	152	-19 SB	<p>eclipsing:: 3.30-4.35, 13 d in V period is increasing at constant rate of ~19 s/y; orbit is seen nearly edge-on; prototype of the β Lyr class of eclipsing systems (but has also been assigned to the new class of “W Ser stars”: 1980AJUS...88..251P); AAVSO supplies information both via VSX database (showing, e.g. the high-precision recent determinations of period) and via www.aavso.org/vsots_betaLyr (a detailed astrophysics discussion, with bibliography): alternating deep and shallow visible-light minima, with the object eclipsed in the deep minima (the “donor”) a Roche-lobe-filling giant, currently ~3 M_⊙ and diminishing, and the object eclipsed in the shallow minima (the “gainer”) embedded in a thick accretion disk, currently ~13 M_⊙ and increasing; mass transfer is copious (~2e-5 M_⊙/y); this disk renders the gainer dim, and its eclipses consequently shallow, even though the (presently dim) gainer is (now, at this rather late stage in mass transfer) already ~4 times more massive than the (bright) donor (cf 1963ApJ...138..342H); further, instabilities in the accretion disk, from which ~20% of the light comes, make the light curve liable to vary slightly from cycle to cycle; the presently dim gainer is destined to be first (1) brightening, and spun up by conservation of angular momentum, as its obscuring accretion disk disappears by being dumped down into photosphere, and then (2) to become a slower rotator, tidally locked with the secondary; at stage “(1)”, the system will be a so-called “Rapidly Rotating Algol,” at stage “(2)”, on the other hand, the system will be simply a “classical Algol” ¶ 2008ApJ...684L..95Z presents the first (CHARA-interferometric) binary-resolving imaging, achieving resolution ~0.5 mas or ~0.7 mas (and for the first time in astrophysics deduces a β Lyr astrometric orbit); the bright low-mass donor, and the presently dim high-mass gainer, are evident, corroborating the overall conception of 1963ApJ...138..342H ; 2008ApJ...684L..95Z discusses also polar outflow jets on the gainer (these do not alter the essential situation: for the gainer, equatorial gain exceeds polar loss), and deduces a distance to ±15% (a distance consistent-to-within-uncertainties with the <i>HIPPARCOS</i> distance) ¶ 2012ApJ...750...59L discusses possible hot spot at edge of accretion disk, on the basis of spectropolarimetry (and 2013MNRAS.432..799M has modelling that provides for hot spot, and additionally for a bright spot, on the accretion disk) ¶ some observations have been made in radio and (a regime especially relevant to hot-spot studies) X-ray ¶ strictly speaking, this is a hierarchical system, Ab experiencing the binary that is Aa1+Aa2 as essentially a point mass; for the Aa+Ab pairing, and for possibility of further pairings (AB, AC, ... , Be, ...), cf WDS and (a source that reports inter alia <i>Gaia</i>) https://en.wikipedia.org/wiki/Beta_Lyrae ¶ although we here, following Garrison, assign a rather straightforward spectral type, this should be taken only as a starting point: cf, eg., 2000A&A...353.1009B, which lists six systems of spectral lines, while repeating an old O. Struve warning that spectrum involves circumstellar matter ¶ Kaler comments in http://stars.astro.illinois.edu/sow/sheliak.html “one of the most confusing, heavily studied, and important stars of the nighttime sky” ¶ the rather long period, with the large magnitude swing,</p>	Sheliak

													and the readily discoverable difference in depths of the alternating minima, make this object a suitable binoculars-or-naked-eye photometry project (using γ Lyr A as a comparison) even from locations suffering rather frequent cloud [THIS STAR ONLY IN ONLINE VERSION OF TABLE]
σ	Sgr Aa	18 56.6 -26 16	2.05 -0.13	B3 IV	14	-2.2	230	0.056	164	-11V			Nunki fast rotator ¶ lunar occultations are possible, and planetary occultation possible-yet-rare (most recently Venus, 1981 Nov. 17) ¶ E(B-V)=+0.02
ξ^2	Sgr	18 59.0 -21 05	3.52 1.15	K1 III	9	-1.7	400	0.034	113	-20			occultations (at any rate lunar) are possible ¶ the angular proximity of ξ^1 Sgr is a mere line-of-sight coincidence [THIS STAR ONLY IN ONLINE VERSION OF TABLE]
γ	Lyr A	18 59.7 +32 43	3.25 -0.05	B9 II [†]	5	-3.1	600	0.003	290	-21 V [†]			Sulafat has been both asserted and denied to be SB ¶ 2001A&A...371.1078A reports many metals underabundant
ζ	Sgr AB [†]	19 04.0 -29 51	2.60 0.06	A2 IV-V + A4:V:	37	0.4	90	n.a.	n.a.	+22 SB			A: 3.2; B: 3.5, 0.5" (2019), orbit 21.1 y separation 10.6 AU min, 16.1 AU max, average 13.4 AU ¶ under IAU rules, "Ascella" designates ζ Sgr A, not ζ Sgr B ¶ http://stars.astro.illinois.edu/sow/ascella.html discusses uncertainty in masses, remarks that temperatures are not yet directly measured ¶ Sgr C (17.6" in 2013) is probably a mere optical companion
ζ	Aql A [†]	19 06.4 +13 54	2.99 0.01	A0 Vann	~39.3	1.0	83	0.096	184	-25 SB			Okab among the most rapidly rotating stars known (period 16 h) ¶ in the angular-proximity grouping ζ Aql A, B, C, D, E, B is considered a gravitationally bound companion of A (mag. 12; angular distance 7.20" in 2009; separation \geq 125 AU, period \geq 800 y); additionally, faint (mag. 16.20) E shares in the AB proper motion, and so is likely gravitationally bound ¶ 2008A&A...487.1041A reports near-IR excess around ζ Aql A, and suggests that an unseen close companion is a more likely source than a close-in hot debris disk
λ	Aql	19 07.4 -4 51	3.43 -0.10	B9 Vnp (kB7HeA0) [†] 26		0.5	120	0.093	192	-12 V [†]			possibly SB rapid rotator (< 21h) ¶ suspected chemically anomalous (metals-weak, in λ Boo class); Astron. Alm. (epoch 2021.5) assigns MK type "A0 IVp (wk 4481)"
τ	Sgr	19 08.3 -27 38	3.32 1.17	K1.5 IIIb [†]	27	0.5	120	0.255 [†]	191	+45 [†]			possibly SB high velocity relative to Sun suggests origin outside galactic thin disk; underabundance of metals is consistent with this conjecture ¶ slow rotator (\leq 270 d)
π	Sgr AB [†]	19 11.0 -20 59	2.88 [†] 0.38	F2 II-III + n.a.	6	-3.1	500	0.036	182	-10			triple system, with AB-C poorly measured Albaldah B is at angular distance 0.10" (1989) from A: PA: 152 $^\circ$ →179 $^\circ$, 1936→1989; separation \geq 13 AU, orbit \geq 15 y; C (mag. 6) was observed in 1936 and 1939 to be at angular distance ~0.3" or ~0.4" from AB (separation \geq 40 AU, orbit \geq 100 y), but seems not to have been more recently measured ¶ π Sgr A, π Sgr B are possibly each of mag. 3.6 ¶ under IAU rules, "Albaldah" designates π Sgr A, not π Sgr B ¶ in HR-diagram terms, π Sgr A lies on blue edge of IS, without being presently observed to pulsate ¶ lunar occultations of ABC are possible, planetary occultations possible yet rare (next by Venus, 2035 Feb. 17)
δ	Dra A	19 12.6 +67 42	3.07 0.99	G9 III	33.5	0.7	97	0.133	46	+25 V			Altair
δ	Aql Aa	19 26.6 +3 10	3.36 [†] 0.32	F2 IV [†]	64	2.4	51	0.268	72	-30 SB [†]			fast rotator (> 0.9 d) ¶ Astron. Alm. (epoch 2021.5) assigns MK type F2 IV-V ¶ binary; 1989AJ...98.686K , addressing the difficulty posed by fast-rotator line broadening, refines previously computed orbital elements for δ Aql Aa, δ Aql Ab spectroscopically, finding period 3.426 d ¶ http://stars.astro.illinois.edu/sow/deltaaql.html discusses points of uncertainty (incl. the just-mentioned binarity, and possible δ Sct variability; although in 2018 δ Aql was not in the AAVSO(VSX) database, it is now (viewed 2021 Jan. 28) listed as a variable of low amplitude, in the

β	Cyg Aa [†]	19 31.6 +28 00	3.36	1.09	K3 II [†]	10 [†]	-2.3	330 [†]	0.009	229	-24 V	<p>γ Dor type, 3.36–3.37 in V, 1.04524 d) B: 4.68, 35"; Aa, Ab, Ac \leq 0.3" (2019) B is of MK type B9.5 Ve ¶ if AB is true binary, orbit is possibly \geq 100 000 y; the competing mere-optical-companions thesis is argued by Bob King in <i>Sky & Telescope</i> 2016 Sep. 21; same conclusion is reached in 2018 by P. Plait at www.syfy.com/syfywire/long-standing-astronomical-mystery-solved-albireo-is-not-a-binary-star, on strength of fresh <i>Gaia</i> data (which yield for β Cyg B π = 8.4 mas \pm2%, implying distance for β Cyg B, to two significant figures, 390 ly; however, further analysis is needed, since astrometry of β Cyg A is potentially perturbed by the multiplicity of A (https://en.wikipedia.org/wiki/Albireo recaps literature, with some reference to recent interferometry) ¶ our values, for β Cyg A, of π = 10 mas (strictly, 9.5 mas \pm 6.0%), with D consequently computed to two significant figures as 330 ly, are taken uncritically from <i>Gaia</i> ~2018, rather than (as in our previous Handbook editions) from <i>HIPPARCOS</i>; we do not here attempt a critical investigation of uncertainties ¶ β Cyg B is a fast rotator (< 0.6 d), and consistently with this is in emission (as "Be", rather than plain "B": being very evolved, this star is not, however, an instance of the "Be phenomenon" as discussed in the final subsection of our accompanying essay) ¶ β Cyg B is itself a tight binary system, and so is strictly β Cyg Ba,Bb (companion of mag. 9.2, at angular distance 0.4"; high eccentricity, with average separation ~40 AU, period almost 100 y) ¶ the name "Albireo," colloquially associated with the AB pairing as visible in a small telescope, applies under IAU rules only to β Cyg Aa B: 6.4, F1 V; 2.8", PA:41°→215°, 1826→2017 orbit 780 y; separation 84 AU min, 230 AU max, 157 AU average, period 780 y ¶ δ Cyg A is a rapid rotator ¶ δ Cyg C is gravitationally bound to the AB pair: mag. 12, angular distance (2017) 62.5", PA (for AC): 66°→67°, 1913→2017 ¶ variability has been suspected both in A and in B ¶ E(B-V)=+0.05</p>	Albireo
δ	Cyg A [†]	19 45.6 +45 11	2.89 [†]	0.00	B9.5 III	20	-0.7	160	-0.066	~42	-20 SB	<p>B: 6.4, F1 V; 2.8", PA:41°→215°, 1826→2017 orbit 780 y; separation 84 AU min, 230 AU max, 157 AU average, period 780 y ¶ δ Cyg A is a rapid rotator ¶ δ Cyg C is gravitationally bound to the AB pair: mag. 12, angular distance (2017) 62.5", PA (for AC): 66°→67°, 1913→2017 ¶ variability has been suspected both in A and in B ¶ E(B-V)=+0.05</p>	Fawaris
γ	Aql A	19 47.3 +10 40	2.72 [†]	1.51	K3 II [†]	~8.3	-2.7	390	0.017	100	-2 V	<p>radius ~0.5 AU ¶ variability has been asserted ¶ a rare instance of a "hybrid" star (possessing a (hot) corona, like our Sun's, and yet also emitting the cool high-mass wind typical in an evolved star)</p>	Tarazed
α	Aql A	19 51.8 +8 56	0.76 [†]	0.22	A7 Vnn	195	2.2	16.7	0.660	54	-26	<p>rapid rotator (~7 h or ~8 h, latitude-dependent) the first MS star, other than the Sun, to yield a measurement of photospheric oblateness (2001ApJ...559.1155V); 2007Sci...317.352M announces CHARA imaging with angular resolution ~0.65 mas (the first direct imaging of any MS star other than the Sun; http://news.bbc.co.uk/2/hi/science/nature/6709345.stm is a news writeup; 2007Sci...317.342M shows oblate rotation-flattened photosphere, brighter at poles than at equator) ¶ found in 2005ApJ...619.1072B, via <i>WIRE</i> salvage mission, to be a δ Sct variable (making this the brightest δ Sct variable, a classification now followed by AAVSO(VSX); second-brightest is β Cas); the 2005ApJ...619.1072B authors suggest that many δ Sct variables, as residents of the IS, may be oscillating at such low amplitudes as to evade detection except by such sensitive facilities as <i>WIRE</i> (their suggestion helps relieve a longstanding astrophysical puzzlement over IS residents that appear, inexplicably, not to be pulsating) ¶ drawing on interferometry, spectroscopy, and the 2005ApJ...619.1072B δ Sct astroseismology, 2020A&A...633A..78B, while conceding a failure of uniqueness, and consequently conceding the need for further spectroscopy, offers a physical model that</p>	Altair

η	Aql A [†]	19 53.6 +1 04	3.87 ^{v†}	0.63	F6–G1 Ib	2	–4.3	1000	0.011	140	–15 SB	<p>takes account of the rapid rotation (by assuming mere cylindrical symmetry, and not the outright spherical symmetry that would be appropriate in the modelling of a slow rotator); Table 5 of the paper summarizes its results, comparing them against earlier modelling; the paper finds a typical core rotation period ~0.6 of the rotation period of the photosphere, and with only modest latitude variation (shearing) in the rotation period in the photosphere (with middle altitudes ~7.7 h, equator ~7.8 h, immediate vicinity of poles ~8.1 h); the paper deduces a value for core metallicity that makes α Aql A young, aged only ~100 My (but some other recent literature proposes instead ~1.2 Gy; both suggested ages are consistent with the failure of α Aql A to have progressed significantly off the MS); the paper ascribes to α Aql A a remarkable variation in envelope temperature, with the envelope convective (because cooler) at low latitudes and radiative (because hotter) at high latitudes (a similar latitude-governed bifurcation in envelope characteristics is believed present in the rapid rotator α Cep A (Alderamin)); consistently with this latitude-dependent temperature variation, 2009A&A...497.511R finds modest coronal X-ray emission, attributed to modest dynamo activity at the low or intermediate latitudes (the authors note that of the stars not in a tight binary system, α Aql A is among the hottest known to have coronal X-ray emission) ¶ 2017A&A...608A.113N reports time-varying IR (K-band) excess, suggestive of tenuous circumstellar material (possibly debris disk: the “Be phenomenon”, present in many hot, young rapid rotators, is believed to involve a gas disk rather than a debris disk) ¶ α Aql B is mag. 9.8, at a wide 196” from A; under IAU rules, the name “Altair” designates just α Aql A</p> <p>Cepheid variable: 3.49–4.30 in V, 7.2 d more precisely, AAVSO(VSX) as at 2021 Jan. 28 gives 7.17679 d (same value as given in January 2019); BSC5 asserts 7.176641 d with period changes; 2002ApJ...140..465B (in centre panel of the author’s Fig 1) gives (1990s?) photometry (to tighter than ± 10 millimag), colour, and radial-velocity curves ¶ hot companion resolved with <i>HST</i> WFC3 (cf 2013AJ...146...93E: the authors, combining this WFC3 work with other work, conclude that η Aql is a triple; their hot-companion binarity result is astrophysically important, as supporting the quest for Cepheid masses, and so ultimately supporting the study of the (astrophysically crucial) Cepheid period-luminosity relation; WDS reports resolution as A, B, C, with B mag. –9, separation 0.7” in 2012 ¶ in the case of novice Northern Hemisphere observers troubled by frequent cloud, its rather long period makes η Aql A a better high-amplitude Cepheid demonstration than the more celebrated δ Cep A [THIS STAR ONLY IN ONLINE VERSION OF TABLE]</p>
γ	Sge	19 59.7 +19 33	3.51 [†]	1.57	M0 III [†]	13	–1.0	260	0.070	71	–33 V?	<p>radius 0.26 AU (from interferometry; the disk subtends an angle of 6.18 mas) ¶ Astron. Alm. (epoch 2021.5) assigns MK type M0–III ¶ slightly variable; already has a dead carbon core, is not yet a Mira [THIS STAR ONLY IN ONLINE VERSION OF TABLE]</p>
θ	Aql Aa	20 12.4 –0 45	3.24	–0.07	B9.5 III [†]	11	–1.5	290	0.036	81	–27 SB2 [†]	<p>a good marker of celestial equator θ Aql Aa, Ab SB 17.1 d, separation ~0.26 AU; Aa, Ab angular distance < 0.1”, Aa, Ac angular distance 1”; 1995AJ...110..376H gives orbital parameters, from interferometry ¶ θ Aql A pairing is metal-rich ¶ Astron. Alm. (epoch 2021.5) assigns MK type B9.5 III+ hierarchical quintuplet (or greater) Dabih https://en.wikipedia.org/wiki/Beta_Capricorni has a diagram summarizing the known gravitationally bound hierarchy: Aa, Ab1 (seen), Ab2 (unseen), Ba, Bb, where Aa is mag. 3.1, Ab1Ab2 is mag. 4.9, Ba is mag. 6.2, Bb is mag. 9.1 (but Wikipedia needs a caveat: since Ab is not yet resolved, even in interferometry, the designations “Ab1”, “Ab2” are not yet WDS-conformant); WDS also lists, as nearby in angular distance,</p>
β	Cap Aa [†]	20 22.2 –14 43	3.05	0.79	K0: II:†	10	–2.0	300	0.046	81	–19 SB	

												<p>C (mag. 8.8, 226"), D (mag. 13.0, 116"), and E (mag. 14.4, 3.9" from D): Ab1, Ab2 period is 8.7 d; Aa experiences Ab1Ab2 as essentially a point mass, recently at angular distance 50 mas (period 3.77 y, separation ~4 AU); Ba, Bb 0.5", according to WDS (and yet https://en.wikipedia.org/wiki/Beta_Capricorni states 3"), PA: 106°→54°, 1884→2019; AB 205", PA: 268°→267°, 1800→2012; each of AaAb, BaBb experiences the other as essentially a point mass, at separation ≥ 0.34 ly, with the AaAb+BaBb orbit ≥ 700,000 y ¶ spectral type of β Cap A is controverted; entire system appears in spectrograph as K0: II: + A5: V:n ¶ β Cap A is overabundant in Hg, Mn, and several other heavy elements ¶ lunar occultations are possible, planetary occultations possible-yet-rare</p>	
γ	Cyg A [†]	20 23.0 +40 20	2.23 [†]	0.67	F8 Ib [†]	2	-6.5	2000	0.003	111	-8 V	<p>BC combined light is mag. 11, with B, C mags 10.0, 11.0, respectively; A, BC angular distance 147" in 2010, with PA unchanged since 1877; however, http://stars.astro.illinois.edu/sow/sadr.html considers the A+BC pairing to be a mere line-of-sight coincidence (and WDS gives the following for BC: 1.9" in 2015, PA: 305°→302°, 1878→2015) ¶ unusual in being not only a supergiant, but a supergiant in MK type F (among supergiants, it is the hotter and the cooler types that are more usually encountered); γ Cyg A resides near the HR diagram Instability Strip: 2010AJ...140.1329G first surveys the observational literature, then discusses spectral variations (possibly pulsation-style oscillation, or alternatively large convection cells are possible; and indeed convection cells can be a driver of oscillation) ¶ radius ~1 AU (http://stars.astro.illinois.edu/sow/sadr.html discusses uncertainty) ¶ BSC5: "no demonstrable connection" between γ Cyg and the so-called γ Cyg supernova remnant</p>	Sadr
α	Pav A	20 27.3 -56 40	1.94	-0.12	B2.5 V	18	-1.8	180	0.086	175	+2 SB [†]	<p>SB 11.753 d, separation 0.21 AU ¶ 1988A&A...201..273V discusses galactic-astronomy implications of this star's puzzling deuterium paucity ¶ E(B-V)=+0.02 ¶ the name, although anomalously English, is nevertheless IAU-official: its origins lie in 1930s RAF Air Almanac project, which directed HM Nautical Almanac Office that no air-navigation star was to be left nameless</p>	Peacock [†]
α	Ind A	20 39.1 -47 13	3.11	1.00	K0 III CN-1 [†]	33	0.7	98	0.083	37	-1	<p>Fe overabundant (α Ind born in metal-rich ISM cloud?)</p>	
α	Cyg A	20 42.2 +45 21	1.25 [†]	0.09 [†]	A2 Ia	2 [†]	-6.9~1400 [†]		0.003	47	-5 SB	<p>blue supergiant, of radius ~0.5 AU or ~1 AU for context pertaining to this particular BSG in the general population of hypergiants and supergiants, cf https://en.wikipedia.org/wiki/List_of_largest_stars (which adopts "~1 AU"); for current state of theoretical investigations into BSG populations (crossing Hertzsprung-Russell diagram for the first time, redward? or, rather, after episode of mass loss, crossing for the second time, blueward?) cf, e.g. 2014MNRAS.439L...6G ¶ the prototype of the α Cyg variables: AAVSO(VSX) gives V ranges 1.21-1.29; seemingly irregular (in the α Cyg variables, many short-period oscillations are superimposed); 2011AJ...141...17R discusses α Cyg, reporting a 1977-through-2001 campaign in both photometric and spectroscopic variability ¶ α Cyg core hydrogen-fusion career started in MK spectral type B, or possibly even in the rare MK spectral type O ¶ present mass-loss rate is ~8e-7 M_o/y ¶ slow rotator (period possibly as long as 0.5 y, consistently with its large radius and its ongoing mass loss)</p>	Deneb

												¶ public-outreach astro audiences enjoy comparing and contrasting distance, and therefore intrinsic luminosity, of α Cyg with distance, and therefore intrinsic luminosity, of the other two Summer Triangle stars (nearby α Lyr, nearby α Aql; all three are similar not only in their apparent magnitudes, but also in falling within MK type A, and consequently in lacking tint, even through binoculars); it is perhaps worth stressing in such lectures that the α Cyg distance, although large (1500 ly? more?), is nevertheless not yet well known; Kaler in http://stars.astro.illinois.edu/sow/deneb.html , accepting ~1500 ly, writes that if placed at distance of α Lyr, α Cyg “would /.../ be as bright as a well-developed crescent Moon, cast shadows on the ground, and easily be visible in broad daylight”
η	Cep A	20 45.7 +61 55	3.41	0.91	K0 IV [†]	70.1	2.6	46.5	0.823 [†]	6	-87 [†]	high velocity relative to Sun indicates interloper status in galactic thin disk (and observed underabundance of Fe is consistent with interloper status)
β	Pav	20 46.9 -66 07	3.42	0.16	A6 IV [†]	~24.1	0.3	135	0.044	283	+10	still a fast rotator (≤ 2.3 d), although core hydrogen fusion is ended or is close to ending
ε	Cyg Aa [†]	20 47.1 +34 03	2.48	1.02	K0 III	44.9	0.7	73	0.486 [†]	47	-11 [†] SB [†]	¶ Astron. Alm. (epoch 2021.5) assigns MK type A6 IV- Aljanah C: common proper motion, 79" (2017) AC PA: 266°→269°, 1959 →2017; AC orbit $\geq 50,000$ y, separation ≥ 1700 AU (where C is a red dwarf, mag. 13.4); the SB pairing (with just one set of lines visible) ε Cyg Aa+Ab has period ≥ 15 y ¶ velocity of AaAb+C relative to Sun is high
ζ	Cyg Aa [†]	21 13.9 +30 19	3.21	0.99	G8 IIIa Ba [†] 0.5	23	0.0	140	0.069	175	+17 SB	in evolutionary terms, possibly a Red Clump resident (stable helium fusion in core); but it might also be the case that core helium fusion has yet to begin ¶ chemically a mild barium star (Astron. Alm. (epoch 2021.5) assigns MK type “G8 ⁺ III-IIIa Ba 0.5”; 1992Obs...112..168G discusses spectroscopy, reviewing history at a level of detail so instructive as to make this a case study for spectroscopy technique more generally, even outside the particular domain of ζ Cyg); consistently with this chemical anomaly, ζ Cyg A has WD companion ζ Cyg B (before becoming a WD, this close companion deposited barium onto ζ Cyg A as it shed mass: WD orbit 17.8 y, separation 8 AU min, 13 AU max, 11 AU average; 2001MNRAS.322..891B announces direct imaging with <i>HST</i> WFPC2 (elongated smear, WD partly resolved, possibly 36 mas))
α	Cep A	21 19.1 +62 41	2.45 [†]	0.26	A7 Van [†]	66.5	1.6	49.1	0.158	72	-10 V	Alderamin fast rotator (< 12 h); the rotational shape distortion, into an oblate spheroid, gives α Cep A a remarkable variation in envelope temperature, with the envelope convective (photosphere ~6600 K) at equator and radiative (photosphere ~8600K) at poles (the transition temperature is ~8300 K): a similar latitude-governed bifurcation in envelope characteristics is present (cf 2011ApJ...732...68C Fig. 9) in the rapid rotator α Aql A (Altair) ¶ Astron. Alm. (epoch 2021.5) assigns MK type A7 V [†] n ¶ listed by AAVSO(VSX) as δ Sct variable, with V mag. range 2.41-2.47 ¶ several factors, including X-ray emission (consistent with corona, as might be expected for convection-harboring latitudes of the envelope) indicate magnetic activity
β	Cep Aa [†]	21 28.9 +70 39	3.23v [†]	-0.20	B1 III [†]	5	-3.4	700	0.015	56	-8 SB	Alfirk PA: 255°→251°, 1779 →2016; orbit $\geq 40,000$ y ¶ the archetype of the β Cep variables (although this same class is sometimes called the “ β CMA variables”), and (as is typical in the class) known to be multiperiodic: AAVSO supplies a 2010 Apr. 13 backgrounder at www.aavso.org/vsots_betacep ; AAVSO(VSX) as viewed 2021 Jan. 16 asserts period 0.1904881 d; AAVSO archives a notice for an August 2009 β Cep

												campaign (coordinated photometry, spectroscopy, CHARA) at www.aavso.org/aavso-special-notice-162	
												¶ β Cep Aa is a magnetic star	
												¶ system comprises at least (the much-studied variable) Aa and Ab (mag. 6.6, probably a Be-phenomenon star, and the origin of the Be-phenomenon behaviour observed in AaAb); Aa+Ab period 85 y (when resolved with speckle interferometry in 1972, angular distance was 250 mas); β Cep B is mag. 8.6, at angular distance 13.5" in 2016; if B is gravitationally bound to AaAb, then period is $\geq 40,000$ y, with separation 3,000 AU	
												¶ MK luminosity class III ("giant") notwithstanding, β Cep Aa is still fusing hydrogen in its core	
β	Aqr A	21 32.7 -5 29	2.9 [†]	0.83 [†]	G0 Ib [†]	6	-3.2	500	0.020	114	+7 V?		Sadalsuud
												a rare instance of a yellow supergiant; possibly now evolving blueward in a second crossing of the HR diagram	
												¶ spectroscopically a "hybrid" star, combining signature of hot corona with signature of cool massive wind; 2005ApJ...627L..53A , in a study jointly covering β Aqr A and the astrophysically similar hypergiant (likewise a hybrid) α Aqr A, reports <i>Chandra</i> observation of coronal X-rays (first X-ray detection from a hybrid G supergiant; such supergiants are X-ray deficient, their coronae notwithstanding)	
												¶ β Aqr lies in the HR diagram Instability Strip, and yet is not known to be a pulsator	
ε	Peg A	21 45.2 +9 58	2.38v [†]	1.52 [†]	K2 Ib [†]	5	-4.2	700	0.027	89	+5 V		Enif
												irregular var.: 2.37–2.45 in V (flare in 1972) 1972IAUC.2392....1W reports extreme flare-like brightening, ~10 minutes, to V mag. 0.7	
												¶ orange-class supergiant	
												¶ Astron. Alm. (epoch 2021.5) assigns MK type K2 Ib–II	
												¶ 1987MNRAS.226..563S discusses abundances, finding that, earlier literature notwithstanding, ε Peg A is unremarkable in its Ba (and unremarkable in its Sr), and therefore discounting an earlier suggestion that ε Peg A outer layers have hosted nucleosynthesis in slow-neutron capture	
												¶ BSC5 suggests "cooler shell surrounding" eclipsing binary: V 2.81–3.05, 1.0 d, 3.2 + 5.2 Deneb Algedi since SB is not resolved (even interferometrically), WDS is not yet able to write "Cap Aa", "Cap Ab"; the δ Cap A pair is classified at AAVSO(VSX) as an Algol-type eclipsing binary, 1.0227688 d (period current as of 2021 Jan. 16; AAVSO(VSX) also yields O-C, i.e. period-monitoring, plotting from 2016); secondary is ~3 mag. fainter than primary and is judged in 1992MNRAS.259..251W to be mildly active, possibly tidally locked, with large spot; A is known to be SB since 1906 (Slipher), and yet is known to be eclipsing only as of 1956PASP...68..541E	
δ	Cap A	21 48.2 -16 02	2.85v [†]	0.18	A3mF2 IV:†	84	2.5	38.7	0.396	139	-6 SB†		
												¶ lunar occultations are possible, planetary occultations possible-yet-rare	
												¶ 1994MNRAS.266L..13L rebuts earlier assertion of δ Sct variability, and remarks that "given the brightness of the system, δ Cap is poorly observed," with period awkward for any one solitary observatory (an implication of this remark is that coordinated intercontinental photometry would now be helpful)	
												¶ Astron. Alm. (epoch 2021.5) assigns MK temperature type F2m and does not assign an MK luminosity class	
γ	Gru	21 55.2 -37 16	3.00	-0.08	B8 IV–Vs	15	-1.1	210	0.099	98	-2 V?		Aldhanab
α	Aqr A	22 06.9 -0 13	2.95 [†]	0.97 [†]	G2 Ib [†]	6	-3.1	~520	0.021	117	+8 V?		Sadalmelik
												a good marker of celestial equator	
												a rare instance of a yellow supergiant; possibly now evolving blueward in a second crossing of the HR plane; resides in the IS (under at least one definition of IS) and yet is nonpulsating (cf further 2017AstL...43..265U)	
												¶ spectroscopically a "hybrid star," combining signature of hot corona with signature of cool, massive wind; 2005ApJ...627L..53A , in a study jointly covering α Aqr A and the astrophysically similar supergiant (likewise a hybrid star) β Aqr A reports <i>Chandra</i> observation of coronal X-rays	

α	Gru A	22 09.6 -46 51	1.73	-0.07	B7 Vn	32	-0.7	101	0.194	139	+12	(first X-ray detection from a hybrid G supergiant; such supergiants are X-ray deficient, their coronae notwithstanding)	Alnair
θ	Peg	22 11.3 +6 18	3.52 [†]	0.09	A2mA1 IV-V [†]	35	1.3	90	0.284	84	-6 SB2	rapid rotator (< 1d) ¶ E(B-V)=-0.02	Biham
ζ	Cep	22 11.6 +58 18	3.39 [†]	1.56 [†]	K1.5 Ib [†]	3.9	-3.7	800	0.014	69	-18 SB	rapid rotator (< 20 h); consistently with rapid rotation, and therefore with a stirred atmosphere, elemental abundances are unremarkable ¶ earlier assertion of δ Sct variability is now discounted [THIS STAR ONLY IN ONLINE VERSION OF TABLE]	
α	Tuc	22 20.0 -60 09	2.87	1.39	K3 III [†]	16	-1.1	200	0.081	241	+42 SB [†]	orange supergiant either approaching core helium fusion or already in core helium fusion ¶ an eclipsing companion has been suggested, with suggestion later questioned ¶ metals somewhat overabundant	
δ	Cep A [†]	22 30.0 +58 32	4.07 ^{v†}	0.78	F5-G2 Ib	4 [†]	-3.0	900 [†]	0.016	77	-15 SB [†]	SB 11.5 y, separation possibly 11.5 AU ¶ primary in the SB is a giant, with carbon underabundant, nitrogen overabundant ¶ http://stars.astro.illinois.edu/sow/alphatuc.html discusses uncertainties in the evolutionary stage of this giant, offering three scenarios the prototype Cepheid variable: 3.49-4.36 in V, 5.4 d second-nearest Cepheid (α UMi is still nearer) ¶ AAVSO offers a tutorial at www.eso.org/public/outreach/eduoff/aol/market/collaboration/varstar/pg2.html and an initial backgrounder at www.aavso.org/vsots_delcep ; the first three sections of a paper directed inter alia to AAVSO observers, 2016JAVSO..44..179N, constitute a deeper backgrounder on the Cepheids ¶ AAVSO(VSX) has, as viewed 2021 Jan. 28, period 5.366266 d (but is this value possibly now stale?); although Cepheids experience both period jitter and (monotonic) period slide, with a slide of even 200 s/y possible, 2014ApJ...794...80E finds δ Cep period sliding slowly, at just -0.1 s/y (period decrease-increase is a signature of evolution, specifically of density increase-decrease, as a Cepheid passes across the HR diagram (δ Cep is now making its second such passage, moving blueward)) ¶ 2015ApJ...804..144A announces that δ Cep is SB, with period 2201 d ¶ accurate distances to Cepheids are foundational in cosmology, which needs independently known (galactic) Cepheid distances before embarking on its external-galaxy distance deductions through applications of the Cepheid Period-Luminosity (PL) Law; it is reassuring that the 2007 HIPPARCOS distance and the distance implied by the usual PL calculation agree to within uncertainties; although we have here stated the 2007 HIPPARCOS parallax, on which distance of δ Cep depends, as 4 mas, our cited 2007 HIPPARCOS determination is more formally, with decimal fractions and the uncertainty made explicit, 3.77±0.16 mas; 2015ApJ...804..144A proposes instead 4.09±0.16 mas, with the remark that impending <i>Gaia</i> may be expected, in part in the light of these authors' SB announcement, to secure an authoritative parallax; an already reassuring state of affairs may thus be expected to improve further ¶ mass loss ~1e-6 M _o /y; bow shock in ISM has now been detected ¶ C: 6.1, 41" (2017), has period ≥ 300,000, separation ≥ 11,000 AU [THIS STAR ONLY IN ONLINE VERSION OF TABLE]	
ζ	Peg A	22 42.5 +10 57	3.41 [†]	-0.09	B8.5 III [†]	16	-0.6	210	0.078	98	+7 V?	our (Garrison) MK type notwithstanding, B8 V has been suggested ¶ fast rotator (< 1.4 d) ¶ microvariable (2007PASP..119.483G discusses satellite detection of amplitude ~0.5 millimag); assigned by AAVSO(VSX) to the class of "slowly pulsating B stars"	Homam

β	Gru	22 43.9 -46 46	2.07 ^{v†}	1.61 [†]	M5 III [†]	18	-1.6	180	0.135	92	+2	semreg. variable 1.90–2.3 in V, 37d among the rather uncommon cool red giants, with radius slightly > 0.8 AU ¶ Astron. Alm. (epoch 2021.5) assigns MK type M4.5 III ¶ classified at AAVSO(VSX) as semireg. late-type giant, perhaps on the basis of 2006JAVSO..34..156O (this paper might serve as a case study for effective amateur-budget intercontinental photometry collaboration)	Tiaki
η	Peg Aa [†]	22 44.0 +30 20	2.93 [†]	0.85	G8 II	15	-1.2	210	0.029	153	+4 SB	we give mag. of η Peg Aa+Ab with also some contribution from BC; Aa alone is mag. ~4, and Ab is mag. ~7, of MK type F0 V ¶ η Peg Aa+Ab period 813 d ¶ Aa is slow rotator (818 d?) ¶ system is possibly more than a binary: cf WDS, which lists, apart from Aa and Ab, also celestial-sphere neighbours B, C, D, E, F, G, H, I	Matar
ε	Gru	22 49.8 -51 12	3.49	0.08	A2 Va	25	0.5	130	0.126	121	0 V	rapid rotator (< 0.65 d) [THIS STAR ONLY IN ONLINE VERSION OF TABLE]	
ι	Cep	22 50.5 +66 19	3.50	1.05	K0 III [†]	28.3	0.8	115	0.141	208	-12	Astron. Alm. (epoch 2021.5) assigns MK type K0 III [THIS STAR ONLY IN ONLINE VERSION OF TABLE]	
μ	Peg	22 51.0 +24 43	3.51	0.93	G8 III [†]	31	0.9	106	0.151	106	+1	Astron. Alm. (epoch 2021.5) assigns MK type G8+ III [THIS STAR ONLY IN ONLINE VERSION OF TABLE]	Sadalbari
δ	Aqr	22 55.8 -15 42	3.27	0.07	A3 IV–V	20	-0.2	160	0.051	237	+18 V	weak $\lambda 4481$ ¶ rapid rotator (< 3.0 d)	Skat
α	PsA Aa [†]	22 58.8 -29 30 [†]	1.17	0.14	A3 Va [†]	130	1.7	25.1	0.368	1 17	+7	2008 (<i>HST</i>) image was debris cloud, not exoplnt <i>HST</i> putative 2008 “exoplanet” was IAU-named Dagon, after a Semitic deity; at ~125 AU, in the outermost of the debris rings; Dagon was in always-wide (albeit eccentric) orbit, making direct imaging, as opposed both to spectroscopy (for star Doppler wobble) and astrometry (for star transverse wobble) the tool of choice: 32 AU min, 320 AU max; period ~1700 y ¶ in more recent years, it was suggested that Dagon could be a mere dust cloud, or an aggregation of rubble, or a single rocky body; an explanation was needed for the fact that Dagon proved so readily <i>HST</i> -visible (e.g. visibility enhanced by circumplanetary dust sphere, or by circumplanetary ring system?); Dagon mass was uncertain (< 2× Jupiter, perhaps even ~Earth); but with Dagon now no longer <i>HST</i> -visible, it would appear that the 2008 <i>HST</i> image was of an expanding debris cloud, now become too tenuous for detection ¶ the nested circumstellar dust rings extend as far as radius ~150 AU (a distance recalling the Solar System Kuiper Belt); 2017ApJ...842...8M reports complete outer debris-ring mapping, via ALMA (230 GHz radio), finding ring mass of 0.015 Earths, eccentric, with α PsA A offset from the ring centroid ¶ α PsA A is a fast rotator (< 1d) ¶ in evolutionary terms, α PsA A is sufficiently young to be undergoing an analogue of the Solar System’s Late Heavy Bombardment (and consistently with this, 2017ApJ...842...9M says exocometary gas is detected, by ALMA 230 GHz radio) ¶ 2017ApJ...842...8M comments that “given its unique characteristics and architecture, the Fomalhaut system is a Rosetta stone for understanding the interaction between planetary systems and debris disks” ¶ α PsA A has low metallicity ¶ 2013AJ...146..154M , working both from proper motion (across the celestial sphere) and from velocities along the line of sight, concludes that α PsA, B, and C belong to the same system: B (a flare star) is V mag. 7.1, at angular distance almost 2° (period ≥ 7.6 My), while C is V mag. 13.2, at enormous angular distance 5.7° (and yet at a sufficiently low separation from AB to have the AB gravitational field dominate the general external gravitational field at its location; period ≥ 35 My) ¶ β Peg, α Peg serve as pointers: since α PsA lies a couple of arcminutes N of	Fomalhaut

β	Peg A	23 04.8 +28 12	2.44 v^{\dagger}	1.66	M2 II–III †	16.6	-1.5	~196	0.232	54	+9 V	DEC=-30°, α PsA rises (if briefly) above the horizon even for such Canadian subarctic communities as Churchill, and for such Scandinavian communities as Stavanger semiregular variable, mag. 2.31–2.74 in V, 43.3 d with period 43.3 d ¶ Astron. Alm. (epoch 2021.5) assigns MK type M2.5 II–III ¶ an intermediary between straightforward red giant and red bright giant (radius ~0.5 AU); mass-loss rate is notably low for such a star ($\leq 1e-8 M_{\odot}/y$; i.e. ~100 \times lower than mass loss rate of α Ori Aa (Betelgeuse); <i>IRAS</i> detected no IR excess)	Scheat
α	Peg	23 05.8 +15 19	2.49	0.00	A0 III–IV	24	-0.6	133	0.073	124	-4 SB	rapid rotator (1.5 d)	Markab
γ	Cep A † +IP †	23 40.2 +77 45	3.21	1.03	K1 III–IV †	71	2.5	46	0.135	339	-42 V?	B: 7.30, 0.9", PA: 257°→256°, 2006→2006 separation 12 AU min, 25 AU max, period 66 y or 67 y; 2007A&A...462..777N reports the first direct imaging of γ Cep B, by Subaru ¶ γ Cep A possible rotation period 781 d (making this star a slow rotator) ¶ exoplanet orbiting γ Cep A (IAU-named Tadmor) is among the few discovered in a binary system; it is circumstellar without being circumbinary: period 2.47 y, average separation 2.05 AU, mass between 3 \times Jupiter and 16 \times Jupiter ¶ Astron. Alm. (epoch 2021.5) assigns MK type K1 III–IV CN 1	Errai