The Insider's Guide to the Galaxy Presents...

Messier Minutes

a guide to completing RASC's Messier Observing Certificate



Part 10 -

May 24th - June 7th, 2022

The following pages include a list of objects discussed in the May 24th, 2022, episode of Messier Minutes. Including finder charts and log pages.

List of Targets Discussed:

May 24 moonless; this is last of the spring Messiers

Messier #	Constellation	Magnitude	Type	NGC	Name
63	CVn	8.6	SG	5055	Sunflower Galaxy
94	CVn	8.2	SG	4736	Cat's Eye or Croc's Eye
101	UMa	7.9	SG	5457	Pinwheel Galaxy
51	CVn	8.1	SG	5194	Whirlpool Galaxy & NGC 5195
102	Dra	9.9	SG	5866	Spindle or Fool's Gold

Notes:

M63:

(13x7 arc-minutes)

Flocculent spiral galaxy (patchy, discontinuous arms) is barely visible in binoculars. Bright core and oval halo can be seen in small aperture telescopes. Larger apertures will reveal structure and dust lanes. Bright foreground star in halo. Close to the line connecting Alkaid to Cor Corali, but much closer to the latter. Or, make a right angle set square towards Alkaid using the Chara – Cor Corali separation. Or, aim your finger a thumb's width north of mag 4.7 star 20 CVn. Watch for shape, structure, and asymmetry. 37 million I-y away.

M94:

(11x9 arc-minutes)

Stubby barred spiral galaxy. Bright core and halo allow sightings in large binoculars under dark skies. Readily visible in smaller aperture telescopes, where it looks like a planetary nebula. large apertures can reveal a bright ring around the core, curving dust lanes, and a faint outer ring composed of two curving arms. Located midway between Cor Corali and Chara, but offset a thumb's width towards Alkaid – a squat isosceles triangle. Watch for orientation, field stars, extent of the halo. 16 million l-y away.

M101:

(29x27 arc-minutes)

Large, bright, face-on loose spiral galaxy near the Big Dipper's handle. Relatively easy in binoculars under dark skies, it's no problem for any size of telescope. More details are seen with larger apertures. Make an isosceles triangle with Alkaid and Mizar on the Polaris side of the Big Dipper's handle. 21 million I-y away. Watch for asymmetry in the spiral arms (making it Arp 26) and small nearby NGC 5477 plus NGCs 5473, 544, and more.

M51:

(11x7 arc-minutes)

Grand design spiral galaxy with a well-defined core and prominent arms). Readily visible in good binoculars under dark skies, and any size of telescope, even from suburban skies. Larger apertures better show the arms and dark dust lanes, and the star bridge connecting it to NGC 5195. Located 3 degrees from Alkaid at a 90° angle from the Big Dipper's handle, in the direction of Cor Corali. Look for shape and structure, knots of H2 regions. Interaction between the two galaxies makes it Arp 85. 23 million I-y away.


M102:

(5x2 arc-minutes)


Edge-on lenticular galaxy. Small and fainter, so not a binocular target. Visible in small telescopes, but larger apertures will show its large halo and bisecting dust lane. Located 4° west of Edasich (lota Dra), on the Arcturus side. Also, a third of the way along the line joining Theta Dra and Alkaid. Or double the line from Kappa Dra to Thuban. Look for the shape and orientation, dust lane, and field stars. 50 million I-y away.

Target Finder Charts:


Overview of Targets -



M63, M94, and M51 Closer Look -




M101 & M102 Closer View -



Sunflower Galaxy

M		7.4	()						
Mi	essier Object		63						
	NGC	50	55						
(Constellation	Ca	ane	s V	⁷ en	ati	ci		
	Type	Sp	oira	ıl C	Fala	аху	(G-S	SAbc)	
	Magnitude	8.	6						
Distance (Kil	o light-years)	37	000	0					
	RA	13	15	.8					
	Dec	+4	12:0)2					
	Size	14	.0	x 8	.0'				
UM I	UM II	75	,76)				37	
	SA	7							
	Remarks	!!	Su	nfl)W	er (Galax	xy; bright,	
		ele	ong	ate	ed				
Ti	ime (hh:mm)								
	Seeing	1	2	3	4	5			
Γ	ransparency	1	2	3	4	5			
Observ	ing Location								
	Telescope								
Date ((dd:mm:yyyy)								



Notes				
	 _	_	_	


PN: Planetary Nebula	RN: (diffu
SNR: Supernova Remnant	EN: (diffus
GC: Globular Cluster	G-: Galaxy
OC: Open Cluster	F/RN: Diff

RN: (diffuse) Reflection Nebula EN: (diffuse) Emission Nebula G-: Galaxy, with Hubble type given E/RN: Diffuse emission and reflection Nebula Seeing: 1 = Best 5 = Poor Transparency: 1 = Best 5 = Poor Time: DD:MM:YYYY

Date: Specify Time Zone or UT

* = Number of stars in cluster ** p = Photographic Magnitude

3.50.4					
M94					
4736					
Canes Venatici					
Spiral Galaxy (G-SAab)					
8.2					
14500					
12 50.9					
+41:07					
13.0' x 11.0'					
75 37					
7					
very bright and comet-like					
1 2 3 4 5					
1 2 3 4 5					



Notes			
		 ·	

PN: Planetary Nebula	F
SNR: Supernova Remnant	F
GC: Globular Cluster	(
OC: Open Cluster	F

RN: (diffuse) Reflection Nebula EN: (diffuse) Emission Nebula G-: Galaxy, with Hubble type given E/RN: Diffuse emission and reflection Nebula Seeing: 1 = Best 5 = PoorTransparency: 1 = Best 5 = Poor Time: DD:MM:YYYY


Date: Specify Time Zone or UT

* = Number of stars in cluster ** p = Photographic Magnitude *** !! = Showpiece Object

http://www.rasc.ca

Pinwheel Galaxy

Messier Obje	ect M101
NG	GC 5457
Constellation	on Ursa Major
Туј	pe Spiral Galaxy (G-SABcd)
Magnitud	de 7.9
Distance (Kilo light-yea	ars) 27000
R	AA 14 03.2
De	ec +54:21
Siz	ze 26' x 26'
UM I UM	II 49 23
S	SA 2, 7
Remarl	ks !! Pinwheel Gallaxy; diffuse face-on
	spiral
Time (hh:mn	n)
Seeir	ng 1 2 3 4 5
Transparence	cy 1 2 3 4 5
Observing Location	on
Telescop	pe
Date (dd:mm:yyyy	y)



Notes			

PN: Planetary Nebula	
SNR: Supernova Remnant	
GC: Globular Cluster	
OC: Open Cluster	

RN: (diffuse) Reflection Nebula
EN: (diffuse) Emission Nebula
G-: Galaxy, with Hubble type given
E/RN: Diffuse emission and reflection Nebula


Seeing: 1 = Best 5 = Poor Transparency: 1 = Best 5 = Poor Time: DD:MM:YYYY

Date: Specify Time Zone or UT

* = Number of stars in cluster ** p = Photographic Magnitude

Whirlpool Galaxy

Messier Obj	ect	M5	51							
N	NGC									
Constellat	ion	Ca	nes	s V	en	ati	ci			
T	ype	Sp	ira	l G	ala	axy	(G -	SAbc)		
Magnitu	ıde	8.4								
Distance (Kilo light-ye	ears)	370	000)						
	RA	13	29.	9						
Ι	Dec	+4'	7:1	2						
S	ize	8.0	' X	7.0	0'					
UM I UM	1 II	76						37		
·	SA	7								
Rema	rks	!! \	Wh	irl	po	ol (Gala	xy; supe	rb in b	ig
		sco	pe							
Time (hh:m	nm)									
See	ing	1	2	3	4	5				
Transparer	ncy	1	2	3	4	5				
Observing Locat	ion									
Telesco										
Date (dd:mm:yy	уу)									



Notes				
	 _	_	_	

PN: Planetary Nebula	RN: (diffuse) R
SNR: Supernova Remnant	EN: (diffuse) E
GC: Globular Cluster	G-: Galaxy, wit
OC: Open Cluster	E/RN: Diffuse


RN: (diffuse) Reflection Nebula EN: (diffuse) Emission Nebula G-: Galaxy, with Hubble type given E/RN: Diffuse emission and reflection Nebula Seeing: 1 = Best 5 = Poor Transparency: 1 = Best 5 = Poor Time: DD:MM:YYYY

Date: Specify Time Zone or UT

* = Number of stars in cluster ** p = Photographic Magnitude

NGC 5866 Spindle Galaxy

1			400							
Messier Ob	ject	M	102	2?_						
N	IGC	5866?								
Constella	tion	Draco								
Т	ype	Spiral Galaxy (G-SA0)								
Magni	tude	9.9								
Distance (Kilo light-y	ears)	40	000)						
	RA	15	06	.5						
	Dec	+55:46								
,	Size	6.6' x 3.2'								
UM I U	M II	50						22		
	SA									
Rem	arks	or	is]	M 1	02	=]	M101	1? (loo	k for 59	07)
Time (hh:	mm)									
Sec	eing	1	2	3	4	5				
Transpare	ency	1	2	3	4	5				
Observing Loca	tion									
Telesc	cope									
Date (dd:mm:y	yyy)									



Notes					
	·		·	·	
			·	_	
	_		_	_	

PN: Planetary Nebula
SNR: Supernova Remnant
GC: Globular Cluster
OC: Open Cluster

RN: (diffuse) Reflection Nebula
EN: (diffuse) Emission Nebula
G-: Galaxy, with Hubble type given
E/RN: Diffuse emission and reflection Nebula

Seeing: 1 = Best 5 = Poor Transparency: 1 = Best 5 = Poor Time: DD:MM:YYYY

Date: Specify Time Zone or UT

* = Number of stars in cluster ** p = Photographic Magnitude